

Downregulation of Ca^{2+} and Mg^{2+} Transport Proteins in the Kidney Explains Tacrolimus (FK506)-Induced Hypercalciuria and Hypomagnesemia

TOM NIJENHUIS, JOOST G.J. HOENDEROP, and RENÉ J.M. BINDELS

Department of Physiology, Nijmegen Center for Molecular Life Sciences, University Medical Center Nijmegen, Nijmegen, the Netherlands

Abstract. FK506 (tacrolimus) and dexamethasone are potent immunosuppressants known to induce significant side effects on mineral homeostasis, including hypercalciuria and hypomagnesemia. However, the underlying molecular mechanisms remain unknown. The present study investigated the effects of FK506 and dexamethasone on the expression of proteins involved in active Ca^{2+} reabsorption: the epithelial Ca^{2+} channel TRPV5 and the cytosolic Ca^{2+} -binding protein calbindin- $\text{D}_{28\text{K}}$. In addition, the renal expression of the putative Mg^{2+} channel TRPM6, suggested to be involved in transcellular Mg^{2+} reabsorption, was determined. Administration of FK506 to rats by daily oral gavage during 7 d significantly enhanced the urinary excretion of Ca^{2+} and Mg^{2+} and induced a significant hypomagnesemia. FK506 significantly decreased the renal mRNA expression of TRPV5 ($62 \pm 7\%$ relative to controls), calbindin- $\text{D}_{28\text{K}}$ ($9 \pm 1\%$), and TRPM6 ($52 \pm 8\%$), as determined by real-time quantitative PCR analysis. Further-

more, semiquantitative immunohistochemistry showed reduced renal protein abundance of TRPV5 ($24 \pm 5\%$) and calbindin- $\text{D}_{28\text{K}}$ ($29 \pm 4\%$), altogether suggesting that downregulation of these transport proteins is responsible for the FK506-induced Ca^{2+} and Mg^{2+} wasting. In contrast, dexamethasone significantly enhanced renal TRPV5 ($150 \pm 15\%$), calbindin- $\text{D}_{28\text{K}}$ ($177 \pm 23\%$), and TRPM6 ($156 \pm 20\%$) mRNA levels along with TRPV5 ($211 \pm 8\%$) and calbindin- $\text{D}_{28\text{K}}$ ($176 \pm 5\%$) protein abundance in the presence of significantly increased Ca^{2+} and Mg^{2+} excretion. This indicated that these proteins are directly or indirectly regulated by dexamethasone. In conclusion, FK506 and dexamethasone induce renal Ca^{2+} and Mg^{2+} wasting, albeit by different mechanisms. Downregulation of specific Ca^{2+} and Mg^{2+} transport proteins provides a molecular mechanism for FK506-induced hypercalciuria and hypomagnesemia, whereas dexamethasone positively regulates these proteins.

Immunosuppressants such as the calcineurin inhibitors cyclosporin A and FK506 (tacrolimus) along with glucocorticoids such as dexamethasone are widely prescribed in numerous disorders and to organ transplant recipients. Although their immunosuppressive actions are achieved by distinct mechanisms, FK506 and dexamethasone both are known to induce significant side effects on mineral homeostasis. These drugs are associated with an increased bone turnover, a negative Ca^{2+} balance and hypercalciuria, perturbations that can ultimately result in osteoporosis (1–3). Furthermore, hypomagnesemia is a widely known additional side effect of FK506 treatment (4, 5). The kidney is crucial to both Ca^{2+} and Mg^{2+} homeostasis by providing the main excretory route for these divalent ions. However, the exact mechanisms by which these

immunosuppressants provoke renal cation wasting are unknown.

The hypercalciuria during treatment with these drugs has been attributed to increased bone resorption as well as decreased renal Ca^{2+} reabsorption (2, 6, 7). In the kidney, the bulk of Ca^{2+} reabsorption is accomplished by Na^{+} -driven passive paracellular Ca^{2+} transport in the proximal tubule and thick ascending limb of Henle (TAL) (8). Fine-tuning of Ca^{2+} excretion is achieved by active transcellular reabsorption of Ca^{2+} in the distal convoluted tubule (DCT) and connecting tubule (CNT), which involves apical Ca^{2+} entry through the epithelial Ca^{2+} channel TRPV5 (previously ECaC1), intracellular buffering and facilitated diffusion by the Ca^{2+} -binding protein calbindin- $\text{D}_{28\text{K}}$, and basolateral extrusion by the $\text{Na}^{+}/\text{Ca}^{2+}$ exchanger (NCX1) and the plasma membrane Ca^{2+} ATPase (PMCA1b) (9, 10). Theoretically, inhibition of active Ca^{2+} reabsorption could be involved in the immunosuppressant-induced hypercalciuria.

FK506 treatment has been associated with an inappropriately high fractional excretion of Mg^{2+} , suggesting that inhibition of passive or active Mg^{2+} reabsorption could contribute to the hypomagnesemia (11). Active reabsorption of Mg^{2+} has been localized to the DCT, but in contrast to Ca^{2+} reabsorption, little is known about the specific proteins that mediate

Received June 13, 2003. Accepted November 27, 2003.

Correspondence to René J.M. Bindels, 160 Cell Physiology, University Medical Center Nijmegen, P.O. Box 9101, NL-6500 HB Nijmegen, the Netherlands. Phone: +31-24-3614211; Fax: +31-24-3616413; E-mail: r.bindels@ncmls.kun.nl

1046-6673/1503-0549

Journal of the American Society of Nephrology

Copyright © 2004 by the American Society of Nephrology

DOI: 10.1097/01.ASN.0000113318.56023.B6

transcellular Mg^{2+} transport (12, 13). Recently, it was shown that autosomal recessive hypomagnesemia, characterized by disturbed intestinal Mg^{2+} absorption and inappropriately high fractional Mg^{2+} excretion rates, is caused by mutations in the gene encoding TRPM6 (14, 15). Like other ion channels in the transient receptor potential (TRP) family, this protein contains six transmembrane domains and is highly homologous to the Mg^{2+} -permeable ion channel TRPM7, and TRPM6 mRNA expression was detected in kidney and intestine (15). Recently, we demonstrated that TRPM6 confers a Mg^{2+} permeable channel that is exclusively expressed in the apical membrane of mouse DCT and small intestine, suggesting that TRPM6 constitutes the apical Mg^{2+} entry channel in transcellular Mg^{2+} (re)absorption (16). The identification of TRPM6 provides an important tool to study renal active Mg^{2+} transport during FK506 treatment on a molecular level.

Taken together, the cascade of cellular and molecular events that lead to impaired renal Ca^{2+} and Mg^{2+} handling during FK506 and glucocorticoid treatment is largely unknown. Hypothetically, downregulation of Ca^{2+} and Mg^{2+} transport proteins in the distal part of the nephron may be involved in the pathogenesis of hypercalciuria and hypermagnesuria during drug treatment. The aim of the present study, therefore, was to determine the effects of FK506 and dexamethasone on the expression of the Ca^{2+} transport proteins TRPV5 and calbindin- D_{28K} and the putative Mg^{2+} channel TRPM6 in the kidney. To this end, rats were treated for 7 d with these immunosuppressants and housed in metabolic cages to collect samples for urine analysis. Subsequently, expression levels of TRPV5, calbindin- D_{28K} , and TRPM6 were determined by real-time quantitative PCR and immunohistochemical analysis.

Materials and Methods

Animal Studies

Young adult male Wistar rats with an initial weight of 225 to 250 g were housed individually in metabolic cages to collect 24-h urine samples. The animals were kept in a light- and temperature-controlled room with *ad libitum* access to standard pelleted diet and water. The animals were randomly assigned to either the control group ($n = 10$) or the two treatment groups, receiving FK506 ($n = 6$) or dexamethasone ($n = 6$). These immunosuppressants were dissolved in peanut oil and administered daily by oral gavage. Control animals received vehicle only, whereas the two treatment groups received either FK506 (Fujisawa Pharmaceutical Co., Osaka, Japan) at a single dose of 1 mg/d or dexamethasone (Sigma-Aldrich, Zwijndrecht, the Netherlands) at 300 μ g/d. Animals were treated for 7 d, after which blood samples were taken and the animals were killed. Kidney cortex and duodenum were sampled and immediately frozen in liquid nitrogen. In addition, kidney cortex was fixed for immunohistochemistry by immersion in 1% (wt/vol) periodate-lysine-paraformaldehyde for 2 h and 15% (wt/vol) sucrose in PBS overnight (17). Subsequently, samples were stored at $-80^{\circ}C$ until further processing. The animal ethics board of the University Medical Center Nijmegen approved all experimental procedures.

Analytical Procedures

Serum and urine Ca^{2+} concentrations were determined using a colorimetric assay as described previously (18). Serum and urine

[Mg^{2+}], [Na^{+}], [creatinine] [glucose], alkaline phosphatase, and γ -glutamyl transpeptidase were measured on a Hitachi autoanalyzer (Hitachi Corp., Tokyo, Japan).

Real-Time Quantitative PCR

Total RNA was extracted from kidney cortex and duodenum using Trizol Total RNA Isolation Reagent (Life Technologies BRL, Breda, the Netherlands). The obtained RNA was subjected to DNase treatment to prevent genomic DNA contamination. Thereafter, 2 μ g of RNA was reverse transcribed by Moloney-murine leukemia virus-reverse transcriptase (Life Technologies BRL) as described previously (19). The obtained cDNA was used to determine TRPV5, calbindin- D_{28K} , and TRPM6 mRNA levels in kidney cortex as well as TRPV6, calbindin- D_{9K} , and TRPM6 mRNA expression in duodenum. Expression levels were quantified by real-time quantitative PCR on an ABI Prism 7700 Sequence Detection System (PE Biosystems, Rotkreuz, Switzerland). PCR primers and fluorescence probes were designed using the computer program Primer Express (Applied Biosystems, Foster City, CA) and are listed in Table 1 (Biogelio, Malden, the Netherlands).

Immunohistochemistry

Immunohistochemical staining was performed as described previously (17). In short, either single or double staining of sections for TRPV5, calbindin- D_{28K} , kallikrein, and the Na^{+} - Cl^{-} co-transporter (NCC) was performed on 7- μ m sections of fixed frozen kidney samples. TRPV5 staining involved immersion of the kidney sections in boiled citrate target retrieval buffer (0.01 M sodium citrate and 0.01 M citric acid [pH 6.0]), which was then left to cool for 30 min and subsequent incubation in 0.3% (vol/vol) H_2O_2 in buffer (0.15 M NaCl, 0.1 M Tris-HCl [pH 7.5]) for 30 min. Sections were incubated for 16 h at $4^{\circ}C$ with primary antibody, affinity-purified guinea pig TRPV5 antibody (1:1000) (17), mouse anti-calbindin- D_{28K} (Swant, Bellinzona, Switzerland; 1:500), rabbit anti-NCC (1:300) (20), and rabbit anti-kallikrein (1:5000; Calbiochem, San Diego, CA). After incubation with biotin-coated goat anti-guinea pig secondary antibody, TRPV5 was visualized using a tyramide signal amplification kit (NEN Life Science Products, Zaventem, Belgium). For detection of calbindin- D_{28K} , NCC, and kallikrein, sections were incubated with an Alexa 488-conjugated goat anti-rabbit or an Alexa 594-conjugated goat anti-mouse secondary antibody. Images were made using a Zeiss fluorescence microscope equipped with a digital photo camera (Nikon DMX1200). For semiquantitative determination of protein levels, images were analyzed with the Image Pro Plus 4.1 image analysis software (Media Cybernetics, Silver Spring, MD), resulting in quantification of the protein levels as the mean of integrated optical density.

Statistical Analyses

Data are expressed as means \pm SEM. Statistical comparisons were tested by one-way ANOVA and Fisher multiple comparison. $P < 0.05$ was considered statistically significant. Statistical analysis was performed using the Statview Statistical Package software (Power PC version 4.51; Berkeley, CA) on a Macintosh computer.

Results

Body Weight, Urine, and Serum Analysis

Net urinary excretion of Ca^{2+} and Mg^{2+} is depicted in Figure 1. Additional biochemical analysis of the serum and 24-h urine samples of the experimental groups is shown in

Table 1. Sequences of primers and Taqman probes for real-time quantitative PCR^a

Gene	Forward Primer	Reverse Primer	Probe
TRPV5	5'-CTTACGGGTTGAACACCACCA-3'	5'-TTGAGAACACACAGAGCCCTTA-3'	5'-TGCTTCTCAGATAGTTGTTCTTGTACTTCTCCTCTTGT-3'
CaBP28	5'-GGAAGCTGGAGCTGACAGAGAT-3'	5'-TGAAGTCTTTCCACACACATTTTGAT-3'	5'-ACCAGTGCAGGAAAATTCCTTCTTAAATTCCA-3'
TRPM6	5'-AAAGCCATGCGAGTTATCAGC-3'	5'-CTTCACAATGAAAACCTGGCC-3'	5'-CCTGGTCTGAGATGATGTTCTCAAGCC-3'
TRPV6	5'-ATCCGCCGCTATGCACA-3'	5'-AGTTTTTCTCCTGAGTCTTTTTCCTCA-3'	5'-TTCCAGCAACAAGATGACCTTACTACTTGA-3'
CaBP9	5'-CCCAGAAATGAAGAGCATTTT-3'	5'-TTCTCCATCACCGTCTTATFCCA-3'	5'-CAAAAATATGCAGCCAAAGGAGCGCA-3'

^a PCR primers and fluorescent probes (5'FAM-3'TAMRA) were designed using the computer program Primer Express (Applied Biosystems) and purchased from Biologig. TRPV5, epithelial Ca²⁺ channel 1 (previously known as ECaC1); CaBP28, calbindin-D_{28K}; TRPM6, putative Mg²⁺ channel; TRPV6, epithelial Ca²⁺ channel 2 (previously known as ECaC2/CaT1); CaBP9, calbindin-D_{9K}.




Figure 1. Effect of FK506 and dexamethasone treatment on net urinary Ca²⁺ and Mg²⁺ excretion in the rat. Net excretion of Ca²⁺ and Mg²⁺ was determined by analysis of 24-h urine samples of rats individually housed in metabolic cages. Ctr, controls; FK506, FK506 1 mg/d by oral gavage; Dex, dexamethasone 300 µg/d by oral gavage. Data are presented as means ± SEM. **P < 0.05 versus controls.

Table 2. In the FK506 group, net and fractional excretion of Ca²⁺ and Mg²⁺ was significantly enhanced. Na⁺ excretion was reduced in this treatment group, whereas urine volume did not significantly differ from controls. Although serum Ca²⁺ and Na⁺ levels were unaffected, FK506 treatment induced a striking hypomagnesemia. Dexamethasone administration enhanced urinary Ca²⁺ as well as Mg²⁺ excretion. Likewise, Na⁺ excretion and urine volume were significantly increased. In the dexamethasone-treated animals, serum Ca²⁺ levels did not significantly differ from controls, whereas serum Na⁺ levels were decreased. Serum Mg²⁺ levels were significantly reduced relative to controls but remained within the normal range. Both serum creatinine and GFR, as determined by creatinine clearance, did not significantly differ from controls in both the FK506 and dexamethasone groups. Serum glucose levels were significantly increased in both the FK506- and dexamethasone-treated groups, resulting in increased glucose excretion in the dexamethasone group only. In addition, urinary excretion of markers of tubular damage (alkaline phosphatase and γ-glutamyl transpeptidase) was detectable, albeit not significantly increased in the FK506 group, whereas the dexamethasone-treated group showed significant enzymuria (21). Although baseline body weight did not differ significantly between the groups (data not shown), at the end of the experiment, dexamethasone-treated animals had a significantly lower mean body weight compared with controls, whereas FK506 treatment did not affect body weight.

Renal mRNA Expression Levels of TRPV5, Calbindin-D_{28K}, and TRPM6

For evaluating the possible association between increased urinary excretion of Ca²⁺ and Mg²⁺ and the expression levels of the respective transport proteins, renal mRNA levels of the epithelial Ca²⁺ channel TRPV5, the cytosolic Ca²⁺-binding protein calbindin-D_{28K} and the putative epithelial Mg²⁺ channel TRPM6 were determined by real-time quantitative PCR analysis. FK506 significantly decreased TRPV5, calbindin-

Table 2. Urine and serum analyses and body weight^a

	Controls	FK506	Dexamethasone
Urine			
FE Ca ²⁺ (%)	0.62 ± 0.07	1.08 ± 0.17 ^b	1.02 ± 0.11 ^b
FE Mg ²⁺ (%)	4.7 ± 1.1	35.1 ± 3.4 ^b	20.4 ± 2.2 ^b
Na ⁺ excretion (mmol/24 h)	1.6 ± 0.1	1.0 ± 0.2 ^b	2.4 ± 0.2 ^b
FE Na ⁺ (%)	0.64 ± 0.06	0.46 ± 0.04 ^b	1.15 ± 0.05 ^b
glucose excretion (mmol/24 h)	0.1 ± 0.1	2.7 ± 1.3	20.8 ± 4.8 ^b
diuresis (ml/24 h)	8.8 ± 1.0	18.6 ± 4.8	42.5 ± 6.2 ^b
C _{cr} (ml/min)	1.1 ± 0.1	1.1 ± 0.1	1.1 ± 0.1 ^b
ALP excretion (U/24 h)	0.1 ± 0.1	0.4 ± 0.2	1.1 ± 0.4 ^b
γ-GT excretion (U/24 h)	0.1 ± 0.1	1.3 ± 0.7	4.9 ± 2.5 ^b
Serum			
Ca ²⁺ (mmol/L)	2.53 ± 0.03	2.51 ± 0.03	2.49 ± 0.02 ^b
Mg ²⁺ (mmol/L)	1.01 ± 0.02	0.65 ± 0.02 ^b	0.92 ± 0.01 ^b
Na ⁺ (mmol/L)	142.1 ± 0.5	141.0 ± 0.4	134.0 ± 0.5 ^b
glucose (mmol/L)	6.9 ± 0.2	11.1 ± 0.9 ^b	18.7 ± 2.4 ^b
creatinine (μmol/L)	37 ± 1	38 ± 1	42 ± 2
Body weight (g)	310 ± 9.3	296 ± 6.1	277 ± 9.7 ^b

^a Controls, animals receiving vehicle only; FK506, FK506 1 mg/d; Dexamethasone, dexamethasone 300 μg/d; FE, fractional excretion; C_{cr}, creatinine clearance; ALP, alkaline phosphatase; γ-GT, gamma-glutamyl transpeptidase. Data are presented as means ± SEM.
^b *P* < 0.05 versus controls.




Figure 2. Effect of FK506 and dexamethasone on mRNA expression levels of Ca²⁺ and Mg²⁺ transport proteins in the rat kidney. Renal mRNA expression levels of the epithelial Ca²⁺ channel TRPV5, the cytosolic Ca²⁺-binding protein calbindin-D_{28K} (CaBP28), and the putative Mg²⁺ channel TRPM6 were determined by real-time quantitative PCR analysis. Ctr, controls; FK506, FK506 1 mg/d by oral gavage; Dex, dexamethasone 300 μg/d by oral gavage. Data are presented as means ± SEM. ***P* < 0.05 versus controls.

D_{28K}, and TRPM6 mRNA levels in the kidney (Figure 2). Conversely, dexamethasone treatment significantly enhanced the renal mRNA expression of both Ca²⁺ transport proteins and TRPM6.

Renal Protein Abundance of TRPV5 and Calbindin-D_{28K}

For assessing whether the changes in renal mRNA levels resulted in altered protein expression, TRPV5 and calbindin-D_{28K} protein abundance was semiquantified by immunohistochemistry (Figure 3A). These results illustrated that the above-mentioned mRNA results were indeed accompanied by similar effects on the protein level. Computerized analysis of the immunohistochemical staining showed significantly reduced TRPV5 and calbindin-D_{28K} protein expression during FK506

treatment, whereas dexamethasone significantly enhanced the abundance of these Ca²⁺ transport proteins in the kidney (Figure 3B). Of note, the effects on TRPV5 protein expression were more pronounced compared with the mRNA level, suggesting that in addition to affecting gene transcription, additional translational changes might have occurred.

For confirming the specificity of the observed effects on the Ca²⁺ and Mg²⁺ transport proteins, protein abundance of kallikrein and NCC were determined by immunohistochemistry (Figure 3A). The renal abundance of tissue kallikrein, a specific marker for the DCT and CNT (22), did not significantly differ from controls in both the FK506 and dexamethasone groups (Figure 3B). In addition, the protein expression of NCC, exclusively expressed in the DCT, was not significantly affected by FK506 treatment but showed a




Figure 3. Effect of FK506 and dexamethasone on protein abundance in the rat kidney. (A) Representative images showing kidney sections stained for the epithelial Ca²⁺ channel TRPV5, the cytosolic Ca²⁺-binding protein calbindin-D_{28K} (CaBP28), kallikrein, and the Na⁺-Cl⁻ cotransporter (NCC). (B) Protein abundance was determined by computerized analysis of immunohistochemical images and is presented as integrated optical density (IOD; arbitrary units). Ctr, controls; FK506, FK506 1 mg/d by oral gavage; Dex, dexamethasone 300 μg/d by oral gavage. Data are presented as means ± SEM. ***P* < 0.05 versus controls.

significant, albeit modest, increase in the dexamethasone-treated animals (Figure 3B). Furthermore, no signs of a general deleterious effect of FK506 (inclusion bodies, tubular vacuolization, and atrophy) were detected on light microscopic analysis (4). GFR was unaffected and enzymuria was not significantly increased, although markers of tubular damage were detectable. Importantly, the unaltered expression of kallikrein and NCC suggested that no overt FK506 nephrotoxicity was present in DCT.

In addition, co-staining of kidney sections for the presence of TRPV5 and kallikrein was performed, which showed a full co-localization of these proteins in DCT and CNT in the control animals (Figure 4A). Importantly, Figure 4B clearly illustrates that TRPV5 protein abundance in DCT and CNT is significantly reduced in the FK506-treated animals, whereas kallikrein expression levels are not affected. Dexamethasone treatment did not influence the co-localization of TRPV5 and kallikrein, substantiating that the dexamethasone-induced increase of TRPV5 abundance is confined to these nephron segments (Figure 4C). Co-staining of kidney sections for kallikrein and calbindin-D_{28K} showed similar results, with the exception that calbindin-D_{28K} expression extended further into the cortical collecting duct (data not shown).

Intestinal mRNA Expression Levels of TRPV6, Calbindin-D_{9K}, and TRPM6

Theoretically, upregulation of Ca²⁺ transport proteins in the intestine could contribute to the increased Ca²⁺ wasting, and decreased intestinal TRPM6 expression could contribute to the hypomagnesemia. Therefore, the duodenal mRNA expression of the epithelial Ca²⁺ channel TRPV6, the cytosolic Ca²⁺-binding protein calbindin-D_{9K}, and TRPM6 were determined (Figure 5). mRNA expression of all three transporters was not significantly altered by FK506 treatment. Dexamethasone significantly increased TRPV6 and calbindin-D_{9K} mRNA expression levels, whereas TRPM6 mRNA expression was not significantly affected.

Discussion

The present study demonstrated that FK506 and dexamethasone significantly enhance renal Ca²⁺ and Mg²⁺ excretion, albeit by different molecular mechanisms. The decreased expression of the Ca²⁺ transport proteins TRPV5 and calbindin-D_{28K} and the putative Mg²⁺ channel TRPM6 during FK506 treatment suggested that downregulation of these transport proteins contributes to the pathogenesis of FK506-induced hypercalciuria and hypomagnesemia. In contrast, dexametha-




Figure 4. Co-immunohistochemical staining of rat kidney for the epithelial Ca^{2+} channel TRPV5 and kallikrein. Co-staining of kidney sections for TRPV5 and kallikrein, a marker for the distal convoluted tubule (DCT) and connecting tubule (CNT), in controls (A), FK506-treated animals (B), and dexamethasone-treated animals (C). Controls, rats receiving vehicle only; FK506, rats receiving FK506 1 mg/d by oral gavage; Dexamethasone, rats receiving dexamethasone 300 $\mu\text{g}/\text{d}$ by oral gavage.

sone treatment increased renal expression levels of TRPV5, calbindin- $\text{D}_{28\text{K}}$, and TRPM6, indicating that these transport proteins are either directly or indirectly regulated by dexamethasone.

FK506 treatment significantly increased urinary Ca^{2+} excretion, accompanied by a downregulation of the renal mRNA expression of TRPV5 and calbindin- $\text{D}_{28\text{K}}$ and a specific reduction of the protein abundance of these Ca^{2+} transport proteins in DCT and CNT, recognized as the main sites of transcellular Ca^{2+} reabsorption. Previous reports showed reduced calbi-

ndin- $\text{D}_{28\text{K}}$ levels during FK506 treatment, suggesting that FK506 could affect active Ca^{2+} transport (23). That serum Ca^{2+} concentrations and GFR did not differ from controls confirmed that impaired Ca^{2+} reabsorption rather than an increased filtered load caused the hypercalciuria. Morphologic features of tubular toxicity were not detected. Furthermore, the excretion of urinary markers of tubular damage was not significantly increased and the expression of specific markers for DCT and CNT was unaltered, excluding that a general deleterious effect of FK506 on DCT is responsible for the observed downregulation. In addition, the expression of the major intestinal Ca^{2+} transport proteins was not increased by FK506 treatment, excluding that hypercalciuria is secondary to up-regulation of active Ca^{2+} absorption. The present data strongly supported our hypothesis that FK506 induces a primary defect of renal active Ca^{2+} reabsorption by specifically downregulating the proteins involved in active Ca^{2+} transport.

The molecular mechanism underlying the downregulation of the Ca^{2+} transport proteins by FK506 remains elusive. In previous studies, plasma calcitriol ($1,25(\text{OH})_2\text{D}_3$) was either unaltered or moderately increased, whereas plasma parathyroid levels were not affected by similar doses of FK506, which excludes that the reduced Ca^{2+} transport protein expression levels are secondary to decreased circulating levels of these calcitropic hormones (11, 23, 24). The immunosuppressive action of FK506 depends on the inhibition of the Ca^{2+} -dependent phosphatase calcineurin in T lymphocytes (25–27). Calcineurin is not known to be involved in renal Ca^{2+} reabsorption, but another calcineurin inhibitor, cyclosporine A, increased urinary Ca^{2+} excretion and decreased calbindin- $\text{D}_{28\text{K}}$ protein levels, suggesting that calcineurin inhibition may play a role in the impairment of Ca^{2+} reabsorption by these drugs (23, 28). In addition, FK506 binds to intracellular immunophilins called FK506-binding proteins (FKBP), which have been implicated as ion channel regulators (29–31). In particular, FKBP4 was shown to bind and regulate the Ca^{2+} -permeable *Drosophila* TRPL channel, and this binding was disrupted by the addition of FK506 (32). Furthermore, several intracellular Ca^{2+} -release channels were shown to be modulated by binding and dissociation of FKBP (33–35). Therefore, it is tempting to speculate that FKBP are potential associated proteins regulating TRPV5 expression or activity.

Hypomagnesemia is a widely known adverse effect of FK506 treatment (4, 5). The increased fractional excretion of Mg^{2+} in the presence of a profound hypomagnesemia indicated that renal Mg^{2+} reabsorption is seriously impaired by FK506. This renal Mg^{2+} wasting was accompanied by reduced renal expression of the putative Mg^{2+} channel TRPM6. Mutations in the gene encoding TRPM6 were recently shown to cause hereditary hypomagnesemia, which was also accompanied by an inappropriately increased fractional excretion of Mg^{2+} (14, 15). Importantly, RT-PCR analysis of microdissected rat nephrons showed strong TRPM6 expression in the DCT, the main site of active Mg^{2+} reabsorption (15). Together with the high structural homology to TRPM7, which has been previously identified to constitute a Mg^{2+} permeable cation channel, these data indicated that this protein constitutes the




Figure 5. Effect of FK506 and dexamethasone treatment on mRNA expression of Ca²⁺ and Mg²⁺ transport proteins in rat duodenum. Duodenal mRNA expression levels of the epithelial Ca²⁺ channel TRPV6, the cytosolic Ca²⁺-binding protein calbindin-D_{9K} (CaBP9), and the putative Mg²⁺ channel TRPM6 were determined by real-time quantitative PCR analysis. Ctr, controls; FK506, FK506 1 mg/d by oral gavage; Dex, dexamethasone 300 μg/d by oral gavage. Data are presented as means ± SEM. * *P* < 0.05 versus controls.

apical Mg²⁺ entry channel in renal transcellular Mg²⁺ transport in DCT (36, 37). Of note, the DCT has been previously indicated as a possible site of the tubular Mg²⁺ leak during FK506 treatment (13). The duodenal TRPM6 mRNA levels were not significantly altered by FK506 treatment, which is at variance with primary role of duodenal TRPM6 in the pathogenesis of hypomagnesemia. Altogether, our data showed that in addition to mutations in TRPM6, drug-induced downregulation of this ion channel in kidney is associated with increased urinary Mg²⁺ excretion and hypomagnesemia, further substantiating the importance of TRPM6 in renal Mg²⁺ reabsorption.

Dexamethasone-treated animals displayed a significant hypercalciuria, but, in contrast to FK506 treatment, dexamethasone increased the expression levels of TRPV5 and calbindin-D_{28K}. However, in line with previous data that showed decreased tubular reabsorption of Ca²⁺ during glucocorticoid treatment, the unaltered serum Ca²⁺ levels and GFR suggested that a primary impairment of Ca²⁺ reabsorption is responsible for the hypercalciuria observed in our study (2, 6). In the proximal tubule and TAL, passive reabsorption of Ca²⁺ takes place, which is altogether responsible for reabsorbing approximately 70 to 90% of the filtered Ca²⁺ (8). In this paracellular pathway, Ca²⁺ is transported down an electrochemical gradient that is dependent on Na⁺ reabsorption, and impairment of the latter is known to result in increased Ca²⁺ excretion. In this study, net Na⁺ excretion was indeed considerably increased and serum Na⁺ levels were reduced, whereas NCC expression was enhanced. Previously, glucocorticoid excess was demonstrated to enhance the expression of the distally located NCC and epithelial Na⁺ channel, whereas dexamethasone was shown to inhibit the function and expression of the proximal tubular Na⁺/phosphate co-transporter (38–41). Therefore, reduced reabsorption of Na⁺ in the proximal tubule or TAL could be responsible for the increased Na⁺ excretion and, by decreasing passive Ca²⁺ reabsorption, the dexamethasone-induced hypercalciuria. In addition, enzymuria was significantly increased by dexamethasone, suggesting that structural damage to the proximal tubule or TAL could contribute to the increased Na⁺ and Ca²⁺ excretion. The increased expression of the proteins involved in active Ca²⁺ transport in DCT indicated

that these proteins are directly or indirectly regulated by dexamethasone. The latter might reflect a compensatory effect secondary to the increased distal Ca²⁺ load, similar to the increased expression of distally located Na⁺ transporters, which has been shown after inhibition of Na⁺ reabsorption in the loop of Henle (42–44). The concomitant increase in expression of the major intestinal Ca²⁺ transport proteins suggests a direct stimulatory effect of dexamethasone on active Ca²⁺ (re)absorption in kidney and intestine.

It is interesting that dexamethasone treatment resulted in an increased urinary excretion of Mg²⁺ with a concomitant decrease of serum Mg²⁺ levels. However, the latter effect was much smaller than that observed in the FK506-treated animals and remained within the normal range, possibly because of the milder renal Mg²⁺ leak. Alternatively, increased intestinal Mg²⁺ absorption mediated by upregulated intestinal TRPM6 expression levels might counterbalance renal Mg²⁺ losses, but this is not supported by the present study. In analogy to Ca²⁺ reabsorption, a substantial part of Mg²⁺ filtered by the glomerulus is reabsorbed via the paracellular Na⁺-driven pathway (12). Therefore, impairment of the passive reabsorption of divalent ions in proximal tubule and TAL could explain both the hypercalciuria and hypermagnesuria. Subsequently, the enhanced TRPM6 expression levels could result from a direct stimulatory effect of dexamethasone or a compensatory response, serving to limit the renal Mg²⁺ wasting.

Indeed, serum glucose levels were significantly increased in both the FK506- and dexamethasone-treated animals. In the FK506-treated group, urinary glucose excretion did not significantly differ from controls, but the trend toward increased glycosuria could very well explain the concomitant trend toward increased urine volume. Furthermore, dexamethasone significantly increased glycosuria, which in conjunction with the increased Na⁺ excretion might cause the observed polyuria.

In conclusion, FK506-induced downregulation of Ca²⁺ and Mg²⁺ transport proteins could be a critical factor in the pathogenesis of hypercalciuria and hypomagnesemia. Elucidation of the molecular mechanism responsible for these inhibitory effects will extend our knowledge of the *in vivo* regulation of these transporters and will identify novel targets for pharmacologic therapy.

Acknowledgments

This study was supported by the Dutch Kidney Foundation (C10.1881) and the Dutch Organization of Scientific Research (Zon-Mw 016.006.001).

We thank the Fujisawa Pharmaceutical Co. for generously supplying tacrolimus (FK506).

References

- Rodino MA, Shane E: Osteoporosis after organ transplantation. *Am J Med* 104: 459–469, 1998
- Reid IR: Glucocorticoid osteoporosis—mechanisms and management. *Eur J Endocrinol* 137: 209–217, 1997
- Stempfle HU, Werner C, Siebert U, Assum T, Wehr U, Rambeck WA, Meiser B, Theisen K, Gartner R: The role of tacrolimus (FK506)-based immunosuppression on bone mineral density and bone turnover after cardiac transplantation: A prospective, longitudinal, randomized, double-blind trial with calcitriol. *Transplantation* 73: 547–552, 2002
- Andoh TF, Burdman EA, Fransechini N, Houghton DC, Bennett WM: Comparison of acute rapamycin nephrotoxicity with cyclosporine and FK506. *Kidney Int* 50: 1110–1117, 1996
- McDiarmid SV, Colonna JO, 2nd, Shaked A, Ament ME, Busuttil RW: A comparison of renal function in cyclosporine- and FK506-treated patients after primary orthotopic liver transplantation. *Transplantation* 56: 847–853, 1993
- Reid IR, Ibbertson HK: Evidence for decreased tubular reabsorption of calcium in glucocorticoid-treated asthmatics. *Horm Res* 27: 200–204, 1987
- Lukert BP, Raisz LG: Glucocorticoid-induced osteoporosis: Pathogenesis and management. *Ann Intern Med* 112: 352–364, 1990
- Friedman PA, Gesek FA: Cellular calcium transport in renal epithelia: Measurement, mechanisms, and regulation. *Physiol Rev* 75: 429–471, 1995
- Hoenderop JG, Nilius B, Bindels RJ: Molecular mechanisms of active Ca^{2+} reabsorption in the distal nephron. *Ann Rev Physiol* 64: 529–549, 2002
- Hoenderop JG, Willems PH, Bindels RJ: Toward a comprehensive molecular model of active calcium reabsorption. *Am J Physiol Renal Physiol* 278: F352–F360, 2000
- Lote CJ, Thewles A, Wood JA, Zafar T: The hypomagnesaemic action of FK506: Urinary excretion of magnesium and calcium and the role of parathyroid hormone. *Clin Sci (Lond)* 99: 285–292, 2000
- de Rouffignac C, Quamme G: Renal magnesium handling and its hormonal control. *Physiol Rev* 74: 305–322, 1994
- Dai LJ, Ritchie G, Kerstan D, Kang HS, Cole DE, Quamme GA: Magnesium transport in the renal distal convoluted tubule. *Physiol Rev* 81: 51–84, 2001
- Walder RY, Landau D, Meyer P, Shalev H, Tsolia M, Borochowitz Z, Boettger MB, Beck GE, Englehardt RK, Carmi R, Sheffield VC: Mutation of TRPM6 causes familial hypomagnesemia with secondary hypocalcemia. *Nat Genet* 31: 171–174, 2002
- Schlingmann KP, Weber S, Peters M, Niemann Nejsum L, Vitzthum H, Klingel K, Kratz M, Haddad E, Ristoff E, Dinour D, Syrrou M, Nielsen S, Sassen M, Waldegger S, Seyberth HW, Konrad M: Hypomagnesemia with secondary hypocalcemia is caused by mutations in TRPM6, a new member of the TRPM gene family. *Nat Genet* 31: 166–170, 2002
- Voets T, Nilius B, Hoefs S, Van Der Kemp AWCM, Droogmans G, Bindels RJM, Hoenderop JGJ: TRPM6 forms the Mg^{2+} influx channel involved in intestinal and renal Mg^{2+} absorption. *J Biol Chem* 279: 19–25, 2004
- Hoenderop JG, Hartog A, Stuiver M, Doucet A, Willems PH, Bindels RJ: Localization of the epithelial Ca^{2+} channel in rabbit kidney and intestine. *J Am Soc Nephrol* 11: 1171–1178, 2000
- Hoenderop JG, Muller D, Van Der Kemp AW, Hartog A, Suzuki M, Ishibashi K, Imai M, Sweep F, Willems PH, Van Os CH, Bindels RJ: Calcitriol controls the epithelial calcium channel in kidney. *J Am Soc Nephrol* 12: 1342–1349, 2001
- van Abel M, Hoenderop JG, Dardenne O, St-Arnaud R, Van Os C, van Leeuwen JP, Bindels RJ: 1, 25(OH) $_2$ D $_3$ -independent stimulatory effect of estrogen on the expression of ECaC1 in kidney. *J Am Soc Nephrol* 13: 2102–2109, 2002
- Nijenhuis T, Hoenderop JG, Loffing J, van der Kemp AW, Van Os C, Bindels RJ: Thiazide-induced hypocalcemia is accompanied by a decreased expression of Ca^{2+} transport proteins in kidney. *Kidney Int* 64: 555–564, 2003
- Hofmeister R, Bhargava AS, Gunzel P: Value of enzyme determinations in urine for the diagnosis of nephrotoxicity in rats. *Clin Chim Acta* 160: 163–167, 1986
- Zolotnitskaya A, Satlin LM: Developmental expression of ROMK in rat kidney. *Am J Physiol* 276: F825–F836, 1999
- Aicher L, Meier G, Norcross AJ, Jakubowski J, Varela MC, Cordier A, Steiner S: Decrease in kidney calbindin-D $_{28\text{kDa}}$ as a possible mechanism mediating cyclosporine A- and FK506-induced calciuria and tubular mineralization. *Biochem Pharmacol* 53: 723–731, 1997
- Hoenderop JG, Dardenne O, van Abel M, van der Kemp AW, Van Os CH, St-Arnaud R, Bindels RJ: Modulation of renal Ca^{2+} transport protein genes by dietary Ca^{2+} and 1, 25-dihydroxyvitamin D $_3$ in 25-hydroxyvitamin D $_3$ -1 α -hydroxylase knockout mice. *FASEB J* 16: 1398–1406, 2002
- Liu J, Farmer JD Jr, Lane WS, Friedman J, Weissman I, Schreiber SL: Calcineurin is a common target of cyclophilin-cyclosporin A and FKBP-FK506 complexes. *Cell* 66: 807–815, 1991
- Fruman DA, Klee CB, Bierer BE, Burakoff SJ: Calcineurin phosphatase activity in T lymphocytes is inhibited by FK 506 and cyclosporin A. *Proc Natl Acad Sci U S A* 89: 3686–3690, 1992
- O'Keefe SJ, Tamura J, Kincaid RL, Tocci MJ, O'Neill EA: FK506- and CsA-sensitive activation of the interleukin-2 promoter by calcineurin. *Nature* 357: 692–694, 1992
- Steiner S, Aicher L, Raymackers J, Meheus L, Esquer-Blasco R, Anderson NL, Cordier A: Cyclosporine A decreases the protein level of the calcium-binding protein calbindin-D $_{28\text{kDa}}$ in rat kidney. *Biochem Pharmacol* 51: 253–258, 1996
- Siekierka JJ, Hung SH, Poe M, Lin CS, Sigal NH: A cytosolic binding protein for the immunosuppressant FK506 has peptidyl-prolyl isomerase activity but is distinct from cyclophilin. *Nature* 341: 755–757, 1989
- Harding MW, Galat A, Uehling DE, Schreiber SL: A receptor for the immunosuppressant FK506 is a cis-trans peptidyl-prolyl isomerase. *Nature* 341: 758–760, 1989
- Schreiber SL: Chemistry and biology of the immunophilins and their immunosuppressive ligands. *Science* 251: 283–287, 1991
- Goel M, Garcia R, Estacion M, Schilling WP: Regulation of Drosophila TRPL channels by immunophilin FKBP59. *J Biol Chem* 276: 38762–38773, 2001
- Masumiya H, Wang R, Zhang J, Xiao B, Chen SR: Localization of the 12.6-kDa FK506-binding protein (FKBP12.6) binding site to the NH $_2$ -terminal domain of the cardiac Ca^{2+} release channel (ryanodine receptor). *J Biol Chem* 278: 3786–3792, 2003

34. Dargan SL, Lea EJ, Dawson AP: Modulation of type-1 Ins(1, 4, 5)P₃ receptor channels by the FK506-binding protein, FKBP12. *Biochem J* 361: 401–407, 2002
35. Cameron AM, Steiner JP, Sabatini DM, Kaplin AI, Walensky LD, Snyder SH: Immunophilin FK506 binding protein associated with inositol 1, 4, 5-trisphosphate receptor modulates calcium flux. *Proc Natl Acad Sci U S A* 92: 1784–1788, 1995
36. Runnels LW, Yue L, Clapham DE: TRP-PLIK, a bifunctional protein with kinase and ion channel activities. *Science* 291: 1043–1047, 2001
37. Nadler MJ, Hermosura MC, Inabe K, Perraud AL, Zhu Q, Stokes AJ, Kurosaki T, Kinet JP, Penner R, Scharenberg AM, Fleig A: LTRPC7 is a Mg²⁺-ATP-regulated divalent cation channel required for cell viability. *Nature* 411: 590–595, 2001
38. MacDonald P, MacKenzie S, Ramage LE, Seckl JR, Brown RW: Corticosteroid regulation of amiloride-sensitive sodium-channel subunit mRNA expression in mouse kidney. *J Endocrinol* 165: 25–37, 2000
39. Velazquez H, Bartiss A, Bernstein P, Ellison DH: Adrenal steroids stimulate thiazide-sensitive NaCl transport by rat renal distal tubules. *Am J Physiol* 270: F211–F219, 1996
40. Chen Z, Vaughn DA, Blakely P, Fanestil DD: Adrenocortical steroids increase renal thiazide diuretic receptor density and response. *J Am Soc Nephrol* 5: 1361–1368, 1994
41. Loffing J, Lotscher M, Kaissling B, Biber J, Murer H, Seikaly M, Alpern RJ, Levi M, Baum M, Moe OW: Renal Na/H exchanger NHE-3 and Na-PO₄ cotransporter NaPi-2 protein expression in glucocorticoid excess and deficient states. *J Am Soc Nephrol* 9: 1560–1567, 1998
42. Na KY, Oh YK, Han JS, Joo KW, Lee JS, Earm JH, Knepper MA, Kim GH: Upregulation of Na⁺ transporter abundances in response to chronic thiazide or loop diuretic treatment in rats. *Am J Physiol Renal Physiol* 284: F133–F143, 2003
43. Abdallah JG, Schrier RW, Edelstein C, Jennings SD, Wyse B, Ellison DH: Loop diuretic infusion increases thiazide-sensitive Na⁺/Cl⁻-cotransporter abundance: Role of aldosterone. *J Am Soc Nephrol* 12: 1335–1341, 2001
44. Ellison DH, Velazquez H, Wright FS: Adaptation of the distal convoluted tubule of the rat. Structural and functional effects of dietary salt intake and chronic diuretic infusion. *J Clin Invest* 83: 113–126, 1989

**Access to UpToDate on-line is available for additional clinical information
at <http://www.jasn.org/>**