

Angiotensin-Converting Enzyme Inhibition in Diabetic Nephropathy: It's All the RAGE

Raymond Harris

Division of Nephrology, Vanderbilt University School of Medicine, Nashville, Tennessee

J Am Soc Nephrol 16: 2251–2253, 2005. doi: 10.1681/ASN.2005060595

Since the landmark multicenter trial in type I diabetics by Lewis *et al.* in 1993 (1), the use of angiotensin-converting enzyme (ACE) inhibitors to retard progression of diabetic nephropathy has become a widely accepted therapeutic strategy. Although initial animal studies determined that ACE inhibitors decrease the increased glomerular capillary pressure (P_{GC}) associated with the hyperfiltration of diabetes (2,3), more recent studies have also attributed to angiotensin II many pathogenic changes observed in the glomerulus and tubulointerstitium in diabetes that are not necessarily related to altered intrarenal hemodynamics, including production of reactive oxygen species, cytokine and growth factor release, and alterations in extracellular matrix production. Recently, a number of studies have also indicated that drugs that interfere with the renin-angiotensin system may also decrease levels of advanced glycation endproducts (AGE) (4–6).

Although diabetic nephropathy is likely to have a multifactorial etiology (7) and to be subject in part to genetic predisposition, there is increasing evidence that AGE play an important role in its development. In the setting of diabetes mellitus and long-term hyperglycemia, free amino groups of proteins are nonenzymatically modified by glucose and its metabolites to form Schiff bases, which then rearrange to form Amadori products. Further modification by the Maillard reaction, nonenzymatic glycation, leads to formation of AGE. The most commonly formed AGE include N- ϵ -(carboxymethyl)lysine (CML), pentosidine, pyrraline, and imidazolone (8–12).

AGE may induce functional changes through nonreceptor-mediated pathways by crosslinking proteins, such as basement membrane. In addition, a number of AGE-binding proteins have been identified, including galectin-3, macrophage scavenger receptors, p60 complex, and p90. The best-described binding protein is receptor for advanced glycation endproducts (RAGE). RAGE is a member of the Ig superfamily of cell surface molecules (13) and is composed of a large N-terminal extracellular portion of 332 amino acid residues consisting of a distal V-type Ig domain, followed by two C-type Ig domains; ligands interact with the V-domain of the receptor (14–16). CML-aducts are a major class of AGE that bind to RAGE (17,18).

Following ligation of the extracellular region, transmembrane signaling by RAGE is thought to be transduced by the single hydrophobic transmembrane-spanning domain and short cytosolic tail. The intracellular portion of the molecule is essential for RAGE-triggered signaling and activates NF- κ B coupled signaling (17,18) as well as Cdc42-Rac-1-MKK6-p38 mitogen-activated protein kinase (MAPK) pathways (19,20). Binding of AGE to RAGE activates cell signaling mechanisms coupled to increased TGF- β and vascular endothelial growth factor (VEGF) expression that are thought to contribute to the pathogenesis of diabetic complications (16,21).

In this issue of *JASN*, Forbes *et al.* undertook studies investigating the potential mechanism underlying the observation that treatment with ACE inhibitors can decrease levels of circulating and tissue AGE (22). Previous *in vitro* studies by Miyata *et al.* had determined that direct incubation of diabetic serum with any of six angiotensin II type 1 receptor (AIIIR) antagonists or four ACE inhibitors inhibited the formation of both pentosidine and CML, with angiotensin receptor blockers (ARB) being somewhat more effective (5). In a rat model of type II diabetes, Nangaku *et al.* further reported that ARB, as well as hydralazine, could decrease kidney pentosidine content (6). In contrast, in both the current as well as former studies by Forbes and coworkers, there was a selective decrease of CML AGE in response to treatment with ACE inhibitors (4,22). These observations were confirmed in cell culture, in experimental animals, and in humans with incipient nephropathy secondary to type I diabetes. This selective decrease in CML and not pentosidine suggested a role for RAGE.

Recent studies have determined that three isoforms of RAGE are expressed through alternative splicing: the full-length transmembrane receptor, a soluble truncated form containing only the extracellular domain (sRAGE), and a truncated isoform containing the transmembrane and cytoplasmic portions of the receptor (N-truncated RAGE) (23). In their current studies, Forbes *et al.* determined that in poorly controlled diabetes, expression of the full length and N-truncated RAGE isoforms increased, while with ACE inhibitor treatment expression and secretion of sRAGE was selectively increased.

This study raises many interesting questions. What is the relationship between the ACE inhibitor-induced increases in sRAGE expression and decreases in circulating and tissue AGE, *i.e.*, is the selective upregulation of sRAGE a cause or effect of decreased circulating AGE? ACE inhibitors can

Published online ahead of print. Publication date available at www.jasn.org.

Address correspondence to: Dr. Raymond Harris, Division of Nephrology, C3121 MCN, Vanderbilt University School of Medicine, Nashville, TN 37232-2372. Phone: 615-343-0030; Fax: 615-343-2675; E-mail: ray.harris@vanderbilt.edu

decrease tissue oxidative stress, and there is evidence that inhibition of free radical production may decrease AGE production (5). Furthermore, there is some evidence that AGE may directly upregulate full-length RAGE (24). In these studies, administration of a mitochondrial reactive oxygen species scavenger to endothelial cells *in vitro* did lead to an increase in sRAGE secretion. Conversely, previous studies indicated that administration of sRAGE prevented diabetic atherosclerotic lesions and diabetic nephropathy in experimental animals (25) (16), and in their studies, Forbes *et al.* were able to identify circulating sRAGE/CML-AGE complexes.

These studies also did not determine whether the ACE inhibitor effect on sRAGE is the result of inhibiting angiotensin II-mediated signaling and, if so, the mechanism by which angiotensin II increases RAGE expression. Against a direct role for angiotensin II in these responses is the finding by Forbes *et al.* in their *in vitro* studies that the ARB valsartan did not increase sRAGE secretion; however, no *in vivo* studies were performed with ARB. If the effect on sRAGE expression does prove to be specific for ACE inhibitors, further studies will be necessary to determine whether it is a class effect, because only two ACE inhibitors (ramipril and perinopril) were used in these studies. If it is a class effect, is it secondary to increased bradykinin production or to some other structural characteristic of ACE inhibitors? Further studies will also be necessary to determine the biologic significance of this intriguing observation and to determine whether the ACE inhibitor-induced alterations in RAGE and AGE represent a previously unappreciated mechanism by which these drugs retard the progression of diabetic nephropathy.

References

- Lewis EJ, Hunsicker LG, Bain RP, Rohde RD: The effect of angiotensin-converting-enzyme inhibition on diabetic nephropathy. The Collaborative Study Group. *N Engl J Med* 329: 1456–1462, 1993
- Anderson S, Meyer TW, Rennke HG, Brenner BM: Control of glomerular hypertension limits glomerular injury in rats with reduced renal mass. *J Clin Invest* 76: 612–619, 1985
- Zatz R, Dunn BR, Meyer TW, Anderson S, Rennke HG, Brenner BM: Prevention of diabetic glomerulopathy by pharmacological amelioration of glomerular capillary hypertension. *J Clin Invest* 77: 1925–1930, 1986
- Forbes JM, Cooper ME, Thallas V, Burns WC, Thomas MC, Brammar GC, Lee F, Grant SL, Burrell LA, Jerums G, Osicka TM: Reduction of the accumulation of advanced glycation end products by ACE inhibition in experimental diabetic nephropathy. *Diabetes* 51: 3274–3282, 2002
- Miyata T, van Ypersele de Strihou C, Ueda Y, Ichimori K, Inagi R, Onogi H, Ishikawa N, Nangaku M, Kurokawa K: Angiotensin II receptor antagonists and angiotensin-converting enzyme inhibitors lower *in vitro* the formation of advanced glycation end products: Biochemical mechanisms. *J Am Soc Nephrol* 13: 2478–2487, 2002
- Nangaku M, Miyata T, Sada T, Mizuno M, Inagi R, Ueda Y, Ishikawa N, Yuzawa H, Koike H, van Ypersele de Strihou C, Kurokawa K: Anti-hypertensive agents inhibit *in vivo* the formation of advanced glycation end products and improve renal damage in a type 2 diabetic nephropathy rat model. *J Am Soc Nephrol* 14: 1212–1222, 2003
- Brownlee M: Biochemistry and molecular cell biology of diabetic complications. *Nature* 414: 813–820, 2001
- Frye EB, Degenhardt TP, Thorpe SR, Baynes JW: Role of the Maillard reaction in aging of tissue proteins. Advanced glycation end product-dependent increase in imidazolium cross-links in human lens proteins. *J Biol Chem* 273: 18714–18719, 1998
- Cohen MP: Intervention strategies to prevent pathogenetic effects of glycated albumin. *Arch Biochem Biophys* 419: 25–30, 2003
- Chen S, Cohen MP, Ziyadeh FN: Amadori-glycated albumin in diabetic nephropathy: Pathophysiologic connections. *Kidney Int Suppl* 77: S40–S44, 2000
- Vlassara H, Brownlee M, Cerami A: Nonenzymatic glycosylation of peripheral nerve protein in diabetes mellitus. *Proc Natl Acad Sci U S A* 78: 5190–5192, 1981
- Niwa T: 3-Deoxyglucosone: Metabolism, analysis, biological activity, and clinical implication. *J Chromatogr B Biomed Sci Appl* 731: 23–36, 1999
- Neeper M, Schmidt AM, Brett J, Yan SD, Wang F, Pan YC, Elliston K, Stern D, Shaw A: Cloning and expression of a cell surface receptor for advanced glycosylation end products of proteins. *J Biol Chem* 267: 14998–15004, 1992
- Schmidt AM, Yan SD, Wautier JL, Stern D: Activation of receptor for advanced glycation end products: A mechanism for chronic vascular dysfunction in diabetic vasculopathy and atherosclerosis. *Circ Res* 84: 489–497, 1999
- Yamagishi S, Takeuchi M, Inagaki Y, Nakamura K, Imaizumi T: Role of advanced glycation end products (AGEs) and their receptor (RAGE) in the pathogenesis of diabetic microangiopathy. *Int J Clin Pharmacol Res* 23: 129–134, 2003
- Wendt TM, Tanji N, Guo J, Kislinger TR, Qu W, Lu Y, Bucciarelli LG, Rong LL, Moser B, Markowitz GS, Stein G, Bierhaus A, Liliensiek B, Arnold B, Nawroth PP, Stern DM, D'Agati VD, Schmidt AM: RAGE drives the development of glomerulosclerosis and implicates podocyte activation in the pathogenesis of diabetic nephropathy. *Am J Pathol* 162: 1123–1137, 2003
- Kislinger T, Fu C, Huber B, Qu W, Taguchi A, Du Yan S, Hofmann M, Yan SF, Pischetsrieder M, Stern D, Schmidt AM: N(epsilon)-(carboxymethyl)lysine adducts of proteins are ligands for receptor for advanced glycation end products that activate cell signaling pathways and modulate gene expression. *J Biol Chem* 274: 31740–31749, 1999
- Hofmann MA, Drury S, Fu C, Qu W, Taguchi A, Lu Y, Avila C, Kambham N, Bierhaus A, Nawroth P, Neurath MF, Slattey T, Beach D, McClary J, Nagashima M, Morser J, Stern D, Schmidt AM: RAGE mediates a novel proinflammatory axis: A central cell surface receptor for S100/calgranulin polypeptides. *Cell* 97: 889–901, 1999
- Lander HM, Tauras JM, Ogiste JS, Hori O, Moss RA, Schmidt AM: Activation of the receptor for advanced glycation end products triggers a p21(ras)-dependent mitogen-activated protein kinase pathway regulated by oxidant stress. *J Biol Chem* 272: 17810–17814, 1997
- Sorci G, Riuzzi F, Arcuri C, Giambanco I, Donato R: Am-

- photerin stimulates myogenesis and counteracts the anti-myogenic factors basic fibroblast growth factor and S100B via RAGE binding. *Mol Cell Biol* 24: 4880–4894, 2004
21. Wendt T, Tanji N, Guo J, Hudson BI, Bierhaus A, Ramasamy R, Arnold B, Nawroth PP, Yan SF, D'Agati V, Schmidt AM: Glucose, glycation, and RAGE: Implications for amplification of cellular dysfunction in diabetic nephropathy. *J Am Soc Nephrol* 14: 1383–1395, 2003
 22. Forbes JM, Thorpe SR, Thallas-Bonke V, Pete J, Thomas MC, Deemer ER, Bassal S, El-Osta A, Long DM, Panagiotopoulos S, Jerums G, Osicka TM, Cooper ME: Modulation of soluble receptor for advanced glycation end products by angiotensin converting enzyme-1 inhibition in diabetic nephropathy. *J Am Soc Nephrol*, 2005
 23. Yonekura H, Yamamoto Y, Sakurai S, Petrova RG, Abedin MJ, Li H, Yasui K, Takeuchi M, Makita Z, Takasawa S, Okamoto H, Watanabe T, Yamamoto H: Novel splice variants of the receptor for advanced glycation end-products expressed in human vascular endothelial cells and pericytes, and their putative roles in diabetes-induced vascular injury. *Biochem J* 370: 1097–1109, 2003
 24. Tanaka N, Yonekura H, Yamagishi S, Fujimori H, Yamamoto Y, Yamamoto H: The receptor for advanced glycation end products is induced by the glycation products themselves and tumor necrosis factor-alpha through nuclear factor-kappa B, and by 17beta-estradiol through Sp-1 in human vascular endothelial cells. *J Biol Chem* 275: 25781–25790, 2000
 25. Park L, Raman KG, Lee KJ, Lu Y, Ferran LJ Jr, Chow WS, Stern D, Schmidt AM: Suppression of accelerated diabetic atherosclerosis by the soluble receptor for advanced glycation endproducts. *Nat Med* 4: 1025–1031, 1998

See related article, "Modulation of Soluble Receptor for Advanced Glycation End Products by Angiotensin Converting Enzyme-1 Inhibition in Diabetic Nephropathy," on pages 2363–2372.