

# Renal Fibroblasts and Myofibroblasts in Chronic Kidney Disease

Frank Strutz\* and Michael Zeisberg<sup>†</sup>

\*Department of Nephrology and Rheumatology, Georg-August-University Medical Center, Göttingen, Germany; and

<sup>†</sup>Division of Matrix Biology, Beth Israel Deaconess Hospital, Harvard Medical School, Boston

J Am Soc Nephrol 17: 2992–2998, 2006. doi: 10.1681/ASN.2006050420

Fibroblasts and myofibroblasts are believed to be the key effector cells in renal fibrogenesis responsible for the synthesis and deposition of extracellular matrix components. In an editorial written 15 yr ago, Gown claimed to have the mysteries of the fibroblast (partially) unmasked (1). Unfortunately, 15 yr later many mysteries of the fibroblast and myofibroblasts remain intact and we are only beginning to solve the riddle this cell poses. This manuscript summarizes current understanding of the origin, phenotypic heterogeneity, and function of renal (myo)fibroblasts.

Renal interstitial fibrosis is considered the hallmark of progressive renal disease. In particular, many studies have determined that the extent of tubulointerstitial involvement correlates better with renal function deterioration than glomerular changes do, thus the extent of damaged tubulointerstitial area in any given renal biopsy has important implications for the renal prognosis of the patient (summarized in [2]). Tubulointerstitial fibrosis is characterized by the accumulation of extracellular matrix (ECM) components including collagen types I, III, and IV, as well as proteoglycans and fibronectin. Fibroblasts are considered the primary matrix-producing cells in the kidney and hence they are clinically relevant as principal mediators of renal fibrosis associated with progressive renal failure.

## Fibroblasts in the Kidney

Originally, resident interstitial fibroblasts were described as contractile cellular elements being the sole source of ECM (3). Today, interstitial fibroblasts can be best defined as the nonvascular, nonepithelial, noninflammatory cells of the interstitium (and of connective tissue in general), and they are its principal cellular component (4). Fibroblasts synthesize many of the constituents of the fibrillar ECM, such as fibronectin and types I, III, and V collagen (5,6), and they are also a major source of

ECM-degrading proteases such as matrix metalloproteinases (MMP), underscoring their crucial role in maintaining an ECM homeostasis *via* regulation of ECM turnover (7,8). Fibroblasts are considered mesenchymal cells that display a spindle-shaped morphology (9). However, due to the relative lack of specific markers, it can be challenging to distinguish fibroblasts from other cells of mesenchymal origin, such as pericytes, vascular smooth muscle cells (VSMC), or mesenchymal stem cells in the kidney (Table 1), and thus renal fibroblasts have remained relatively poorly characterized in molecular terms.

## What Is the Myofibroblast?

The term “myofibroblast” is used for fibroblasts that possess unique contractile properties (6). Myofibroblasts were originally identified as the cells being responsible for wound contraction (10). In the setting of renal fibrosis, myofibroblasts are considered to represent an activated fibroblast phenotype, which is mainly responsible for ECM deposition in tubulointerstitial fibrosis (11). Myofibroblasts are large cells with long processes resembling fibroblasts and VSMC (12). In addition, they are characterized by bundled microfilaments, a well-developed rough endoplasmic reticulum, hemidesmosomes, and other intercellular attachments (11). In the absence of electron microscopy validation, myofibroblasts are commonly identified by their expression of  $\alpha$ -smooth muscle actin ( $\alpha$ -SMA) (6,13). Identification of myofibroblasts based on the expression of  $\alpha$ -SMA can be challenging, however, because  $\alpha$ -SMA is also expressed by other cells of mesenchymal lineage such as VSMC and pericytes (1,4) (Table 1).

Traditionally, myofibroblasts in the kidney have been considered to represent an activated population of resident fibroblasts (14). However, recent studies questioned whether the increase in the number of  $\alpha$ -SMA-positive cells is the result of residual proliferating smooth muscle cells, proliferation of the few fibroblasts constitutively expressing the protein, or by *de novo* expression of formerly nonexpressing cells. This issue is further complicated by the observation that in anti-glomerular basement membrane antibody-induced nephritis (15), as well as in periglomerular areas in a study by the same group (16), early expansion of  $\alpha$ -SMA-positive myofibroblasts is observed in perivascular regions similar to what had been described

Published online ahead of print. Publication date available at [www.jasn.org](http://www.jasn.org).

**Address correspondence to:** Dr. Frank Strutz, Department of Nephrology and Rheumatology, Georg-August-University Medical Center, Robert-Koch-Str. 40, 37099 Göttingen, Germany. Phone: +49-551-396981; Fax: +49-551-398906; E-mail: [fstrutz@gwdg.de](mailto:fstrutz@gwdg.de); or Dr. Michael Zeisberg, Division of Matrix Biology, Beth Israel Deaconess Medical Center, 330 Brookline Ave., RW511, Boston, MA 02215. Phone: 617-667-0467; Fax: 617-667-2562; E-mail: [mzeisber@bidmc.harvard.edu](mailto:mzeisber@bidmc.harvard.edu)

Table 1. Cell types with mesenchymal phenotypes detected in the kidney<sup>a</sup>

Cell Type	Markers	Specific Characteristics	References
Resident Fibroblasts	FSP1, HSP47, 5'ectonucleotidase, CD44, ICAM, DDR2	Phenotypic variability between cortical and medullary fibroblasts	41, 44, 63
Myofibroblasts	$\alpha$ -SMA		13
Pericytes	$\alpha$ -SMA, NG2, PDGFR-2, Desmin	Associated capillaries and venules	64, 65
Vascular smooth muscle cells	$\alpha$ -SMA, caldesmon, calponin	Associated arteries, arterioles, veins	64
EMT-derived fibroblasts	FSP1 (residual epithelial markers?)	Exclusively found in fibrotic kidneys	41, 44
Mesenchymal stem cells	FSP1?		44

<sup>a</sup>At least 6 distinct cell types with mesenchymal phenotypes can be detected in the kidney. Due to the lack of specific markers, identification of the different lineages can be challenging. In the normal kidney, myofibroblasts can be distinguished from vascular smooth muscle cells and pericytes based on topographical criteria. Without genetic markers, indisputable identification of EMT-derived cells and mesenchymal stem cells is not feasible as of yet. HSP47, heat shock protein 47; ICAM, intracellular adhesion molecule; DDR2, discoidin domain receptor;  $\alpha$ -SMA,  $\alpha$ -smooth muscle actin; NG2, neuron glial antigen 2; PDGFR, PDGF receptor; EMT, epithelial–mesenchymal transition.

Wiggins *et al.* (15), suggesting that shedding of VSMC and pericytes contributes to the population of  $\alpha$ -SMA–positive cells in the fibrotic interstitium. Furthermore, at least for some fibroblasts, activation is not associated with *de novo* expression of  $\alpha$ -SMA (17). Ru and colleagues described that even among so-called myofibroblasts a higher degree of variability and complexity due to different levels of  $\alpha$ -SMA expression may exist (18). In summary, all these studies support the notion that fibroblasts from kidneys with interstitial fibrosis differ from fibroblasts in normal kidneys.

#### Myofibroblasts and Nonmyofibroblasts

*De novo* expression of  $\alpha$ -SMA is not the only change during the transition from fibroblast to myofibroblast. In fact, the radical change in phenotype is often underappreciated (19). The differences between fibroblasts from normal or fibrotic kidneys extend to their proliferative activity and matrix synthesis capacity (20). Müller and Rodemann determined that the relative percentage of fibroblasts was much higher in primary cell cultures from kidneys with fibrosis compared with their normal counterparts (47 versus 7%) (21). This finding was confirmed in a study by our group on primary human cortical fibroblasts (22), indicating an increased proliferative activity of activated fibroblasts. Furthermore, Rodemann and colleagues demonstrated that fibroblasts from fibrotic kidneys displayed a higher mitogenic potential, a higher resistance to the mitogenic drug mitomycin C, and a three- to five-fold increase in the rate of collagen synthesis as determined by <sup>3</sup>H-proline incorporation (23), suggesting that such an activated phenotype of fibroblasts observed in the cell culture experiments was representative of their behavior *in vivo*. However, the preservation of this activation state is relatively poorly understood in molecular terms.

#### Fibroblast Activation

Various stimuli have been found to induce fibroblast activation (24). In general, fibroblasts respond to stimuli associated with tissue injury by acquiring an activated phenotype (Figure 1). For example, infiltration of inflammatory cells results in the




Figure 1. Activation of resident fibroblasts by stimuli associated with tissue injury.

activation and proliferation of resident fibroblasts. These resident fibroblasts become activated by stimulation with cytokines including TGF- $\beta$ 1 (25), PDGF (26), and fibroblast growth factor 2 (FGF-2) (20), among others. Additional activation mechanisms include direct cell–cell contact (leukocytes and macrophages), ECM–integrin interaction (mainly  $\alpha$ 1 and  $\beta$ 1), as well as environmental stimuli such as hypoxia and hyperglycemia (11). Most authors concur that cortical fibroblasts can proliferate (20), although no fibroblast proliferation was observed in a recent report by Yamashita *et al.* (27). In the setting of tubulo-interstitial fibrosis, renal fibroblasts maintain their activated phenotype, even after the initial insult no longer exists (28). How such an activated state is maintained is only incompletely understood. Several studies demonstrated that autocrine secretion of fibroblasts is involved in this process. For example, our research group demonstrated that autocrine secretion of FGF-2 contributes to maintenance of an activated phenotype (29).

Several recent studies in the cancer field suggested that epigenetic changes in response to cell injury could causally underlie such imprinted activation (4,30). Whether such epigenetic changes are involved in fibroblast activation associated with tubulointerstitial fibrosis remains to be determined.

There is still a question of how these activated fibroblasts are cleared during resolution of a fibrotic process. One potential pathway is apoptosis, as was shown for activated Ito cells in the liver (31). In addition, reversion to a quiescent phenotype may be a second possibility, but this remains to be proven. If this possibility exists, then there may be a point of no return, after which autonomous fibroblast proliferation cannot be stopped and fibrogenesis invariably progresses. This is again of clinical importance because it was recently shown that fibrotic processes are potentially reversible in the kidney (32) as well as in other organs such as the liver (33). Understanding of physiologic clearance of activated fibroblasts is particularly important to develop therapeutic strategies to inhibit the progression of tubulointerstitial fibrosis. While ablation of activated fibroblasts has been proven to be effective in the setting of fibroblast-specific protein 1 (FSP1)-tk transgenic mice, due to the lack of fibroblast-specific targets no such approach has been developed for potential therapy as yet. Most therapeutic approaches are thus designed to “normalize” activated fibroblasts, reverting them to their quiescent physiologic phenotype. In this regard, our group has successfully demonstrated that excessive matrix synthesis and proliferative activity could be inhibited by various molecules such as pentoxifylline, IFN- $\gamma$ , relaxin, or bone morphogenic protein-7 (BMP-7) (34–36). More understanding of fibroblast-specific biology is required to design more specific therapeutic agents.

#### Origin of Fibroblasts in the Kidney

Due to the lack of systematic lineage tracing studies, the origin of fibroblasts is still controversial (4). As early as 1867, Cohnheim wrote a classic manuscript on chronic inflammation, in which he stated that fibroblasts (he called them contractile cellular elements at the time) are direct descendents of migrating leukocytes (3). However, over 100 years later, Ross *et al.* demonstrated, in a very elegant set of experiments using parabiotic rats, that fibroblasts are not derived from blood cells but were instead of local origin (37). Such thinking was corroborated by studies that demonstrated that adult fibroblasts localized to the sites of embryonic mesenchymal cells, suggesting that adult fibroblasts are direct derivatives of embryonic progenitor cells. For example, a study, which evaluated mesenchymal cells and fibroblasts in rat kidneys from E14 to E28 and postnatally until day 28 suggested that the interstitial kidney fibroblasts are derivatives from the uninduced mesenchyme (38). While molecular evidence for such thinking is lacking in the kidney, an elegant study that used retrovirus-mediated lineage analysis of the inner ear revealed that fibroblasts have a common mesenchymal progenitor (39), suggesting that adult resident fibroblasts arise locally during embryogenesis *via* division and differentiation of mesenchymal progenitor cells. However, recent studies point again to a possible bone marrow origin of at least some fibroblasts, and the concept of fibroblast

stem cells has gained wide popularity, although its exact contribution remains to be determined (40). Moreover, periadventitial cells (15) and tubular epithelial cells (41) have been implicated as the origin of matrix-producing cells (Figure 2). In summary, evidence is evolving that the origin of kidney fibroblasts is as diverse as their phenotypic heterogeneity.

As opposed to fibroblasts being direct derivatives from resident embryonic mesenchymal cells, a number of recent studies have confirmed the existence of bone marrow–derived cells within the interstitium (42) and the mesenchymal–epithelial transition of bone marrow cells to tubular epithelium (43). Preliminary evidence using genetically tagged fibroblasts and tubular epithelial cells indicates that fibroblasts derived from bone marrow comprise about 12% of the resident interstitial population in normal murine kidneys (44). In that study, this percentage did not change when an experimental model of progressive renal disease was induced. Conversely, up to 36% of additional ECM-producing cells derived from tubular epithelial cells by epithelial–mesenchymal “transition” (EMT; see below). Moreover, in a study in human biopsies from patients with chronic allograft nephropathy, it was suggested that up to 30% of interstitial mesenchymal cells were of bone marrow origin (42). However, a recent and very elegant study by Roufosse and coworkers used a transgenic mouse expressing both luciferase as well as  $\beta$ -galactosidase reporter molecules under control of the promoter for the  $\alpha$ 2-chain of type I collagen (45). Using these mice as male bone marrow donors into female wildtype mice, a mean of 8.6% of  $\alpha$ -SMA–positive cells within the tubulointerstitial space were identified by positivity for the Y chromosome after induction of unilateral ureteral obstruction (UUO) in recipients. The bone marrow–derived cells, however, were completely negative for luciferase, indicating a lack of collagen synthesis. In summary, with respect to the technical difficulties that come with the available reporter gene techniques, the origin of fibroblasts remains an unsolved question and further molecular studies are required to determine the contribution of bone marrow–derived cells to fibroblasts in the kidney.




Figure 2. Possible origin of matrix-producing cells in the kidney.

### *Epithelial-Mesenchymal Transition in Renal Fibrogenesis*

The term “EMT” is used preferentially to describe the conversion of terminally differentiated epithelia into cells with a mesenchymal phenotype (*i.e.*, cells with a fibroblastoid appearance and positivity for mesenchymal cell markers) (44). EMT is a variant of transdifferentiation and a well recognized mechanism for dispersing cells in vertebrate embryos (46), forming fibroblasts in injured tissues (41,44), or initiating metastases in epithelial cancer (47). The conversion of epithelial cells to mesenchymal cells was first described in 1982 in a seminal study by Greenburg and Hay (48).

Regarding renal fibrogenesis, EMT was first demonstrated by our group in a murine model of anti-tubular basement membrane disease by cloning of FSP1, a member of the S100 family (41). FSP1 expression is constitutive in tissue fibroblasts under physiologic conditions (41), and its promoter contains a *cis*-acting element (FTS-1) highly specific for fibroblasts (49). Although there has been some controversy regarding the exclusivity of FSP1 expression in fibroblasts (50,51), this specificity was demonstrated in an elegant experiment, in which cells expressing green fluorescent protein (GFP) under control of the FSP1 promoter were isolated from kidneys of FSP1-GFP transgenic mice and analyzed by FACS, demonstrating that the FSP1-GFP-positive cells are indeed exclusively fibroblasts (51). Furthermore, by placing the herpes virus thymidine kinase under the control of the FSP1 promoter in transgenic mice, and by the specific killing of fibroblasts *via* addition of nucleoside analogs, interstitial fibrosis is prevented, further indicating the specificity of FSP1-positive fibroblasts (52). Analyzing two mouse models of chronic progressive renal disease, FSP1 expression was robustly upregulated in the tubulointerstitium. The initial staining pattern was very similar to the distribution of collagen-producing cells in a rabbit model of anti-glomerular basement membrane disease with a perivascular accentuation (15). However, *de novo* expression of this fibroblast-specific protein could be detected in tubular epithelial cells in both models of chronic progressive renal disease, suggesting a possible transition of these cells to a mesenchymal phenotype (53). This phenomenon had been described in a number of organs, including the thyroid and mammary glands, as well as the retina (reviewed in [46]). EMT is a dynamic process that involves the gradual loss of epithelial cell markers associated with a gain of mesenchymal markers. This process is associated with various intermediate stages of cells that co-express epithelial cell markers as well as mesenchymal markers. To distinguish stress fiber-expressing epithelial cells from cells that are actively undergoing EMT, the loss of the epithelial adhesion molecules such as E-cadherin and ZO-1 paired with increased expression of mesenchymal markers such as vimentin,  $\alpha$ -SMA, and/or FSP1, in addition to migratory activity, are used as criteria to detect EMT. However, absolute criteria have not been defined. Recently, Nishitani and colleagues described the use of the FSP1-antibody as a prognostic marker in patients with IgA nephropathy, indicating its potential usefulness even in human biopsies (54).

Regarding the situation in the kidney, the occurrence of this

phenomenon is not surprising. Because tubular epithelial cells (with the exception of collecting duct cells) are derivatives of the metanephric mesenchyme, they seem to be particularly well suited to undergo phenotypic changes toward a mesenchymal phenotype. Thus, this process can be described as a reiteration of renal developmental programs (55). *De novo* expression of FSP1 is not the only change observed in tubular epithelial cells undergoing EMT. Ng and coworkers described the *de novo* expression of the myofibroblast marker  $\alpha$ -SMA in tubular epithelial cells starting at day 21 after 5/6 nephrectomy (56). Tubular epithelial cells lost their apical-basal polarity and seemed to migrate into the interstitium as demonstrated by electron microscopy. However, it is interesting to note that FSP1 and  $\alpha$ -SMA seem to identify different fibroblast populations, since the coexpression of both molecules seem to be the exception rather than the rule (17).

It is still unclear whether all tubular epithelial cells can potentially undergo EMT. Most commonly, proximal tubular but also distal tubular epithelial cells have been implicated in EMT. In a recent study, Yamashita and colleagues suggested that there is a distinct population of renal progenitor tubular cells that may undergo EMT (27). However, studies by our group and by others predominantly suggest that all tubular epithelial cells possess the capacity to undergo EMT.

### *Is the Origin of the Renal Fibroblast Comparable in Different Models?*

Even considering the heterogeneity of fibroblasts in the kidney and the various markers that are commonly used to detect them, there is a remarkable variability in the results of recent studies that evaluated renal fibroblasts in kidney disease. A possible explanation for such discrepancy is that fibroblast recruitment occurs in a disease-specific manner. In this regard, Faulkner and colleagues analyzed the origin of  $\alpha$ -SMA-positive cells in an accelerated model of angiotensin II-induced renal fibrosis after Habu venom injury. Faulkner *et al.* concluded that most if not all of the tubulointerstitial  $\alpha$ -SMA-expressing cells were of local origin (57). There were no disruptions of tubular basement integrity, and no traffic of proximal tubular epithelial cells into the interstitial compartment was observed by labeling with Texas Red dextran (57). Similarly, EMT was not observed in a model of overload proteinuria (58). However, using  $\alpha$ -SMA expression alone as a marker for matrix-producing cells may not sample all these cells, as was nicely demonstrated by Okada and coworkers in a model of murine polycystic kidney disease (17). Thus, the origin of matrix-producing cells may vary according to the model used, although we need more studies that apply transgenic techniques given the difficulties in identifying matrix-synthesizing cells by common immunohistochemistry techniques. Iwano and colleagues showed very convincingly the relative importance of EMT for the UUO model (44). Moreover, the specific significance of EMT for the progression of renal disease was shown very convincingly by Yang *et al.*, who demonstrated that EMT was critical for the progression of renal disease in the UUO model compared with the tissue plasminogen activator-deficient mice (59). The relative contribution of EMT to interstitial matrix deposition may also have therapeutic

implications. Bone morphogenetic protein (BMP)-7 is a morphogen that was recently shown to be capable of halting and reverting renal fibrosis in experimental animal models (35,60,61). One of the main effects of BMP-7 is the inhibition of EMT (61). Additional therapeutic agents, such as hepatocyte growth factor, which may halt progression in experimental models, also act at least in part by inhibition of EMT (62).

### Conclusion

In summary, the mysteries of the renal fibroblast are difficult to unmask and many aspects of these cells remain unsolved. The relative lack of reliable marker proteins, particularly for human cortical fibroblasts, may be an indication of the heterogeneity of this cell type. Conversely, myofibroblasts can be easily detected *in vivo* via expression of  $\alpha$ -SMA, but it is probably only one of many activation states of fibroblasts if we define these cells as interstitial matrix-producing cells. Matrix-synthesizing cells may be derived from resident interstitial cells, mesenchymal and/or hematopoietic stem cells, perivascular cells, and/or by the process of epithelial mesenchymal transition. The renal community has learned in recent years that the relative contribution of these various cellular elements may vary according to the model of progressive renal disease studied. From a clinical point of view, it is interesting that this contribution may have therapeutic implications. Research into fibroblasts and their origin will be an exciting area in which to work in the years to come.

### References

- Gown AM: The mysteries of the myofibroblast (partially) unmasked. *Lab Invest* 63: 1–3, 1990
- Bohle A, Strutz F, Muller GA: On the pathogenesis of chronic renal failure in primary glomerulopathies. *Exp Nephrol* 2: 205–210, 1994
- Cohnheim J: Über Entzündung und Eiterung. *Virchows Arch* 40: 1–79, 1867
- Kalluri R, Zeisberg M: Fibroblasts in cancer. *Nat Rev Cancer* 6: 392–401, 2006
- Rodemann HP, Muller GA: Characterization of human renal fibroblasts in health and disease: II. In vitro growth, differentiation, and collagen synthesis of fibroblasts from kidneys with interstitial fibrosis. *Am J Kidney Dis* 17: 684–686, 1991
- Tomasek JJ, Gabbiani G, Hinz B, Chaponnier C, Brown RA: Myofibroblasts and mechano-regulation of connective tissue remodelling. *Nat Rev Mol Cell Biol* 3: 349–363, 2002
- Chang HY, Chi JT, Dudoit S, Bondre C, van de Rijn M, Botstein D, Brown PO: Diversity, topographic differentiation, and positional memory in human fibroblasts. *Proc Natl Acad Sci U S A* 99: 12877–12882, 2002
- Simian M, Hirai Y, Navre M, Werb Z, Lochter A, Bissell MJ: The interplay of matrix metalloproteinases, morphogens and growth factors is necessary for branching of mammary epithelial cells. *Development* 128: 3117–3131, 2001
- Strutz F, Muller GA: Renal fibrosis and the origin of the renal fibroblast. *Nephrol Dial Transplant* August 3, 2006 [pub ahead of print]
- Majno G, Gabbiani G, Hirschel BJ, Ryan GB, Statkov PR: Contraction of granulation tissue in vitro: Similarity to smooth muscle. *Science* 173: 548–550, 1971
- Qi W, Chen X, Poronnik P, Pollock CA: The renal cortical fibroblast in renal tubulointerstitial fibrosis. *Int J Biochem Cell Biol* 38: 1–5, 2006
- Ina K, Kitamura H, Tatsukawa S, Takayama T, Fujikura Y, Shimada T: Transformation of interstitial fibroblasts and tubulointerstitial fibrosis in diabetic nephropathy. *Med Electron Microsc* 35: 87–95, 2002
- Sappino AP, Schurch W, Gabbiani G: Biology of disease. Differentiation repertoire of fibroblastic cells: Expression of cytoskeletal proteins as marker of phenotypic modulations. *Lab Invest* 63: 144–161, 1990
- Zeisberg M, Strutz F, Muller GA: Role of fibroblast activation in inducing interstitial fibrosis. *J Nephrol* 13[Suppl 3]: S111–S120, 2000
- Wiggins R, Goyal M, Merritt S, Killen PD: Vascular adventitial cell expression of collagen I messenger ribonucleic acid in anti-glomerular basement membrane antibody-induced crescentic nephritis in the rabbit. *Lab Invest* 68: 557–565, 1993
- Barnes VL, Musa J, Mitchell RJ, Barnes JL: Expression of embryonic fibronectin isoform EIIIA parallels alpha-smooth muscle actin in maturing and diseased kidney. *J Histochem Cytochem* 47: 787–798, 1999
- Okada H, Ban S, Nagao S, Takahashi H, Suzuki H, Neilson EG: Progressive renal fibrosis in murine polycystic kidney disease: An immunohistochemical observation. *Kidney Int* 58: 587–597, 2000
- Ru Y, Eyden B, Curry A, McWilliam LJ, Coyne JD: Actin filaments in human renal tubulo-interstitial fibrosis: Significance for the concept of epithelial-myofibroblast transformation. *J Submicrosc Cytol Pathol* 35: 221–233, 2003
- Grinnell F: Fibroblasts, myofibroblasts, and wound contraction. *J Cell Biol* 124: 401–404, 1994
- Strutz F, Zeisberg M, Hemmerlein B, Sattler B, Hummel K, Becker V, Muller GA: Basic fibroblast growth factor (FGF-2) expression is increased in human renal fibrogenesis and may mediate autocrine fibroblast proliferation. *Kidney Int* 57: 1521–1538, 2000
- Muller GA, Rodemann HP: Characterization of human renal fibroblasts in health and disease: I. Immunophenotyping of cultured tubular epithelial cells and fibroblasts derived from kidneys with histologically proven interstitial fibrosis. *Am J Kidney Dis* 17: 680–683, 1991
- Strutz F, Renziehausen A, Dietrich M, Amin J, Becker V, Heeg M, Rastaldi MP, Muller GA: Cortical fibroblast culture from human biopsies. *J Nephrol* 14: 190–197, 2001
- Rodemann HP, Muller GA, Knecht A, Norman JT, Fine LG: Fibroblasts of rabbit kidney culture I. Characterization and identification of cell-specific markers. *Am J Physiol* 261: F283–F291, 1991
- Zeisberg M, Strutz F, Muller GA: Renal fibrosis: An update. *Curr Opin Nephrol Hypertens* 10: 315–320, 2001
- Alvarez RJ, Sun MJ, Haverty TP, Iozzo RV, Meyers JC, Neilson EG: Biosynthetic and proliferative characteristics of tubulointerstitial fibroblasts probed with paracrine cytokines. *Kidney Int* 41: 14–23, 1992
- Alpers CE, Seifert RA, Hudkins KL, Johnson RJ, Bowen-Pope DF: PDGF-receptor localizes to mesangial, parietal epithelial, and interstitial cells in human and primate kidneys. *Kidney Int* 43: 286–294, 1993

27. Yamashita S, Maeshima A, Nojima Y: Involvement of renal progenitor tubular cells in epithelial-to-mesenchymal transition in fibrotic rat kidneys. *J Am Soc Nephrol* 16: 2044–2051, 2005
28. Strutz F, Muller GA: On the progression of chronic renal disease. *Nephron* 69: 371–379, 1995
29. Strutz F, Zeisberg M, Ziyadeh FN, Yang CQ, Kalluri R, Muller GA, Neilson EG: Role of basic fibroblast growth factor-2 in epithelial-mesenchymal transformation. *Kidney Int* 61: 1714–1728, 2002
30. Orimo A, Gupta PB, Sgroi DC, Arenzana-Seisdedos F, Delaunay T, Naeem R, Carey VJ, Richardson AL, Weinberg RA: Stromal fibroblasts present in invasive human breast carcinomas promote tumor growth and angiogenesis through elevated SDF-1/CXCL12 secretion. *Cell* 121: 335–348, 2005
31. Desmouliere A, Gabbiani G: Myofibroblast differentiation during fibrosis. *Exp Nephrol* 3: 134–139, 1995
32. Strutz F, Zeisberg M: Epithelial-mesenchymal transition and the reversal of renal fibrosis. *Transworld Net* 2006, in press
33. Bataller R, Brenner DA: Liver fibrosis. *J Clin Invest* 115: 209–218, 2005
34. Strutz F, Heeg M, Kochsiek T, Siemers G, Zeisberg M, Muller GA: Effects of pentoxifylline, pentifylline and gamma-interferon on proliferation, differentiation, and matrix synthesis of human renal fibroblasts. *Nephrol Dial Transplant* 15: 1535–1546, 2000
35. Zeisberg M, Bottiglio C, Kumar N, Maeshima Y, Strutz F, Muller GA, Kalluri R: Bone morphogenic protein-7 inhibits progression of chronic renal fibrosis associated with two genetic mouse models. *Am J Physiol Renal Physiol* 285: F1060–F1067, 2003
36. Heeg MH, Koziolk MJ, Vasko R, Schaefer L, Sharma K, Muller GA, Strutz F: The antifibrotic effects of relaxin in human renal fibroblasts are mediated in part by inhibition of the Smad2 pathway. *Kidney Int* 68: 96–109, 2005
37. Ross R, Everett NB, Tyler R: Wound healing and collagen formation. VI. The origin of the wound. Fibroblast studied in parabiosis. *J Cell Biol* 44: 645–654, 1970
38. Ekblom P, Weller A: Ontogeny of tubulointerstitial cells. *Kidney Int* 39: 394–400, 1991
39. Lang H, Fekete DM: Lineage analysis in the chicken inner ear shows differences in clonal dispersion for epithelial, neuronal, and mesenchymal cells. *Dev Biol* 234: 120–137, 2001
40. Powell DW, Mifflin RC, Valentich JD, Crowe SE, Saada JI, West AB: Myofibroblasts. II. Intestinal subepithelial myofibroblasts. *Am J Physiol* 277: C183–201, 1999
41. Strutz F, Okada H, Lo CW, Danoff T, Carone B, Tomaszewski J, Neilson EG: Identification and characterization of fibroblast-specific protein 1 (FSP1). *J Cell Biol* 130: 393–405, 1995
42. Grimm PC, Nickerson P, Jeffery J, Savani RC, Gough J, McKenna RM, Stern E, Rush DN: Neointimal and tubulointerstitial infiltration by recipient mesenchymal cells in chronic renal-allograft rejection. *N Engl J Med* 345: 93–97, 2001
43. Poulosom R, Forbes SJ, Hodivala-Dilke K, Ryan E, Wyles S, Navaratnasah S, Jeffery R, Hunt T, Alison M, Cook T, Pusey C, Wright NA: Bone marrow contributes to renal parenchymal turnover and regeneration. *J Pathol* 195: 229–235, 2001
44. Iwano M, Plieth D, Danoff TM, Xue C, Okada H, Neilson EG: Evidence that fibroblasts derive from epithelium during tissue fibrosis. *J Clin Invest* 110: 341–350, 2002
45. Roufosse C, Bou-Gharios G, Prodromidi E, Alexakis C, Jeffery R, Khan S, Otto WR, Alter J, Poulosom R, Cook HT: Bone marrow-derived cells do not contribute significantly to collagen I synthesis in a murine model of renal fibrosis. *J Am Soc Nephrol* 17: 775–782, 2006
46. Hay ED, Zuk A: Transformations between epithelium and mesenchyme: Normal, pathological, and experimentally induced. *Am J Kidney Dis* 26: 678–690, 1995
47. Yang J, Mani SA, Donaher JL, Ramaswamy S, Itzykson RA, Come C, Savagner P, Gitelman I, Richardson A, Weinberg RA: Twist, a master regulator of morphogenesis, plays an essential role in tumor metastasis. *Cell* 117: 927–939, 2004
48. Greenburg G, Hay ED: Epithelia suspended in collagen gels can lose polarity and express characteristics of migrating mesenchymal cells. *J Cell Biol* 95: 333–339, 1982
49. Okada H, Danoff TM, Fischer A, Lopez-Guisa JM, Strutz F, Neilson EG: Identification of a novel cis-acting element for fibroblast-specific transcription of the FSP1 gene. *Am J Physiol* 275: F306–314, 1998
50. Le Hir M, Hegyi I, Cueni-Loffing D, Loffing J, Kaissling B: Characterization of renal interstitial fibroblast-specific protein 1/S100A4-positive cells in healthy and inflamed rodent kidneys. *Histochem Cell Biol* 123: 335–346, 2005
51. Inoue T, Plieth D, Venkov CD, Xu C, Neilson EG: Antibodies against macrophages that overlap in specificity with fibroblasts. *Kidney Int* 67: 2488–2493, 2005
52. Iwano M, Fischer A, Okada H, Plieth D, Xue C, Danoff TM, Neilson EG: Conditional abatement of tissue fibrosis using nucleoside analogs to selectively corrupt dna replication in transgenic fibroblasts. *Mol Ther* 3: 149–159, 2001
53. Strutz F, Muller GA, Neilson EG: Transdifferentiation: A new angle on renal fibrosis. *Exp Nephrol* 4: 267–270, 1996
54. Nishitani Y, Iwano M, Yamaguchi Y, Harada K, Nakatani K, Akai Y, Nishino T, Shiiki H, Kanauchi M, Saito Y, Neilson EG: Fibroblast-specific protein 1 is a specific prognostic marker for renal survival in patients with IgAN. *Kidney Int* 68: 1078–1085, 2005
55. Zeisberg M, Kalluri R: The role of epithelial-to-mesenchymal transition in renal fibrosis. *J Mol Med* 82: 175–181, 2004
56. Ng YY, Huang TP, Yang WC, Chen ZP, Yang AH, Mu W, Nikolic-Paterson DJ, Atkins RC, Lan HY: Tubular epithelial-myofibroblast transdifferentiation in progressive tubulointerstitial fibrosis in 5/6 nephrectomized rats. *Kidney Int* 54: 864–876, 1998
57. Faulkner JL, Szykalski LM, Springer F, Barnes JL: Origin of interstitial fibroblasts in an accelerated model of angiotensin II-induced renal fibrosis. *Am J Pathol* 167: 1193–1205, 2005
58. Ikeda Y, Jung YO, Kim H, Oda T, Lopez-Guisa J, Maruvada R, Diamond DL, Martin KJ, Wing D, Cai X, Eddy AA: Exogenous bone morphogenetic protein-7 fails to attenuate renal fibrosis in rats with overload proteinuria. *Nephron Exp Nephrol* 97: e123–e135, 2004
59. Yang J, Shultz RW, Mars WM, Wegner RE, Li Y, Dai C, Nejak K, Liu Y: Disruption of tissue-type plasminogen activator gene in mice reduces renal interstitial fibrosis in obstructive nephropathy. *J Clin Invest* 110: 1525–1538, 2002

60. Hruska KA, Guo G, Wozniak M, Martin D, Miller S, Liapis H, Loveday K, Klahr S, Sampath TK, Morrissey J: Osteogenic protein-1 prevents renal fibrogenesis associated with ureteral obstruction. *Am J Physiol Renal Physiol* 279: F130–F143, 2000
61. Zeisberg M, Hanai J, Sugimoto H, Mammoto T, Charytan D, Strutz F, Kalluri R: BMP-7 counteracts TGF-beta1-induced epithelial-to-mesenchymal transition and reverses chronic renal injury. *Nat Med* 9: 964–968, 2003
62. Yang J, Liu Y: Blockage of tubular epithelial to myofibroblast transition by hepatocyte growth factor prevents renal interstitial fibrosis. *J Am Soc Nephrol* 13: 96–107, 2002
63. Ohba K, Miyata Y, Koga S, Nishikido M, Kanetake H, Nazneen A, Razzaque MS, Taguchi T: Interstitial expression of heat-shock protein 47 correlates with capillary deposition of complement split product C4d in chronic allograft nephropathy. *Clin Transplant* 19: 810–816, 2005
64. Hughes S, Chan-Ling T: Characterization of smooth muscle cell and pericyte differentiation in the rat retina in vivo. *Invest Ophthalmol Vis Sci* 45: 2795–2806, 2004
65. Bjarnegard M, Enge M, Norlin J, Gustafsdottir S, Fredriksson S, Abramsson A, Takemoto M, Gustafsson E, Fassler R, Betsholtz C: Endothelium-specific ablation of PDGFB leads to pericyte loss and glomerular, cardiac and placental abnormalities. *Development* 131: 1847–1857, 2004