

- reexpression in Alport's mouse glomerular basement membranes. *Kidney Int* 63: 826–834, 2003
16. Cosgrove D, Meehan DT, Grunkemeyer JA, Kornak JM, Sayers R, Hunter WJ, Samuelson GC: Collagen COL4A3 knockout: A mouse model of autosomal Alport's syndrome. *Genes Dev* 10: 2981–2992, 1996
 17. Jarad G, Cunningham J, Shaw AS, Miner JH: Proteinuria precedes podocyte abnormalities in *Lamb2*^{-/-} mice, implicating the glomerular basement membrane as an albumin barrier. *J Clin Invest* 116: 2272–2279, 2006
 18. Miner JH, Li C: Defective glomerulogenesis in the absence of laminin alpha5 demonstrates a developmental role for the kidney glomerular basement membrane. *Dev Biol* 217: 278–289, 2000
 19. Kojima K, Davidovitas A, Poczewski H, Langer B, Uchida S, Nagy-Bojarski K, Hovorka A, Sedivy R, Kerjaschki D: Podocyte flattening and disorder of glomerular basement membrane are associated with splitting of dystroglycan-matrix interaction. *J Am Soc Nephrol* 15: 2079–2089, 2004
 20. Timpl R, Tisi D, Talts FJ, Andac Z, Sasaki T, Hohenester E: Structure and function of laminin LG modules. *Matrix Biol* 19: 309–317, 2000

See the related article, "Partial Rescue of Glomerular Laminin α 5 Mutation by Wild-Type Endothelial Cells Produce Hybrid Glomeruli," on pages 2285–2293.

Angiotensin-2 and Glomerular Proteinuria

Mark P. de Caestecker

Departments of Medicine, Cell and Developmental Biology, and Cancer Biology, Vanderbilt University School of Medicine, Nashville, Tennessee

J Am Soc Nephrol 18: 2217–2218, 2007.
doi: 10.1681/ASN.2007060650

The close spatial apposition between podocytes, fenestrated endothelia, and mesangial cells within the glomerulus has led to speculation that paracrine growth factors secreted by these cells are required to maintain structural and functional integrity of glomerular permselectivity in adults. This line of thinking has advanced the concept that disruption in the normal balance of these paracrine growth factors might give rise to proteinuric renal disease.

Angiotensin-2 is an antiangiogenic growth factor that is secreted by endothelial cells during periods of active vascular remodeling and opposes the proangiogenic effects of angiotensin-1 mediated through activation of the endothelial tyrosine kinase receptor Tie-2.^{1,2} Previous studies have shown that angiotensin-2 is expressed in developing glomeruli, where it is normally downregulated after birth³ but is upregulated in a variety of

experimental models of glomerular disease, including diabetes.^{4–6} As such, angiotensin-2 is a candidate growth factor that might play a role in destabilizing glomerular endothelia, causing a breakdown of glomerular permselectivity in proteinuric renal diseases.

In this issue of *JASN*, Davis *et al.*⁷ address this question using an inducible transgenic strategy to promote prolonged (5 to 10 wk) ectopic expression of angiotensin-2 in adult mouse podocytes. These mice develop low levels of nonselective proteinuria, indicating that angiotensin-2 has the capacity to modify glomerular permselectivity. These observations raise two important questions that warrant further discussion: How does angiotensin-2 cause proteinuria, and what is the significance of these findings for pathogenesis of glomerular disease?

Electron microscopic studies in these angiotensin-2–overexpressing mice demonstrate glomerular endothelial apoptosis, but there is no evidence of glomerular capillary collapse or foot process effacement. These findings are consistent with the role of angiotensin-2 in destabilizing endothelial cell integrity⁸ but raise questions about the mechanism of proteinuria.

The authors provide evidence that the slit diaphragm protein nephrin, an essential component of the glomerular permselectivity barrier,⁹ is downregulated in angiotensin-2–overexpressing mice. On the basis of the observation that proteinuria has been described in the absence of foot process effacement, the authors argue that these changes in the expression of nephrin may give rise to a defect in slit diaphragm function without inducing a structural abnormality in podocytes. This is certainly a possibility that might be confirmed by more detailed ultrastructural analysis of the slit diaphragm. However, an alternative possibility is that the primary defect in these mice results from loss of glomerular endothelial integrity.

This speculation is consistent with Davis's observations of endothelial cell apoptosis and that expression of the angiotensin-1/2 receptor Tie-2 is generally restricted to endothelial cells.¹⁰ Furthermore, despite the important focus on podocyte biology in the pathogenesis of proteinuric renal disease,⁹ there is emerging evidence that the specialized, fenestrated endothelia along the glomerular capillary also play a significant role in maintaining the charge selective barrier to proteinuria.^{11,12} In addition, it could be argued that the low levels of proteinuria observed in the angiotensin-2–overexpressing mice are more consistent with human microalbuminuria that is thought to reflect a primary defect in endothelial as opposed to glomerular epithelial function.¹³

The functional significance of these changes in angiotensin-2–overexpressing mice for glomerular pathology is even less clear cut. For example, it is uncertain whether mild proteinuria without evidence of structural abnormalities in glomerular architecture will give rise to progressive renal disease. Long-term studies using this mouse model would establish whether this is the case. More important, however, is that it is unclear whether persistent, isolated expression of angiotensin-2 in podocytes reflects the more complex environment of

Published online ahead of print. Publication date available at www.jasn.org.

Correspondence: Dr. Mark de Caestecker, Division of Nephrology, S-3223, Medical Center North, 1161 21st Street S., Nashville, TN 37232-2372. Phone: 615-343-2844; Fax: 615-343-2675; E-mail: mark.de.caestecker@vanderbilt.edu

Copyright © 2007 by the American Society of Nephrology

the glomerulus in disease states. There is evidence, for example, that glomerular expression of a key regulator of angiotensin-2 function, vascular endothelial growth factor A (VEGF-A), is upregulated in some glomerular diseases (notably diabetic nephropathy) and downregulated in others.¹⁴

This is a critical issue because angiotensin-2 has different effects on vascular remodeling and endothelial integrity depending on the coincident levels of VEGF-A.¹⁰ On this basis, we might expect increased levels of angiotensin-2 and VEGF-A in diabetic glomeruli to have different effects from those of angiotensin-2 overexpression in other glomerular diseases. These issues can be addressed only by evaluating the impact of decreasing angiotensin-2 expression in appropriate experimental models. However, these studies cannot be performed using germline *angiotensin-2* null mice, because these mice die within the first 2 weeks of life with severe defects in the lymphatic system.¹⁵ Studies in *angiotensin-2* heterozygous null mice might be informative if gene dosage has an impact on angiotensin-2 expression and function. Otherwise, we will have to wait for definitive studies in which the effect of temporally controlled, cell-specific deletion of *angiotensin-2* is evaluated experimentally. It may be some time before we obtain a definitive answer to these questions, as the necessary floxed *Ang-2* mutants have yet to be generated.

DISCLOSURES

None.

REFERENCES

- Maisonpierre PC, Suri C, Jones PF, Bartunkova S, Wiegand SJ, Radziejewski C, Compton D, McClain J, Aldrich TH, Papadopoulos N, Daly TJ, Davis S, Sato TN, Yancopoulos GD: Angiotensin-2, a natural antagonist for Tie2 that disrupts in vivo angiogenesis. *Science* 277: 55–60, 1997
- Fiedler U, Scharpfenecker M, Koidl S, Hegen A, Grunow V, Schmidt JM, Kriz W, Thurston G, Augustin HG: The Tie-2 ligand angiotensin-2 is stored in and rapidly released upon stimulation from endothelial cell Weibel-Palade bodies. *Blood* 103: 4150–4156, 2004
- Yuan HT, Suri C, Landon DN, Yancopoulos GD, Woolf AS: Angiotensin-2 is a site-specific factor in differentiation of mouse renal vasculature. *J Am Soc Nephrol* 11: 1055–1066, 2000
- Yuan HT, Tipping PG, Li XZ, Long DA, Woolf AS: Angiotensin correlates with glomerular capillary loss in anti-glomerular basement membrane glomerulonephritis. *Kidney Int* 61: 2078–2089, 2002
- Lu YH, Deng AG, Li N, Song MN, Yang X, Liu JS: Changes in angiotensin expression in glomeruli involved in glomerulosclerosis in rats with daunorubicin-induced nephrosis. *Acta Pharmacol Sin* 27: 579–587, 2006
- Rizkalla B, Forbes JM, Cao Z, Boner G, Cooper ME: Temporal renal expression of angiogenic growth factors and their receptors in experimental diabetes: Role of the renin-angiotensin system. *J Hypertens* 23: 153–164, 2005
- Davis B, Dei Cas A, Long DA, White KE, Hayward A, Ku CH, Woolf AS, Bilous R, Viberti G, Gnudi L: Podocyte-specific induced overexpression of angiotensin-2 causes proteinuria and apoptosis of glomerular endothelia. *J Am Soc Nephrol* 17: 2320–2329, 2007
- Scharpfenecker M, Fiedler U, Reiss Y, Augustin HG: The Tie-2 ligand angiotensin-2 destabilizes quiescent endothelium through an internal autocrine loop mechanism. *J Cell Sci* 118: 771–780, 2005
- Tryggvason K, Wartiovaara J: How does the kidney filter plasma? *Physiology (Bethesda)* 20: 96–101, 2005
- Eklund L, Olsen BR: Tie receptors and their angiotensin ligands are context-dependent regulators of vascular remodeling. *Exp Cell Res* 312: 630–641, 2006
- Camici M: Renal glomerular permselectivity and vascular endothelium. *Biomed Pharmacother* 59: 30–37, 2005
- Haraldsson B, Sorensson J: Why do we not all have proteinuria? An update of our current understanding of the glomerular barrier. *News Physiol Sci* 19: 7–10, 2004
- Amann K, Wanner C, Ritz E: Cross-talk between the kidney and the cardiovascular system. *J Am Soc Nephrol* 17: 2112–2119, 2006
- Schrijvers BF, Flyvbjerg A, De Zeeuw AS: The role of vascular endothelial growth factor (VEGF) in renal pathophysiology. *Kidney Int* 65: 2003–2017, 2004
- Gale NW, Thurston G, Hackett SF, Renard R, Wang Q, McClain J, Martin C, Witte C, Witte MH, Jackson D, Suri C, Campochiaro PA, Wiegand SJ, Yancopoulos GD: Angiotensin-2 is required for postnatal angiogenesis and lymphatic patterning, and only the latter role is rescued by Angiotensin-1. *Dev Cell* 3: 411–423, 2002

See the related article, "Podocyte-Specific Expression of Angiotensin-2 Causes Proteinuria and Apoptosis of Glomerular Endothelia," on pages 2320–2329.

Hemoglobin Variability in Dialysis Patients

K. Scott Brimble* and Catherine M. Clase*[†]

*Department of Medicine and [†]Department of Clinical Epidemiology and Biostatistics, McMaster University, Hamilton, Ontario, Canada

J Am Soc Nephrol 18: 2218–2220, 2007.

doi: 10.1681/ASN.2007060643

Hemoglobin variability is the extent to which multiple measured hemoglobin values differ from each other. Variability may be assessed within the same patient or between patients in a group; in the context of clinical practice, it is generally the variability within a patient that is important, whereas for quality assurance purposes, both variability within patients (an index of individual stability) and between patients (an index of the extent to which values differ between patients) may be relevant. As West *et al.* observe in the current issue of *JASN*,¹ the adjustment of epoetins in the management of anemia in renal disease, whether done by clinical judgment, the use of simple clinical decision rules, or a more complex computer program essentially follows the principles of a negative-feed-

Published online ahead of print. Publication date available at www.jasn.org.

Correspondence: Prof. Catherine M Clase, Department of Medicine, Associate Member, Department of Clinical Epidemiology and Biostatistics, McMaster University, 708-25 Charlton Avenue East, Hamilton, Ontario L8N 1Y2, Canada. Phone: 905-521-6094; Fax: 905-521-6153; E-mail: clase@mcmaster.ca

Copyright © 2007 by the American Society of Nephrology