

containing integrins in glomerular function? There has not been a detailed examination of kidneys in  $\alpha v$  null mice or in conditional knockouts of the gene encoding  $\alpha v$  integrin in any kidney lineages; therefore, it is possible some kidney defects have been missed. Conversely, much of the  $\alpha v$ -containing integrin in the glomerulus is thought to be  $\alpha v\beta 3$ , and  $\beta 3$  integrin null mice have relatively normal kidneys<sup>11</sup> that would argue against a developmental role for  $\alpha v\beta 3$  in kidneys. Borza *et al.* also demonstrated that  $\alpha 3\beta 1$  integrins bind the  $\alpha 3$ NC1 collagen domain and transdominantly inhibit adhesion mediated by  $\alpha v\beta 3$  integrin,<sup>12</sup> suggesting regulatory interplay between integrins on podocytes. In addition, some recent evidence posits an important role for  $\alpha v\beta 3$  integrin in kidney injury.<sup>11</sup> It is known that GPI-linked urokinase-type plasminogen activator receptors (uPAR) interact with several integrins that modulate their ligand-binding activities.<sup>13</sup> Wei *et al.*<sup>11</sup> showed that uPAR and  $\alpha v\beta 3$  integrin co-localize in podocyte foot processes and that uPAR-deficient mice are resistant to LPS-induced foot process effacement.  $\alpha v\beta 3$  integrin is among those that are modulated by uPAR, and activation of  $\alpha v\beta 3$  integrin in glomeruli (engaged by an activation-specific antibody) is decreased in uPAR-deficient mice.<sup>13</sup> Finally, expression of a constitutively active  $\beta 3$  integrin in podocytes results in proteinuria.

Together, the studies of Borza *et al.* and Wei *et al.* now indicate a role for  $\alpha v\beta 3$  integrin in glomerular disease and indicate a binding site for  $\alpha v\beta 3$  integrin in the KRGDS motif of human and primate  $\alpha 3$  type IV collagen. Although the latter study indicates a role for  $\alpha v\beta 3$  in glomerular disease even in the absence of this motif in rodents, this does not preclude a role for the motif in modulating glomerular function in humans or from its having a role in human kidney disease. Further studies will need to examine the phosphorylation of this site in various models of glomerular disease and the effects of mutagenizing this site to prevent phosphorylation or integrin binding. Genetic studies in animals are difficult because this motif is not found in rodents. Perhaps immortalized human podocytes can be used as an *in vitro* model for additional studies.

## DISCLOSURES

None.

## REFERENCES

1. Miner JH: Developmental biology of glomerular basement membrane components. *Curr Opin Nephrol Hypertens* 7: 13–19, 1998
2. Noakes PG, Miner JH, Gautam M, Cunningham JM, Sanes JR, Merlie JP: The renal glomerulus of mice lacking s-laminin/laminin beta 2: Nephrosis despite molecular compensation by laminin beta 1. *Nat Genet* 10: 400–406, 1995
3. Miner JH, Li C: Defective glomerulogenesis in the absence of laminin alpha5 demonstrates a developmental role for the kidney glomerular

- basement membrane. *Dev Biol* 217: 278–289, 2000
4. Kreidberg JA, Donovan MJ, Goldstein SL, Rennke H, Shepherd K, Jones RC, Jaenisch R: alpha3beta1 integrin has a crucial role in kidney and lung organogenesis. *Development* 122: 3537–3547, 1996
5. Kang JS, Wang XP, Miner JH, Morello R, Sado Y, Abrahamson DR, Borza DB: Loss of alpha3/alpha4(IV) collagen from the glomerular basement membrane induces a strain-dependent isoform switch to alpha5/alpha6(IV) collagen associated with longer renal survival in Col4a3-/- Alport mice. *J Am Soc Nephrol* 17: 1962–1969, 2006
6. Borza CM, Borza DB, Pedchenko V, Saleem MA, Mathieson PW, Sado Y, Hudson HM, Pozzi A, Saus J, Abrahamson DR, Zent R, Hudson BG: Human podocytes adhere to the KRGDS motif of the  $\alpha 3\alpha 4\alpha 5$  collagen IV network. *J Am Soc Nephrol* 19: 677–684, 2008
7. Raya A, Revert F, Navarro S, Saus J: Characterization of a novel type of serine/threonine kinase that specifically phosphorylates the human Goodpasture antigen. *J Biol Chem* 274: 12642–12649, 1999
8. Chen X, Moeckel G, Morrow JD, Cosgrove D, Harris RC, Fogo AB, Zent R, Pozzi A: Lack of integrin alpha1beta1 leads to severe glomerulosclerosis after glomerular injury. *Am J Pathol* 165: 617–630, 2004
9. Brooks PC, Clark RA, Cheresh DA: Requirement of vascular integrin alphavbeta3 for angiogenesis. *Science* 264: 569–571, 1994
10. Bader BL, Rayburn H, Crowley D, Hynes RO: Extensive vasculogenesis, angiogenesis, and organogenesis precede lethality in mice lacking all alphav integrins. *Cell* 95: 507–519, 1998
11. Wei C, Moller CC, Altintas MM, Li J, Schwarz K, Zacchigna S, Xie L, Henger A, Schmid H, Rastaldi MP, Cowan P, Kretzler M, Parrilla R, Bendayan M, Gupta V, Nikolic B, Kalluri R, Carmeliet P, Mundel P, Reiser J: Modification of kidney barrier function by the urokinase receptor. *Nat Med* 14: 55–63, 2008
12. Borza CM, Pozzi A, Borza DB, Pedchenko V, Hellmark T, Hudson BG, Zent R: Integrin alpha3beta1, a novel receptor for alpha3(IV) noncollagenous domain and a trans-dominant inhibitor for integrin alphavbeta3. *J Biol Chem* 281: 20932–20939, 2006
13. Chapman HA, Wei Y: Protease crosstalk with integrins: The urokinase receptor paradigm. *Thromb Haemost* 86: 124–129, 2001

See related article, "Human Podocytes Adhere to the KRGDS Motif of the  $\alpha 3\alpha 4\alpha 5$  Collagen IV Network," on pages 677–684.

## Are Podocytes Passive or Provocative in Proteinuric Glomerular Pathology?

Peter G. Tipping

Department of Medicine, Monash University, Victoria, Australia

*J Am Soc Nephrol* 19: 651–653, 2008.  
doi: 10.1681/ASN.2008020156

Podocytes are highly specialized epithelial cells with unique structure and function that are essential to maintenance of the

Published online ahead of print. Publication date available at [www.jasn.org](http://www.jasn.org).

**Correspondence:** Dr. Peter G. Tipping, Monash University, Department of Medicine, Monash Medical Centre, 246 Clayton Road, Clayton, Victoria, Australia. Phone: 011-61-39-594-5547; Fax: 011-61-39-594-6495; E-mail: [peter.tipping@med.monash.edu.au](mailto:peter.tipping@med.monash.edu.au)

Copyright © 2008 by the American Society of Nephrology

glomerular filtration barrier. Because of their pivotal role in glomerular filtration, their delicate fimbriated structure, and their limited ability for regeneration and repair, podocytes have been considered critical and vulnerable targets of glomerular injury. Disruption of the anatomic relationships between adjacent foot processes and between foot processes and the glomerular basement membrane (GBM) are among the earliest morphologic features of glomerular injury where proteinuria is a hallmark.

Podocytes are generally regarded as passive targets of both immune and nonimmune injury. In experimental models, they are highly susceptible to a variety of injurious agents, including complement, reactive oxygen species (ROS), and toxins such as puromycin aminonucleoside. Podocytes upregulate their expression of C5a receptors<sup>1</sup> and become the major target of complement-mediated immune injury in membranous nephropathy.<sup>2</sup> In Heymann nephritis, a rat model of human membranous nephritis, the combined insults of sublytic amounts of complement membrane attack complex and ROS induce podocyte injury and proteinuria.<sup>3</sup> Albuminuria in diabetic nephropathy results from nonimmune glomerular injury. The loss of  $\alpha 3\beta 1$  integrin anchoring of foot processes to the GBM followed by podocyte detachment and shedding into the urine is now recognized as an early pathologic feature in experimental<sup>4,5</sup> and human diabetes.<sup>4</sup> In puromycin aminonucleoside-induced nephrosis, used as a model of idiopathic nephrotic syndrome, direct oxidative mechanisms induce podocyte foot process effacement<sup>6</sup> and apoptosis<sup>7</sup> leading to proteinuria.

Recent studies, including that by Banas *et al.*<sup>8</sup> in this issue of *JASN*, suggest podocytes may aggravate immune and nonimmune glomerular injury and contribute to their own demise through expression of receptors linked to pathways that induce proinflammatory molecules. Banas *et al.* studied immune complex-initiated injury in thymic stromal lymphopoietin transgenic mice that develop cryoglobulinemia and membranoproliferative glomerulonephritis.<sup>8</sup> Expression of mRNA encoding TLR-1, -2, and -4 are markedly upregulated early in the development of disease, and their expression is further increased when the severity of glomerulonephritis is augmented by deletion of an inhibitory Fc receptor (FcRIIb). TLR-4 expression is found on podocytes in nephritic glomeruli, and *in vitro* studies using immortalized mouse podocytes demonstrate induction of chemokine expression in response to lipid A, the specific TLR-4-binding component of LPS ligand. Interestingly, fibrinogen, an endogenous TLR-4 ligand, prominently deposited in glomeruli in this model and exhibits a similar ability to induce chemokine production by podocytes. These studies demonstrate the potential of podocytes to contribute actively to recruitment of inflammatory cells and glomerular injury by upregulating TLR-4 and production of proinflammatory chemokines in response to stimulation by either endogenous or exogenous TLR-4 ligands.

Although the studies by Banas *et al.* did not investigate

the separate *in vivo* contributions to renal injury of TLR-4 expression by podocytes and leukocytes, a recent study by Brown *et al.*<sup>9</sup> addressed this issue using an acute anti-GBM antibody/lipid A-induced model of glomerular injury. Glomerular TLR-4 expression, rather than expression on leukocytes, is principally required for development of proteinuria in this model. TLR-4 expression on podocytes and mesangial cells and involvement of CXC chemokines in TLR-4-dependent injury was demonstrated, adding further weight to the argument that intrinsic renal cells, including podocytes, amplify glomerular injury through TLR-4-dependent pathways. Glomerular epithelial cells produce the proinflammatory cytokines TNF- $\alpha$  and IL-6 after LPS stimulation *in vitro*.<sup>10</sup> The ability of intrinsic renal cells to amplify inflammatory glomerular injury by their capacity to produce cytokines, particularly TNF- $\alpha$ , has been previously reported in murine anti-GBM nephritis,<sup>11</sup> and in human membranous nephritis, podocytes are a prominent source of TNF- $\alpha$ .<sup>12</sup>

Rapid induction of B7-1 (CD80) on podocytes and development of proteinuria in mice after administration of LPS and puromycin provides more evidence of the ability of podocytes to acquire a proinflammatory phenotype.<sup>13</sup> B7-1 is better known as a co-stimulatory molecule involved in antigen presentation to T cells, but its contribution to development of proteinuria in this study was independent of T cell participation. LPS-stimulated TLR-4 induces upregulation of B7-1 expression by podocytes *in vitro* and results in rearrangement of their actin cytoskeleton. In light of the work by Banas *et al.*,<sup>8</sup> induction of chemokine expression may have been expected, but this end point was not reported. Higher B7-1 expression on podocytes is also associated with more severe World Health Organization classes of glomerulonephritis in human lupus nephritis and more severe proteinuria and renal injury in murine lupus, suggesting involvement in inflammatory cell recruitment.

Podocytes amplify both immune and nonimmune glomerular injury by their capacity to produce and release ROS. Increased production of ROS by podocytes is also implicated in various experimental proteinuric diseases, including puromycin nephrosis,<sup>14</sup> Heymann nephritis,<sup>15</sup> Mpv17 gene-inactivated mice (a model of focal segmental sclerosis),<sup>16</sup> type 1 diabetes,<sup>17</sup> and human membranous nephropathy.<sup>18</sup> Podocytes express chemokine receptors and produce ROS in response to chemokine stimulation *in vitro*.<sup>19</sup> In turn, ROS stimulate podocyte production of the proinflammatory cytokine GM-CSF,<sup>20</sup> induce podocyte cell-cycle checkpoint proteins,<sup>14</sup> induce apoptosis in podocytes,<sup>17</sup> and directly injure the GBM.<sup>3,15</sup>

In diabetic nephropathy, evidence is emerging that acquisition of proinflammatory capacities by podocytes contributes to perpetuation of injury. Both hemodynamic and metabolic stimuli contribute to these changes in podocyte phenotype. For example, mechanical stretch can induce angiotensin II (AngII) expression in cultured podocytes.<sup>21</sup> An-

gII stimulates podocyte vascular endothelial growth factor production, which, in an autocrine manner together with TGF- $\beta$ , stimulates production of  $\alpha$ 5(IV) collagen.<sup>22</sup> Targeted overexpression of AngII in podocytes leads to proteinuria and glomerulosclerosis in rats.<sup>23</sup> Exposure of cultured podocytes to high glucose concentrations increases ROS production and podocyte hypertrophy,<sup>24</sup> modulates production of matrix metalloproteases and  $\alpha$ 5(IV) collagen,<sup>25</sup> and stimulates 12-lipoxygenase and p38 mitogen-activated protein kinase signaling pathways.<sup>26</sup> The role of podocytes in diabetic nephropathy has been covered extensively in recent reviews.<sup>27,28</sup>

The study by Banas *et al.*<sup>8</sup> adds to accumulating evidence that podocytes provide can acquire functions that augment both immune and nonimmune glomerular injury in response to various pathogenic stimuli. This evidence challenges the view that podocytes are passive participants and merely a common final pathway or target of injury in proteinuric renal diseases.

## REFERENCES

- Abe K, Miyazaki M, Koji T, Furusu A, Nakamura-Kurashige T, Nishino T, Ozono Y, Harada T, Sakai H, Kohno S: Enhanced expression of complement C5a receptor mRNA in human diseased kidney assessed by in situ hybridization. *Kidney Int* 60: 137–146, 2001
- Couser WG, Nangaku M: Cellular and molecular biology of membranous nephropathy. *J Nephrol* 19: 699–705, 2006
- Kerjaschki D, Neale TJ: Molecular mechanisms of glomerular injury in rat experimental membranous nephropathy (Heymann nephritis). *J Am Soc Nephrol* 7: 2518–2526, 1996
- Chen HC, Chen CA, Guh JY, Chang JM, Shin SJ, Lai YH: Altering expression of alpha3beta1 integrin on podocytes of human and rats with diabetes. *Life Sci* 67: 2345–2353, 2000
- Regoli M, Bendayan M: Alterations in the expression of the alpha 3 beta 1 integrin in certain membrane domains of the glomerular epithelial cells (podocytes) in diabetes mellitus. *Diabetologia* 40: 15–22, 1997
- Inokuchi S, Shirato I, Kobayashi N, Koide H, Tomino Y, Sakai T: Re-evaluation of foot process effacement in acute puromycin aminonucleoside nephrosis. *Kidney Int* 50: 1278–1287, 1996
- Sanwal V, Pandya M, Bhaskaran M, Franki N, Reddy K, Ding G, Kapasi A, Valderrama E, Singhal PC: Puromycin aminonucleoside induces glomerular epithelial cell apoptosis. *Exp Mol Pathol* 70: 54–64, 2001
- Banas MC, Banas B, Hudkins KL, Wietecha TA, Iyoda M, Bock E, Hauser P, Pippin JW, Shankland SJ, Smith KD, Stoelcker B, Liu G, Gröne H-J, Krämer BK, Alpers CE: TLR4 links podocytes with the innate immune system to mediate glomerular injury. *J Am Soc Nephrol* 19: 704–713, 2008
- Brown HJ, Lock HR, Wolfs TG, Buurman WA, Sacks SH, Robson MG: Toll-like receptor 4 ligation on intrinsic renal cells contributes to the induction of antibody-mediated glomerulonephritis via CXCL1 and CXCL2. *J Am Soc Nephrol* 18: 1732–1739, 2007
- Hughes AK, Stricklett PK, Kohan DE: Shiga toxin-1 regulation of cytokine production by human glomerular epithelial cells. *Nephron* 88: 14–23, 2001
- Timoshanko JR, Sedgwick JD, Holdsworth SR, Tipping PG: Intrinsic renal cells are the major source of tumor necrosis factor contributing to renal injury in murine crescentic glomerulonephritis. *J Am Soc Nephrol* 14: 1785–1793, 2003
- Neale TJ, Ruger BM, Macaulay H, Dunbar PR, Hasan Q, Bourke A, Murray-McIntosh RP, Kitching AR: Tumor necrosis factor-alpha is expressed by glomerular visceral epithelial cells in human membranous nephropathy. *Am J Pathol* 146: 1444–1454, 1995
- Reiser J, von Gersdorff G, Loos M, Oh J, Asanuma K, Giardino L, Rastaldi MP, Calvaresi N, Watanabe H, Schwarz K, Faul C, Kretzler M, Davidson A, Sugimoto H, Kalluri R, Sharpe AH, Kreidberg JA, Mundel P: Induction of B7-1 in podocytes is associated with nephrotic syndrome. *J Clin Invest* 113: 1390–1397, 2004
- Marshall CB, Pippin JW, Krofft RD, Shankland SJ: Puromycin aminonucleoside induces oxidant-dependent DNA damage in podocytes in vitro and in vivo. *Kidney Int* 70: 1962–1973, 2006
- Neale TJ, Ullrich R, Ojha P, Poczewski H, Verhoeven AJ, Kerjaschki D: Reactive oxygen species and neutrophil respiratory burst cytochrome b558 are produced by kidney glomerular cells in passive Heymann nephritis. *Proc Natl Acad Sci U S A* 90: 3645–3649, 1993
- Binder CJ, Weiher H, Exner M, Kerjaschki D: Glomerular overproduction of oxygen radicals in Mpv17 gene-inactivated mice causes podocyte foot process flattening and proteinuria: A model of steroid-resistant nephrosis sensitive to radical scavenger therapy. *Am J Pathol* 154: 1067–1075, 1999
- Susztak K, Raff AC, Schiffer M, Böttinger EP: Glucose-induced reactive oxygen species cause apoptosis of podocytes and podocyte depletion at the onset of diabetic nephropathy. *Diabetes* 55: 225–233, 2006
- Grone HJ, Walli AK, Grone EF: The role of oxidatively modified lipoproteins in lipid nephropathy. *Contrib Nephrol* 120: 160–175, 1997
- Huber TB, Reinhardt HC, Exner M, Burger JA, Kerjaschki D, Saleem MA, Pavenstädt H: Expression of functional CCR and CXCR chemokine receptors in podocytes. *J Immunol* 168: 6244–6252, 2002
- Greiber S, Muller B, Daemisch P, Pavenstädt H: Reactive oxygen species alter gene expression in podocytes: Induction of granulocyte macrophage-colony-stimulating factor. *J Am Soc Nephrol* 13: 86–95, 2002
- Durvasula RV, Petermann AT, Hiromura K, Blonski M, Pippin J, Mundel P, Pichler R, Griffin S, Couser WG, Shankland SJ: Activation of a local tissue angiotensin system in podocytes by mechanical strain. *Kidney Int* 65: 30–39, 2004
- Chen S, Lee JS, Iglesias-de la Cruz MC, Wang A, Izquierdo-Lahuerta A, Gandhi NK, Danesh FR, Wolf G, Ziyadeh FN: Angiotensin II stimulates alpha3(IV) collagen production in mouse podocytes via TGF-beta and VEGF signalling: Implications for diabetic glomerulopathy. *Nephrol Dial Transplant* 20: 1320–1328, 2005
- Hoffmann S, Podlich D, Hahnel B, Kriz W, Gretz N: Angiotensin II type 1 receptor overexpression in podocytes induces glomerulosclerosis in transgenic rats. *J Am Soc Nephrol* 15: 1475–1487, 2004
- Kim NH, Rincon-Choles H, Bhandari B, Choudhury GG, Abboud HE, Gorin Y: Redox dependence of glomerular epithelial cell hypertrophy in response to glucose. *Am J Physiol Renal Physiol* 290: F741–F751, 2006
- Bai Y, Wang L, Li Y, Liu S, Li J, Wang H, Huang H: High ambient glucose levels modulates the production of MMP-9 and alpha5(IV) collagen by cultured podocytes. *Cell Physiol Biochem* 17: 57–68, 2006
- Kang SW, Natarajan R, Shahed A, Nast CC, LaPage J, Mundel P, Kashtan C, Adler SG: Role of 12-lipoxygenase in the stimulation of p38 mitogen-activated protein kinase and collagen alpha5(IV) in experimental diabetic nephropathy and in glucose-stimulated podocytes. *J Am Soc Nephrol* 14: 3178–3187, 2003
- Marshall SM: The podocyte: A potential therapeutic target in diabetic nephropathy? *Curr Pharm Des* 13: 2713–2720, 2007
- Reddy GR, Kotlyarevska K, Ransom RF, Menon RK: The podocyte and diabetes mellitus: Is the podocyte the key to the origins of diabetic nephropathy? *Curr Opin Nephrol Hypertens* 17: 32–36, 2008

See related article, "TLR4 Links Podocytes with the Innate Immune System to Mediate Glomerular Injury," on pages 704–713.