

- Urinary volume, water and recurrences in idiopathic calcium nephrolithiasis: A 5-year randomized prospective study. *J Urol* 155: 839–843, 1996
7. Valtin H: "Drink at least eight glasses of water a day." Really? Is there scientific evidence for "8 x 8"? *Am J Physiol Regul Integr Comp Physiol* 283: R993–R1004, 2002
 8. Anastasio P, Cirillo M, Spitali L, Frangiosa A, Pollastro RM, De Santo NG: Level of hydration and renal function in healthy humans. *Kidney Int* 60: 748–756, 2001
 9. Spigt MG, Knottnerus JA, Westerterp KR, Olde Rikkert MG, Schayck CP: The effects of 6 months of increased water intake on blood sodium, glomerular filtration rate, blood pressure, and quality of life in elderly (aged 55–75) men. *J Am Geriatr Soc* 54: 438–443, 2006
 10. Spigt M, van Schayck O, Knipschild P, Westerterp K, van de Beek C, van Kerrebroeck P, Pel J, van Mastrigt R, Knottnerus A: Is it possible to improve elderly male bladder function by having them drink more water? A randomized trial of effects of increased fluid intake/urine output on male lower urinary tract function. *Urology* 68: 1031–1036, 2006
 11. Adroque HJ, Madias NE: Sodium and potassium in the pathogenesis of hypertension. *N Engl J Med* 356: 1966–1978, 2007
 12. Shafiee MA, Charest AF, Cheema-Dhadli S, Glick DN, Napolova O, Roozbeh J, Semenova E, Sharman A, Halperin ML: Defining conditions that lead to the retention of water: The importance of the arterial sodium concentration. *Kidney Int* 67: 613–621, 2005
 13. Claybaugh JR, Sato AK, Crosswhite LK, Hassell LH: Effects of time of day, gender, and menstrual cycle phase on the human response to a water load. *Am J Physiol Regul Integr Comp Physiol* 279: R966–R973, 2000
 14. Lappalainen R, Mennen L, van Weert L, Mykkanen H: Drinking water with a meal: A simple method of coping with feelings of hunger, satiety and desire to eat. *Eur J Clin Nutr* 47: 815–819, 1993
 15. Rolls BJ, Castellanos VH, Halford JC, Kilara A, Panyam D, Pelkman CL, Smith GP, Thorwart ML: Volume of food consumed affects satiety in men. *Am J Clin Nutr* 67: 1170–1177, 1998
 16. Rolls BJ, Bell EA, Thorwart ML: Water incorporated into a food but not served with a food decreases energy intake in lean women. *Am J Clin Nutr* 70: 448–455, 1999
 17. Boschmann M, Steiniger J, Hille U, Tank J, Adams F, Sharma AM, Klaus S, Luft FC, Jordan J: Water-induced thermogenesis. *J Clin Endocrinol Metab* 88: 6015–6019, 2003
 18. Boschmann M, Steiniger J, Franke G, Birkenfeld AL, Luft FC, Jordan J: Water drinking induces thermogenesis through osmosensitive mechanisms. *J Clin Endocrinol Metab* 92: 3334–3337, 2007
 19. Bergeron MF, Waller JL, Marinik EL: Voluntary fluid intake and core temperature responses in adolescent tennis players: Sports beverage versus water. *Br J Sports Med* 40: 406–410, 2006
 20. Brown CM, Dulloo AG, Montani JP: Water-induced thermogenesis reconsidered: The effects of osmolality and water temperature on energy expenditure after drinking. *J Clin Endocrinol Metab* 91: 3598–3602, 2006
 21. Brown CM, Barberini L, Dulloo AG, Montani JP: Cardiovascular responses to water drinking: Does osmolality play a role? *Am J Physiol Regul Integr Comp Physiol* 289: R1687–R1692, 2005
 22. Lu CC, Diedrich A, Tung CS, Paranjape SY, Harris PA, Byrne DW, Jordan J, Robertson D: Water ingestion as prophylaxis against syncope. *Circulation* 108: 2660–2665, 2003
 23. Chan J, Knutsen SF, Blix GG, Lee JW, Fraser GE: Water, other fluids, and fatal coronary heart disease: The Adventist health study. *Am J Epidemiol* 155: 827–833, 2002
 24. Altieri A, La Vecchia C, Negri E: Fluid intake and risk of bladder and other cancers. *Eur J Clin Nutr* 57[Suppl 2]: S59–S68, 2003
 25. Blau JN, Kell CA, Sperling JM: Water-deprivation headache: A new headache with two variants. *Headache* 44: 79–83, 2004
 26. Spigt MG, Kuijper EC, Schayck CP, Troost J, Knipschild PG, Linssen VM, Knottnerus JA: Increasing the daily water intake for the prophylactic treatment of headache: A pilot trial. *Eur J Neurol* 12: 715–718, 2005
 27. Wipke-Tevis DD, Williams DA: Effect of oral hydration on skin microcirculation in healthy young and midlife and older adults. *Wound Repair Regen* 15: 174–185, 2007

See related article, "Impact of Solute Intake on Urine Flow and Water Excretion," on pages 1076–1078.

Integrins, Extracellular Matrix, and Terminal Differentiation of Renal Epithelial Cells

Ambra Pozzi*^{†‡} and Roy Zent*^{†§}

Departments of *Medicine, Division of Nephrology, [†]Cancer Biology, and [§]Cell Biology, Vanderbilt University Medical Center, and [‡]Department of Medicine, Veterans Affairs Hospital, Nashville, Tennessee

J Am Soc Nephrol 19: 1043–1044, 2008.
doi: 10.1681/ASN.2008040370

The mechanism whereby epithelial cells terminally differentiate is an active area of investigation. One potential interface is the spatial and temporal expression of the transmembrane receptors known as integrins and the extracellular matrix (ECM) proteins to which they bind. In this issue of *JASN*, Vijayakumar *et al.*¹ propose that activation of integrin $\alpha\beta1$ causes synthesis and deposition of hensin, an ECM protein that forms 50- to 100-nm-long fibers composed of several fibrils. Upon polymerization and deposition into the ECM, hensin binds to $\alpha6$ -containing integrins, a key step in mediating the conversion of epithelial cells to a cuboid-like phenotype capable of apical endocytosis. These novel studies suggest that integrin–hensin interactions play an important role in the terminal differentiation of intercalated cells of the collecting duct.

The collecting system of the kidney is derived from the ureteric bud, which undergoes multiple iterations of branching morphogenesis followed by a phase of growth, maturation, and differentiation.² Many mechanisms regulate this branching and tubular expansion. Multiple transcription factors, growth factors, ECM proteins, and various cognate receptors play a critical role in these processes. Less information is available on the moieties that halt collecting duct growth and induce terminal differentiation. A number of transcription factors modulate the terminal differentiation of epithelial cells in

Published online ahead of print. Publication date available at www.jasn.org.

Correspondence: Dr. Ambra Pozzi or Dr. Roy Zent, Medical Center North, C3210, Department of Medicine, Division of Nephrology, Vanderbilt University, Nashville, TN 37232. Phone: 615-322-4632; Fax: 615-322-4690; E-mail: ambra.pozzi@vanderbilt.edu or roy.zent@vanderbilt.edu

Copyright © 2008 by the American Society of Nephrology

various nephron segments. In this context, the DNA-binding protein/tumor suppressor p53 is one such factor, and its stabilization by phosphorylation and acetylation enhances the transcription of genes associated with terminal differentiation, including aquaporin 2 and the Na⁺/K⁺ ATPase.^{3,4} In addition, p53 cooperates with at least two other transcription factors, cAMP response element-binding protein and Kruppel-like factor 4, to mediate its transcriptional regulation.⁵ Another example of transcriptional regulation of collecting duct differentiation is the forkhead transcription factor Foxi1, which mediates differentiation of intercalated cells from precursor epithelial cells. When Foxi1 is deleted in mice, they exhibit impaired expression of intercalated cell-specific genes such as Pendrin, H⁺-ATPase, and AE1, with no altered expression of principal cell markers.⁶ Thus, transcriptional processes or factors/co-factors that regulate the nuclear chromatin surrounding transcription are critical for modulating the expression of genes responsible for induction and maintenance of a differentiated cell phenotype.

Among ECM proteins, hensin modulates terminal differentiation by favoring the conversion of β to α intercalated cells through a process that requires polymerization of hensin by galectin 3.^{7,8} A key question is how hensin might regulate this terminal differentiation of intercalated cells. Although the answer is not clear, some clues are provided by Vijayakumar *et al.*¹ In this study, the authors show that polymerized hensin signals intercalated cells *in vitro* through $\alpha 6$ -containing integrins. Although they were unable to delineate clearly which $\alpha 6$ -containing integrin ($\alpha 6\beta 1$ and/or $\alpha 6\beta 4$) mediates this signaling, blocking anti- $\beta 1$ antibody experiments suggest the hensin-mediated signal is integrin $\alpha 6\beta 1$ independent, which leaves integrin $\alpha 6\beta 4$.

Integrin $\alpha 6\beta 4$ is unique among the 24 known mammalian integrins, because the $\beta 4$ subunit cytoplasmic tail is composed of 1017 amino acids and has distinctive cytoskeletal and signaling functions.⁹ Furthermore, the $\beta 4$ subunit is unique because it promotes the assembly of hemidesmosomes, specialized structures present in epithelial cells that create stable adhesion by connecting the intracellular keratin cytoskeleton to basement membrane.¹⁰ Thus, $\alpha 6\beta 4$ not only performs a major adhesive function in cells but also promotes polarization, a final event in epithelial cell differentiation. The highly polarized and tight junction epithelial cells found within the medulla of the kidney certainly benefit from such structures. Although the best described ligand for integrin $\alpha 6\beta 4$ is laminin-332 (also known as laminin-5), it is possible that hensin is also a ligand for this receptor. Alternative explanations could be that hensin interacts with laminin-332, thereby enhancing its interaction with integrin $\alpha 6\beta 4$, or it might induce laminin-332 expression directly.

The mechanisms whereby integrin $\alpha 6\beta 4$ regulates the terminal differentiation of intercalated cells are purely speculative at present. Hemidesmosome formation in epithelial cells along the collecting duct might favor not only cell polarization but also resistance to apoptosis. In this context, nonpolarized cells are sensitive to induction of apoptosis, whereas polarized cells

are resistant with ligation of the $\beta 4$ integrin.¹¹ It is also conceivable that ligation of integrin $\alpha 6\beta 4$ by hensin induces the activation of specific integrin $\alpha 6\beta 4$ -mediated signaling pathways, or the $\beta 4$ subunit works as an “adaptor” for growth factor receptor signaling.⁹ Both of these pathways might alter the transcription of genes required for the terminal differentiation of intercalated cells.

Although hensin is the first ECM protein shown to regulate terminal differentiation of intercalated cells within the collecting duct, it is highly likely that other ECM proteins along with basement membranes also regulate terminal differentiation of epithelial cells in different segments of the nephron and collecting system. Thus, the novel finding of Vijayakumar *et al.* suggests a critical role for the spatial/temporal expression of ECM proteins and their receptors in segmental renal function.

DISCLOSURES

None.

REFERENCES

- Vijayakumar S, Erdjument-Bromage H, Tempst P, Al-Awqati Q: Role of integrins in the assembly and function of hensin in intercalated cells. *J Am Soc Nephrol* 19: 1079–1091, 2008
- Cebrian C, Borodo K, Charles N, Herzlinger DA: Morphometric index of the developing murine kidney. *Dev Dyn* 231: 601–608, 2004
- Saifudeen Z, Dipp S, El-Dahr SS: A role for p53 in terminal epithelial cell differentiation. *J Clin Invest* 109: 1021–1030, 2002
- El-Dahr SS, Aboudehen K, Saifudeen Z: Transcriptional control of terminal nephron differentiation. *Am J Physiol Renal Physiol* 2008, in press
- Saifudeen Z, Dipp S, Fan H, El-Dahr SS: Combinatorial control of the bradykinin B2 receptor promoter by p53, CREB, KLF-4, and CBP: Implications for terminal nephron differentiation. *Am J Physiol Renal Physiol* 288: F899–F909, 2005
- Blomqvist SR, Vidarsson H, Fitzgerald S, Johansson BR, Ollerstam A, Brown R, Persson AE, Bergstrom GG, Enerback S: Distal renal tubular acidosis in mice that lack the forkhead transcription factor Foxi1. *J Clin Invest* 113: 1560–1570, 2004
- Al-Awqati Q, Vijayakumar S, Takito J: Terminal differentiation of epithelia from trophoctoderm to the intercalated cell: the role of hensin. *J Am Soc Nephrol* 14[Suppl 1]: S16–S21, 2003
- Hikita C, Vijayakumar S, Takito J, Erdjument-Bromage H, Tempst P, Al-Awqati Q: Induction of terminal differentiation in epithelial cells requires polymerization of hensin by galectin 3. *J Cell Biol* 151: 1235–1246, 2000
- Giancotti FG: Targeting integrin beta4 for cancer and anti-angiogenic therapy. *Trends Pharmacol Sci* 28: 506–511, 2007
- Litjens SH, de Pereda JM, Sonnenberg A: Current insights into the formation and breakdown of hemidesmosomes. *Trends Cell Biol* 16: 376–383, 2006
- Weaver VM, Lelievre S, Lakins JN, Chrenek MA, Jones JC, Giancotti F, Werb Z, Bissell MJ: Beta4 integrin-dependent formation of polarized three-dimensional architecture confers resistance to apoptosis in normal and malignant mammary epithelium. *Cancer Cell* 2: 205–216, 2002

See related article, “Role of Integrins in the Assembly and Function of Hensin in Intercalated Cells,” on pages 1079–1091.