

6. Huber TB, Kottgen M, Schilling B, Walz G, Benzing T: Interaction with podocin facilitates nephrin signaling. *J Biol Chem* 276: 41543–41546, 2001
7. Huber TB, Simons M, Hartleben B, Sernetz L, Schmidts M, Gundlach E, Saleem MA, Walz G, Benzing T: Molecular basis of the functional podocin-nephrin complex: Mutations in the NPHS2 gene disrupt nephrin targeting to lipid raft microdomains. *Hum Mol Genet* 12: 3397–3405, 2003
8. Huber TB, Schermer B, Muller RU, Hohne M, Bartram M, Calixto A, Hagmann H, Reinhardt C, Koos F, Kunzelmann K, Shirokova E, Krautwurst D, Harteneck C, Simons M, Pavenstadt H, Kerjaschki D, Thiele C, Walz G, Chalfie M, Benzing T: Podocin and MEC-2 bind cholesterol to regulate the activity of associated ion channels. *Proc Natl Acad Sci U S A* 103: 17079–17086, 2006
9. Hinkes BG, Mucha B, Vlangos CN, Gbadegesin R, Liu J, Hasselbacher K, Hangan D, Ozaltin F, Zenker M, Hildebrandt F: Nephrotic syndrome in the first year of life: Two thirds of cases are caused by mutations in 4 genes (NPHS1, NPHS2, WT1, and LAMB2). *Pediatrics* 119: e907–e919, 2007
10. Franceschini N, North KE, Kopp JB, McKenzie L, Winkler C: NPHS2 gene, nephrotic syndrome and focal segmental glomerulosclerosis: A HuGE review. *Genet Med* 8: 63–75, 2006
11. Roselli S, Heidet L, Sich M, Henger A, Kretzler M, Gubler MC, Antignac C: Early glomerular filtration defect and severe renal disease in podocin-deficient mice. *Mol Cell Biol* 24: 550–560, 2004
12. Philippe A, Weber S, Esquivel EL, Houbron C, Hamard G, Ratelade J, Kriz W, Schaefer F, Gubler MC, Antignac C: A missense mutation in podocin leads to early and severe renal disease in mice. *Kidney Int* 73: 1038–1047, 2008
13. Mollet G, Ratelade J, Boyer O, Muda AO, Morisset L, Lavin TA, Kitzis D, Dallman MJ, Bugeon L, Hubner N, Gubler MC, Antignac C, Esquivel EL: Podocin inactivation in mature kidneys causes focal segmental glomerulosclerosis and nephrotic syndrome. *J Am Soc Nephrol* 20: 2181–2189, 2009

See related article, "Podocin Inactivation in Mature Kidneys Causes Focal Segmental Glomerulosclerosis and Nephrotic Syndrome," on pages 2181–2189.

## More Evidence that Cystatin C Predicts Mortality Better than Creatinine

Peter P. Reese and Harold I. Feldman

Department of Medicine, Renal Division, Center for Clinical Epidemiology and Biostatistics, and Leonard Davis Institute, University of Pennsylvania, Philadelphia, Pennsylvania

*J Am Soc Nephrol* 20: 2088–2090, 2009.  
doi: 10.1681/ASN.2009080832

The main clinical benefit of estimating renal function in the general population is to identify and treat patients at risk for

Published online ahead of print. Publication date available at [www.jasn.org](http://www.jasn.org).

**Correspondence:** Dr. Peter P. Reese, Renal Electrolyte and Hypertension Division, Department of Medicine, University of Pennsylvania School of Medicine, 1 Founders, 3400 Spruce Street, Philadelphia, PA 19104. Phone: 215-662-7934; Fax: 215-615-0349; E-mail: [peter.reese@uphs.upenn.edu](mailto:peter.reese@uphs.upenn.edu)

Copyright © 2009 by the American Society of Nephrology

developing ESRD, cardiovascular events, and death.<sup>1</sup> Unfortunately, the most widely used equations to estimate GFR are limited, in part, by their reliance on serum creatinine as the filtration marker (estimated GFR [eGFR]). The Modification of Diet in Renal Disease (MDRD) equation, for example, overestimates renal function among individuals with low muscle mass and underestimates it among those with GFR >60 ml/min per 1.73 m<sup>2</sup>.<sup>2–4</sup> The use of serum cystatin C, a cysteine protease inhibitor that is freely filtered by the glomerulus, has potential advantages over creatinine as a filtration marker in that its production is not dependent on muscle mass.<sup>2</sup> As a result, cystatin C offers opportunities to estimate GFR more accurately than creatinine-based equations and additionally may predict worsening kidney function even when the GFR is actually near the normal range.<sup>5</sup>

Beyond the relationship to measured GFR, cystatin-C–derived eGFR (ecGFR) was a better predictor of mortality than creatinine-based estimates in a study of elderly individuals<sup>6</sup>; however, the ability to generalize these findings to a younger and broader set of individuals is unknown. In this issue of *JASN*, Astor *et al.*<sup>7</sup> extend previous observations and demonstrate convincingly that ecGFR predicts mortality more accurately than the MDRD equation in a sample from the general population in the United States. If replicated, then these findings should spur trials to determine whether the use of cystatin C improves patient outcomes by identifying those with an elevated risk for death, both within and outside the setting of diagnosed chronic kidney disease (CKD).

Astor *et al.*<sup>7</sup> focus on the relationship of ecGFR to mortality, but it is instructive to examine the challenges of using creatinine-based equations to assess renal function—the original reason for interest in cystatin C as a diagnostic test. The accuracy of a creatinine-derived eGFR and its value for detecting progressive kidney disease depends on the patient's level of renal function and associated level of proteinuria.<sup>1</sup> Patients with an eGFR <30 ml/min per 1.73 m<sup>2</sup> using MDRD are at substantial risk for later developing ESRD, but among patients with less severely diminished eGFR (45 to 90 ml/min per 1.73 m<sup>2</sup>), the prognostic value of an eGFR is limited, particularly when albuminuria is absent. For instance, follow-up of individuals who participated in the Multiple Risk Factor Intervention Trial (MRFIT) revealed that a subset with eGFR <60 (mean 55 ml/min per 1.73 m<sup>2</sup>) have only a 5.6% absolute risk for ESRD over 25 yr. In comparison, among participants with eGFR 60 to 75 ml/min per 1.73 m<sup>2</sup>, approximately 5.7% of those with 1+ proteinuria on urine dipstick evaluation and 17.7% of those with 2+ proteinuria develop ESRD.<sup>8</sup> The creatinine-based eGFR alone, therefore, has limited utility in predicting which patients with mildly diminished eGFR and no proteinuria will experience renal deterioration in the future.<sup>9</sup>

There are theoretical reasons to believe that cystatin C is a superior filtration marker compared with creatinine,<sup>10</sup> however, a comparison by Stevens *et al.*<sup>11</sup> of the performance characteristics of the MDRD equation with a cystatin C–based estimate in a large pooled population of patients with established

CKD revealed that cystatin C offers only modest improvement in accuracy. The performance of cystatin C–based measures to estimate renal function in the general population or among those with mild reductions in kidney function has not been explored fully.

Although eGFR derived from creatinine is also thought to be an important risk factor for mortality,<sup>12</sup> previous studies indicated cystatin C may be a powerful predictor of death independent of creatinine. In a cohort of 4637 community-dwelling elderly patients, Shlipak *et al.*<sup>6</sup> demonstrated a J-shaped relationship between the eGFR using the MDRD equation and mortality: Diminishing eGFR was associated with a graded increase in the risk for death, with this relationship reversing direction at higher levels of renal function such that patients with eGFR >74 ml/min per 1.73 m<sup>2</sup> had a higher (albeit NS) adjusted rate of mortality compared with patients with an eGFR from 66 to 74 ml/min per 1.73 m<sup>2</sup>. Without an apparent biologic basis for the relationship between greater renal function and a greater rate of death, this J-shaped relationship has been explained by the overestimation of renal function by creatinine-based equations among patients with more comorbidities and muscle wasting. By contrast, in the same cohort, diminishing ecGFR showed a monotonic relationship to mortality across all strata of renal function. Notably, this relationship is evident even in the absence of data on urinary protein excretion.<sup>6</sup>

Astor *et al.*<sup>7</sup> assembled a cohort of patients from the general population using the Third National Health Examination Survey (NHANES) to evaluate the risk of death associated with lower levels of eGFR estimated from cystatin C, from serum creatinine, or from both. Compared with previous published reports, the strengths of this work include median follow-up of 8 yr, inclusion of patients with ethnic diversity, a wide age range, and analytical rigor with particular focus on the implications of cystatin C among patients with eGFR >60 ml/min per 1.73 m<sup>2</sup> as calculated using the MDRD equation. Similar to the study by Shlipak *et al.*,<sup>6</sup> the results by Astor *et al.*<sup>7</sup> show an inverse and generally monotonic relationship between ecGFR and mortality, whereas the estimated filtration rates derived from MDRD and from cystatin and creatinine had more J-shaped relationships to all-cause mortality (Figure 1 in the accompanying article<sup>7</sup>).

Lower ecGFR is also informative in predicting mortality, in particular in the range in which creatinine-based eGFR underestimates renal function and is prone to misclassifying patients as having stage 3 CKD according to the Kidney Disease Outcomes Quality Initiative (KDOQI) staging system.<sup>13</sup> Among patients classified by the MDRD equation as having eGFR 30 to 59 or eGFR 60 to 89 ml/min per 1.73 m<sup>2</sup>, overall mortality is significantly elevated only when cystatin C also categorizes patients as having an ecGFR <60 ml/min per 1.73 m<sup>2</sup>; therefore, cystatin C is able to stratify patients with respect to the risk for death even among those with mild or even no reduction in renal function as assessed by a creatinine-based equation.

Astor *et al.*<sup>7</sup> also address the question of whether serum

cystatin C should be used alone or in combination with creatinine to determine patients who are at risk for adverse health events. Previous studies that examined the performance characteristics of the MDRD and other equations, such as the pooled analysis of patients with CKD by Stevens *et al.*,<sup>11</sup> found the combination of creatinine and cystatin C perform best in estimating measured GFR. In contrast, the receiver operating characteristic curves presented by Astor *et al.*<sup>7</sup> reveal that estimated filtration rates derived from the combination of cystatin C and creatinine levels is actually *worse* at predicting death compared with cystatin C alone. Potential explanations for the divergence of cystatin C from creatinine with respect to predicting death include the possibility that cystatin C is affected by systemic inflammation or other pathophysiologic pathways that amplify the mortality risk independent of GFR.<sup>14,15</sup> A second possibility is that limitations of creatinine as a filtration marker (particularly when GFR is mildly diminished) stand in the way of attempts to elucidate relationships between renal function and mortality that cystatin C–based measures alone can reveal.

Astor *et al.*<sup>7</sup> have strengthened the evidence linking cystatin C to mortality risk, but more work is required before cystatin C can be integrated into clinical practice. First, we need studies showing whether using cystatin C adds value in predicting risk for progression from CKD to ESRD. Given that most patients with CKD die before needing renal replacement therapy,<sup>12</sup> it is possible that risk factors for death differ from risk factors for further loss of renal function. New analyses of cystatin C and renal events should, like this study by Astor *et al.*,<sup>7</sup> also focus on the general population with inclusion of patients with GFR >60 ml/min per 1.73 m<sup>2</sup>. Second, future studies that relate cystatin C to mortality and renal deterioration should ideally be informed by data on proteinuria. Third, to justify bringing cystatin C to routine clinical care, trials showing that the use of cystatin C leads to better selection of patients for treatments that reduce morbidity and mortality in CKD must be performed.

Astor *et al.*<sup>7</sup> confirm that ecGFR is strongly associated with subsequent mortality in a large community-based population. Cystatin C holds promise as a prognostic marker that could eventually be incorporated into existing measures that risk-stratify patients for cardiovascular events or death. Additional careful research, like the study by Astor *et al.*,<sup>7</sup> is needed to clarify whether the association between cystatin C and mortality is mediated by renal function alone. Although further studies are required, Astor *et al.*<sup>7</sup> have moved us closer to sorting out the complex set of relationships among contemporary filtration markers, progressive kidney disease, and mortality risk.

## ACKNOWLEDGMENTS

Dr. Reese is supported by National Institutes of Health grant K23-DK078688-01; Dr. Feldman is supported by National Institutes of Health grant K24-DK002651.

## DISCLOSURES

None.

## REFERENCES

1. Glassock RJ, Winearls C: Screening for CKD with eGFR: Doubts and dangers. *Clin J Am Soc Nephrol* 3: 1563–1568, 2008
2. Baxmann AC, Ahmed MS, Marques NC, Menon VB, Pereira AB, Kirsztajn GM, Heilberg IP: Influence of muscle mass and physical activity on serum and urinary creatinine and serum cystatin C. *Clin J Am Soc Nephrol* 3: 348–354, 2008
3. Rule AD, Gussak HM, Pond GR, Bergstralh EJ, Stegall MD, Cosio FG, Larson TS: Measured and estimated GFR in healthy potential kidney donors. *Am J Kidney Dis* 43: 112–119, 2004
4. Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF 3rd, Feldman HI, Kusek JW, Eggers P, Van Lente F, Greene T, Coresh J: A new equation to estimate glomerular filtration rate. *Ann Intern Med* 150: 604–612, 2009
5. Shlipak MG, Katz R, Sarnak MJ, Fried LF, Newman AB, Stehman-Breen C, Seliger SL, Kestenbaum B, Psaty B, Tracy RP, Siscovick DS: Cystatin C and prognosis for cardiovascular and kidney outcomes in elderly persons without chronic kidney disease. *Ann Intern Med* 145: 237–246, 2006
6. Shlipak MG, Sarnak MJ, Katz R, Fried LF, Seliger SL, Newman AB, Siscovick DS, Stehman-Breen C: Cystatin C and the risk of death and cardiovascular events among elderly persons. *N Engl J Med* 352: 2049–2060, 2005
7. Astor BC, Levey AS, Stevens LA, Van Lente F, Selvin E, Coresh J: Method of glomerular filtration rate estimation affects prediction of mortality risk. *J Am Soc Nephrol* 20: 2214–2222, 2009
8. Ishani A, Grandits GA, Grimm RH, Svendsen KH, Collins AJ, Prineas RJ, Neaton JD: Association of single measurements of dipstick proteinuria, estimated glomerular filtration rate, and hematocrit with 25-year incidence of end-stage renal disease in the multiple risk factor intervention trial. *J Am Soc Nephrol* 17: 1444–1452, 2006
9. Gansevoort RT, de Jong PE: The case for using albuminuria in staging chronic kidney disease. *J Am Soc Nephrol* 20: 465–468, 2009
10. Dhamidharka VR, Kwon C, Stevens G: Serum cystatin C is superior to serum creatinine as a marker of kidney function: A meta-analysis. *Am J Kidney Dis* 40: 221–226, 2002
11. Stevens LA, Coresh J, Schmid CH, Feldman HI, Froissart M, Kusek J, Rossert J, Van Lente F, Bruce RD 3rd, Zhang YL, Greene T, Levey AS: Estimating GFR using serum cystatin C alone and in combination with serum creatinine: A pooled analysis of 3,418 individuals with CKD. *Am J Kidney Dis* 51: 395–406, 2008
12. Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY: Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med* 351: 1296–1305, 2004
13. den Hartog JR, Reese PP, Cizman B, Feldman HI: The costs and benefits of automatic estimated glomerular filtration rate reporting. *Clin J Am Soc Nephrol* 4: 419–427, 2009
14. Fried LF: Creatinine and cystatin C: What are the values? *Kidney Int* 75: 578–580, 2009
15. Knight EL, Verhave JC, Spiegelman D, Hillege HL, de Zeeuw D, Curhan GC, de Jong PE: Factors influencing serum cystatin C levels other than renal function and the impact on renal function measurement. *Kidney Int* 65: 1416–1421, 2004

See related article, “Method of Glomerular Filtration Rate Estimation Affects Prediction of Mortality Risk,” on pages 2214–2222.

## Warfarin and Stroke Outcomes in Hemodialysis Patients with Atrial Fibrillation

Simonetta Genovesi\* and Antonio Santoro<sup>†</sup>

\*Department of Clinical Medicine and Prevention, University of Milano-Bicocca, Milano, Italy, and Nephrology Unit, San Gerardo Hospital, Monza, Italy; and <sup>†</sup>Nephrology, Dialysis and Hypertension Department, University of Bologna and S. Orsola-Malpighi Hospital, Bologna, Italy

*J Am Soc Nephrol* 20: 2090–2092, 2009.  
doi: 10.1681/ASN.2009070754

The prevalence of atrial fibrillation in ESRD is extremely high, reaching 27%,<sup>1</sup> and fibrillation, as in the general population, also is associated with increased mortality in hemodialysis patients.<sup>2</sup> A large number of trials show the usefulness of oral anticoagulation therapy<sup>1</sup> for primary and secondary prevention of stroke in patient populations with atrial fibrillation absent ESRD.<sup>3</sup> Recently, a large study demonstrated the superiority of oral anticoagulation therapy compared with the combination of clopidogrel plus aspirin with regard to stroke prevention, with no added risk of bleeding.<sup>4</sup> Even trials performed in patients with high hemorrhagic risk who took warfarin, particularly the elderly, show that benefits of treatment exceed the risks when the international normalized ratio (INR) is monitored correctly.

The decision to use oral anticoagulation therapy, particularly warfarin, in patients with atrial fibrillation involves weighing the risk of a thromboembolic event without therapy, or with inadequate anticoagulation, against the risk of a hemorrhagic event on therapy, particularly over-anticoagulation. Efficacy and safety of anticoagulation in atrial fibrillation depend on maintaining the INR between 2 and 3,<sup>5</sup> as recommended by most practice guidelines.<sup>6</sup> Recently an INR of 3.0 to 3.4 has been proposed to achieve optimal anticoagulation intensity in patients with atrial fibrillation.<sup>7</sup> However, dialysis populations are different from the general population.

The association between renal dysfunction and bleeding has long been recognized, even as long as 200 yr ago,<sup>8</sup> and morbidity and mortality from bleeding remain a significant clinical problem in ESRD. Impaired platelet function is one of the main determinants of uremic bleeding. This impairment is multifactorial and includes defects that are intrinsic to platelets and abnormal platelet–endothelial interactions. Uremic toxins and anemia also play a role.<sup>9</sup> Moreover, hemodialysis patients, unlike other patient settings, are exposed to continuous

Published online ahead of print. Publication date available at www.jasn.org.

**Correspondence:** Dr. Simonetta Genovesi, Dipartimento di Medicina Clinica e Prevenzione, Via Cadore 48, Monza, Milano 20052, Italy. Phone: +390392332426; Fax: +390392332376; E-mail: simonetta.genovesi@unimib.it

Copyright © 2009 by the American Society of Nephrology