

37. Tradtrantip L, Yangthara B, Padmawar P, Morrison C, Verkman AS: ThiopheneCaSRboxylate suppressor of cyclic nucleotides discovered in a small-molecule screen blocks toxin-induced intestinal fluid secretion. *Mol Pharmacol* 75: 134–142, 2009
38. Brown EM: Clinical lessons from the calcium-sensing receptor. *Nat Clin Pract Endocrinol Metab* 3: 122–133, 2007
39. Brown EM, MacLeod RJ: Extracellular calcium sensing and extracellular calcium signaling. *Physiol Rev* 81: 239–297, 2001
40. Huang C, Miller RT: Regulation of renal ion transport by the calcium-sensing receptor: An update. *Curr Opin Nephrol Hypertens* 16: 437–443, 2007
41. Abdullah HI, Pedraza PL, McGiff JC, Ferreri NR: Calcium-sensing receptor signaling pathways in medullary thick ascending limb cells mediate COX-2-derived PGE2 production: functional significance. *Am J Physiol Renal Physiol* 295: F1082–F1089, 2008
42. Wang W, Lu M, Balazy M, Hebert SC: Phospholipase A2 is involved in mediating the effect of extracellular Ca<sup>2+</sup> on apical K<sup>+</sup> channels in rat TAL. *Am J Physiol* 273: F421–F429, 1997
43. Wang WH, Lu M, Hebert SC: Cytochrome P-450 metabolites mediate extracellular Ca(2+)-induced inhibition of apical K<sup>+</sup> channels in the TAL. *Am J Physiol* 271: C103–C111, 1996
44. Park F, Sweeney WE, Jia G, Roman RJ, Avner ED: 20-HETE mediates proliferation of renal epithelial cells in polycystic kidney disease. *J Am Soc Nephrol* 19: 1929–1939, 2008
45. Ortiz-Capisano MC, Ortiz PA, Garvin JL, Harding P, Beierwaltes WH: Expression and function of the calcium-sensing receptor in juxtaglomerular cells. *Hypertension* 50: 737–743, 2007
46. Sands JM, Flores FX, Kato A, Baum MA, Brown EM, Ward DT, Hebert SC, Harris HW: Vasopressin-elicited water and urea permeabilities are altered in IMCD in hypercalcemic rats. *Am J Physiol* 274: F978–F985, 1998
47. Lee M, Partridge NC: Parathyroid hormone signaling in bone and kidney. *Curr Opin Nephrol Hypertens* April 22, 2009 [epub ahead of print]
48. Bajwa A, Forster MN, Maiti A, Woolbright BL, Beckman MJ: Specific regulation of CYP27B1 and VDR in proximal versus distal renal cells. *Arch Biochem Biophys* 477: 33–42, 2008
49. Gattone VH II, Chen NX, Sinderson RM, Seifert MF, Duan D, Martin D, Henley C, Moe SM: Calcimimetic inhibits late-stage cyst growth in ADPKD. *J Am Soc Nephrol* 20: 1527–1532, 2009
50. Wang X, Harris PC, Somlo S, Batlle D, Torres VE: Effect of calcium-sensing receptor activation in models of autosomal recessive or dominant polycystic kidney disease. *Nephrol Dial Transplant* 24: 526–534, 2009
51. Cowley B Jr, Gudapaty S, Kraybill A, Barash B, Harding M, Calvet J, Gattone VD: Autosomal dominant polycystic kidney disease in the rat. *Kidney Int* 43: 522–534, 1993
52. Schäfer K, Gretz N, Bader M, Oberbäumer I, Eckardt K-U, Kriz W, Bachmann S: Characterisation of the Han-SPRD rat model for hereditary polycystic kidney disease. *Kidney Int* 46: 134–152, 1994
53. Grantham JJ, Torres VE, Chapman AB, Guay-Woodford LM, Bae KT, King BF Jr, Wetzell LH, Baumgarten DA, Kenney PJ, Harris PC, Klahr S, Bennett WM, Hirschman GN, Meyers CM, Zhang X, Zhu F, Miller JP: Volume progression in polycystic kidney disease. *N Engl J Med* 354: 2122–2130, 2006
54. Lager DJ, Qian Q, Bengal RJ, Ishibashi M, Torres VE: The pck rat: A new model that resembles human autosomal dominant polycystic kidney and liver disease. *Kidney Int* 59: 126–136, 2001
55. Torres VE: Vasopressin antagonists in polycystic kidney disease. *Kidney Int* 68: 2405–2418, 2005
56. Wu G, D'Agati V, Cai Y, Markowitz G, Park J, Reynolds D, Maeda Y, Le T, Hou H Jr, Kucherlapati R, Edelmann W, Somlo S: Somatic inactivation of PKD2 results in polycystic kidney disease. *Cell* 93: 177–188, 1998
57. Costa MZ, Bacchi CE, Franco M: Histogenesis of the acquired cystic kidney disease: An immunohistochemical study. *Appl Immunohistochem Mol Morphol* 14: 348–352, 2006
58. Ishikawa I, Horiguchi T, Shikura N: Lectin peroxidase conjugate reactivity in acquired cystic disease of the kidney. *Nephron* 51: 211–214, 1989
59. Ogata H, Ritz E, Odoni G, Amann K, Orth SR: Beneficial effects of calcimimetics on progression of renal failure and CaSRdiovacular risk factors. *J Am Soc Nephrol* 14: 959–967, 2003
60. Piecha G, Kokeny G, Nakagawa K, Koleganova N, Geldyev A, Berger I, Ritz E, Schmitt CP, Gross ML: Calcimimetic R-568 or calcitriol: Equally beneficial on progression of renal damage in subtotal nephrectomized rats. *Am J Physiol Renal Physiol* 294: F748–F757, 2008
61. Koleganova N, Piecha G, Ritz E, Bekeredjian R, Schirmacher P, Schmitt CP, Gross ML: Interstitial fibrosis and microvascular disease of the heart in uremia: Amelioration by a calcimimetic. *Lab Invest* 89: 520–530, 2009
62. Charytan C, Coburn JW, Chonchol M, Herman J, Lien YH, Liu W, Klassen PS, McCaSRy LC, Pichette V: Cinacalcet hydrochloride is an effective treatment for secondary hyperparathyroidism in patients with CKD not receiving dialysis. *Am J Kidney Dis* 46: 58–67, 2005
63. Chonchol M, Locatelli F, Abboud HE, Charytan C, de Francisco AL, Jolly S, Kaplan M, Roger SD, Sarkar S, Albizem MB, Mix TC, Kubo Y, Block GA: A randomized, double-blind, placebo-controlled study to assess the efficacy and safety of cinacalcet HCl in participants with CKD not receiving dialysis. *Am J Kidney Dis* 53: 197–207, 2009
64. Kalantar-Zadeh K, Kovesdy CP: Is it worth correcting hyperparathyroidism if hyperphosphatemia and hypocalcemia worsen? A cinacalcet story. *Am J Kidney Dis* 53: 183–188, 2009

See related article, "Calcimimetic Inhibits Late-Stage Cyst Growth in ADPKD," on pages 1527–1532.

## The Promise of Well-Being: Stay in Shape with N(i)ck

Thomas Benzing

Department of Medicine, Centre for Molecular Medicine, and Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases, University of Cologne, Cologne, Germany

*J Am Soc Nephrol* 20: 1425–1427, 2009.  
doi: 10.1681/ASN.2009040453

Just more than a decade ago, a report of a rare human genetic disorder initiated a new spurt of research that profoundly changed our understanding of the glomerular filtration barrier. In that landmark article,<sup>1</sup> mutations in the gene encoding the adhesion protein nephrin identified the cause of congenital nephrotic syndrome of the Finnish type. Subsequent studies localized nephrin to the podocyte slit diaphragm, a specialized

Published online ahead of print. Publication date available at www.jasn.org.

**Correspondence:** Dr. Thomas Benzing, Renal Division, Department of Medicine, University of Cologne, Kerpener Strasse 62, 50937 Cologne, Germany. Phone: +49-221-478-4480; Fax: +49-221-478-5959; E-mail: thomas.benzing@uk-koeln.de

Copyright © 2009 by the American Society of Nephrology

cell junction that connects neighboring podocytes, the visceral epithelial cells of the glomerulus. These neighboring podocytes elaborate long, regularly spaced, interdigitated foot processes that enwrap the glomerular capillaries and form a 40-nm-wide filtration slit bridged by the slit diaphragm.

Further studies suggested that nephrin and associated proteins, in addition to their role as structural components of the slit diaphragm, act as signaling molecules to control podocyte development, cytoskeletal architecture, cell shape and viability, and the function of the glomerular filtration barrier.<sup>2–4</sup> Much insight into the signaling function of the slit diaphragm protein complex resulted from the identification of nephrin-interacting proteins,<sup>5–9</sup> yet the mechanisms that control ultrastructure and shape of the highly organized podocyte foot process remain poorly understood. This changed with the finding that nephrin binds Src homology 2 (SH2)/SH3 domain-containing Nck adaptor proteins, which in turn seems to control the podocyte cytoskeleton *in vivo*.<sup>10,11</sup>

The Nck family of adaptors (mammals carry two related *Nck* genes, *Nck1* and *Nck2*) functions in coupling phosphotyrosine signals to actin cytoskeletal reorganization. Each of these proteins contain multiple protein interaction domains: three SH3 domains that are known to interact with proline-rich binding partners such as the actin organizers WASP, SPIN90, and dynamin and an SH2 domain that mediates binding of transmembrane receptors and adhesion proteins through phosphotyrosine-dependent protein interactions.<sup>12</sup> Nephrin ectodomains recruit the Src family tyrosine kinase Fyn to phosphorylate the nephrin cytoplasmic tail, allowing recruitment of Nck and inducing localized Nck-driven actin polymerization.<sup>13</sup> Notably, selective gene deletion of *Nck1* and *Nck2* from podocytes in transgenic mice results in severe defects in the formation of podocyte processes and in congenital nephrotic syndrome.<sup>10</sup> Although the profound changes in cell shape and podocyte morphology in this model mirror the stereotypical reaction of podocytes to injury or damage,<sup>14,15</sup> the phenotype is clearly a developmental defect—mice are born without Nck1 or Nck2 in podocytes—and does not clarify the important question of whether slit diaphragm signaling and nephrin-mediated recruitment of Nck are involved in actively regulating podocyte cell shape and maintaining the ultrastructural and functional integrity of the glomerular filtration barrier *in vivo*.

In a remarkable article in this issue of *JASN*, Jones *et al.*<sup>16</sup> now show that inducible gene deletion of *Nck* in adult transgenic mice results in podocyte cell shape changes, massive proteinuria, and the development of glomerular scarring. Jones *et al.* use an inducible transgenic strategy that allows selective deletion of Nck expression in podocytes of adult mice. The authors carefully analyze the ultrastructural changes of podocytes in adult mice lacking Nck expression and definitively demonstrate that Nck1 and Nck2 maintain the shape and integrity of foot processes.

This is interesting for a number of reasons. In response to injury, podocytes undergo a dramatic change in morphol-

ogy—termed foot process effacement—resulting from alterations in cytoskeletal and intercellular junctional architecture. Effacement results from simplification of foot processes, namely loss of foot process shape and retraction, widening, and shortening of the processes of each podocyte. The frequency of filtration slits is also reduced, giving the appearance of a continuous cytoplasmic sheet covering the glomerular basement membrane.<sup>15</sup>

Effacement is not specific to one disease but rather is synonymous with podocyte injury of many forms, providing a fluid and reversible process that directly correlates with the development of proteinuria both in human disease and in experimental models. Thus, it is interesting that *Nck* gene deletion and, accordingly, loss of nephrin-dependent Nck recruitment exactly mirror this phenotype. The glomerular morphology of gene-deleted mice undergo the widely known continuum from mild foot process widening to focal areas of fusion and the development of extensive effacement that mirrors minimal-change disease in humans; however, these changes obviously precede the loss of podocytes and the development of focal areas of glomerulosclerosis,<sup>16</sup> which is consistent with FSGS in humans, raising the hypothesis that minimal-change disease and FSGS in many cases are two extremes of an ellipse.

Nephrin and associated proteins are conserved through evolution and play important roles in patterning of the eye in *Drosophila melanogaster* or synapse targeting in *Caenorhabditis elegans*. All of these changes require defined cytoskeletal rearrangements; therefore, it will be important to determine whether nephrin-induced Nck activity is also required in these events. As Jones *et al.*<sup>16</sup> clearly demonstrate, nephrin phosphorylation is maintained in the adult glomerulus supporting the importance of signaling at the slit diaphragm for regulation of podocyte cell shape, cytoskeletal architecture, and cell viability.

It is worth noting that these new findings have the same limitations as the original report concerning a developmental role of nephrin-Nck signaling. Although Jones *et al.* clearly demonstrate that *Nck1* and *Nck2* are essential for developing and maintaining the complex ultrastructure of podocyte secondary processes, both articles lack direct proof that Nck proteins exclusively act through the nephrin/slit diaphragm axis *in vivo*. In fact, this has to be addressed in future studies, for instance using knock-in mouse models of nephrin with mutated tyrosine residues. Despite this limitation, Jones *et al.*<sup>16</sup> undoubtedly demonstrate the importance of Nck in controlling the ultrastructure of podocyte foot processes in maintaining the integrity of the glomerular filtration barrier.

## ACKNOWLEDGMENTS

I thank members of the Benzing laboratory for helpful discussions and critically reading the manuscript.

## DISCLOSURES

None.

## REFERENCES

1. Kestila M, Lenkkeri U, Mannikko M, Lamerdin J, McCready P, Putaala H, Ruotsalainen V, Morita T, Nissinen M, Herva R, Kashtan CE, Peltonen L, Holmberg C, Olsen A, Tryggvason K: Positionally cloned gene for a novel glomerular protein—nephrin—is mutated in congenital nephrotic syndrome. *Mol Cell* 1: 575–582, 1998
2. Schermer B, Benzing T: Lipid-protein interactions along the slit-diaphragm of podocytes. *J Am Soc Nephrol* 20: 473–478, 2009
3. Benzing T: Signaling at the slit-diaphragm. *J Am Soc Nephrol* 15: 1382–1391, 2004
4. Johnstone DB, Holzman LB: Clinical impact of research on the podocyte slit-diaphragm. *Nat Clin Pract Nephrol* 2: 271–282, 2006
5. Huber TB, Hartleben B, Kim J, Schmidts M, Schermer B, Keil A, Egger L, Lecha RL, Borner C, Pavenstadt H, Shaw AS, Walz G, Benzing T: Nephrin and CD2AP associate with phosphoinositide 3-OH kinase and stimulate AKT-dependent signaling. *Mol Cell Biol* 23: 4917–4928, 2003
6. Huber TB, Kottgen M, Schilling B, Walz G, Benzing T: Interaction with podocin facilitates nephrin signaling. *J Biol Chem* 276: 41543–41546, 2001
7. Schwarz K, Simons M, Reiser J, Saleem MA, Faul C, Kriz W, Shaw AS, Holzman LB, Mundel P: Podocin, a raft-associated component of the glomerular slit-diaphragm, interacts with CD2AP and nephrin. *J Clin Invest* 108: 1621–1629, 2001
8. Verma R, Wharram B, Kovari I, Kunkel R, Nihalani D, Wary KK, Wiggins RC, Killen P, Holzman LB: Fyn binds to and phosphorylates the kidney slit-diaphragm component Nephrin. *J Biol Chem* 278: 20716–20723, 2003
9. Harita Y, Kurihara H, Kosako H, Tezuka T, Sekine T, Igarashi T, Ohsawa I, Ohta S, Hattori S: Phosphorylation of nephrin triggers Ca<sup>2+</sup> signaling by recruitment and activation of phospholipase C-gamma1. *J Biol Chem* 284: 8951–8962, 2009
10. Jones N, Blasutig IM, Eremina V, Ruston JM, Bladt F, Li H, Huang H, Larose L, Li SS, Takano T, Quaggin SE, Pawson T: Nck adaptor proteins link nephrin to the actin cytoskeleton of kidney podocytes. *Nature* 440: 818–823, 2006
11. Verma R, Kovari I, Soofi A, Nihalani D, Patrie K, Holzman LB: Nephrin ectodomain engagement results in Src kinase activation, nephrin phosphorylation, Nck recruitment, and actin polymerization. *J Clin Invest* 116: 1346–1359, 2006
12. Songyang Z, Shoelson SE, Chaudhuri M, Gish G, Pawson T, Haser WG, King F, Roberts T, Ratnofsky S, Lechleider RJ, et al.: SH2 domains recognize specific phosphopeptide sequences. *Cell* 72: 767–778, 1993
13. Blasutig IM, New LA, Thanabalasuriar A, Dayarathna TK, Goudreault M, Quaggin SE, Li SS, Gruenheid S, Jones N, Pawson T: Phosphorylated YDXV motifs and Nck SH2/SH3 adaptors act cooperatively to induce actin reorganization. *Mol Cell Biol* 28: 2035–2046, 2008
14. Faul C, Asanuma K, Yanagida-Asanuma E, Kim K, Mundel P: Actin up: Regulation of podocyte structure and function by components of the actin cytoskeleton. *Trends Cell Biol* 17: 428–437, 2007
15. Shankland SJ: The podocyte's response to injury: Role in proteinuria and glomerulosclerosis. *Kidney Int* 69: 2131–2147, 2006
16. Jones N, New LA, Fortino MA, Eremina V, Ruston J, Blasutig IM, Aoudjit L, Zou Y, Liu X, Yu G-L, Takano T, Quaggin SE, Pawson T: Nck proteins maintain the adult glomerular filtration barrier. *J Am Soc Nephrol* 20: 1533–1543, 2009

See related article, "Nck Proteins Maintain the Adult Glomerular Filtration Barrier," on pages 1533–1543.

## Glomerular Filtration: Still Sympathetic to Endothelin's Influence?

Tracy E. Hunley and Valentina Kon

Division of Nephrology, Department of Pediatrics, Vanderbilt University School of Medicine, Nashville, Tennessee

*J Am Soc Nephrol* 20: 1427–1429, 2009.

doi: 10.1681/ASN.2009050503

Methodologic and analytic advances in molecular genetics identify many disease-predisposing variants in the human genome. One of the challenges presented by these studies is to define the mechanisms by which a given genetic variant affects a pathophysiologic disorder. This inevitable quandary confronts the study of Chen *et al.*<sup>1</sup> in this issue of *JASN*. The authors previously reported that a genetic polymorphism of chromatogranin A (CHGA), a soluble protein released from secretory granules of chromaffin cells and sympathetic nerves, predicts hypertensive ESRD in black patients. Using population genetic techniques, they further showed that the *CHGA* locus is a *trans*-QTL for endothelin-1 secretion and that CHGA stimulates endothelin release from endothelial cells. In view of the major role for endothelin postulated in the pathogenesis of a variety of cardiovascular and renal diseases, the authors questioned whether endothelin secretion is the mechanistic link between CHGA and renal dysfunction.

Using cultured human umbilical vein endothelial cells, Chen *et al.*<sup>1</sup> now show that CHGA increases release of endothelin along with other peptides typically stored in Weibel-Palade bodies, including von Willebrand factor and angiotensin 2. CHGA also increases endothelin secretion from mouse glomerular endothelial cells, and co-culture of CHGA-exposed glomerular endothelial cells with mesangial cells potentiates secretion of TGF- $\beta$ 1. In humans, plasma CHGA correlates positively with plasma endothelin and negatively with GFR. A promoter haplotype for *CHGA* predicts the level of GFR in individuals with GFRs in the normal range, whereas *CHGA* haplotypes in the 3' untranslated region associate with GFR in individuals with progressive renal disease. The conclusion that CHGA triggers exocytotic release of endothelin from the glomerular endothelium and that this, in turn, regulates the level of GFR in normal and declining renal function offers some novel implications. These include genetic modulation of GFR in normal and diseased kidneys under the potential influence of the sympathochromaffin protein, the utility of circulating

Published online ahead of print. Publication date available at [www.jasn.org](http://www.jasn.org).

**Correspondence:** Dr. Valentina Kon, Pediatric Nephrology, Vanderbilt University Medical Center, C-4204 Medical Center North, Nashville, TN 37232-2584. Phone: 615-322-7416; Fax: 615-322-7929; E-mail: [valentina.kon@vanderbilt.edu](mailto:valentina.kon@vanderbilt.edu)

Copyright © 2009 by the American Society of Nephrology