

Sodium and Potassium Balance Depends on α ENaC Expression in Connecting Tubule

Birgitte Mønster Christensen,* Romain Perrier,* Qing Wang,[†] Annie Mercier Zuber,* Marc Maillard,[†] David Mordasini,[†] Sumedha Malsure,* Caroline Ronzaud,[‡] Jean-Christophe Stehle,[§] Bernard C. Rossier,* and Edith Hummler*

Departments of *Pharmacology and Toxicology and [§]Pathology, University of Lausanne, Lausanne, Switzerland; [†]Service of Nephrology, Lausanne University Hospital, Lausanne, Switzerland; and [‡]German Cancer Research Center, Heidelberg, Germany

ABSTRACT

Mutations in α , β , or γ subunits of the epithelial sodium channel (ENaC) can downregulate ENaC activity and cause a severe salt-losing syndrome with hyperkalemia and metabolic acidosis, designated pseudohypoaldosteronism type 1 in humans. In contrast, mice with selective inactivation of α ENaC in the collecting duct (CD) maintain sodium and potassium balance, suggesting that the late distal convoluted tubule (DCT2) and/or the connecting tubule (CNT) participates in sodium homeostasis. To investigate the relative importance of ENaC-mediated sodium absorption in the CNT, we used Cre-lox technology to generate mice lacking α ENaC in the aquaporin 2-expressing CNT and CD. Western blot analysis of microdissected cortical CD (CCD) and CNT revealed absence of α ENaC in the CCD and weak α ENaC expression in the CNT. These mice exhibited a significantly higher urinary sodium excretion, a lower urine osmolality, and an increased urine volume compared with control mice. Furthermore, serum sodium was lower and potassium levels were higher in the genetically modified mice. With dietary sodium restriction, these mice experienced significant weight loss, increased urinary sodium excretion, and hyperkalemia. Plasma aldosterone levels were significantly elevated under both standard and sodium-restricted diets. In summary, α ENaC expression within the CNT/CD is crucial for sodium and potassium homeostasis and causes signs and symptoms of pseudohypoaldosteronism type 1 if missing.

J Am Soc Nephrol 21: 1942–1951, 2010. doi: 10.1681/ASN.2009101077

Sodium reabsorption in the kidney is essential for maintaining fluid and electrolyte homeostasis as well as regulation of blood pressure (BP). Renal sodium reabsorption is under tight control of aldosterone in the late distal convoluted tubule (DCT2), the connecting tubule (CNT), and the collecting duct (CD).¹ Sodium enters the aldosterone-sensitive epithelial cell through the epithelial sodium channel (ENaC) at the apical plasma membrane, and sodium is extruded to the interstitial fluid *via* the basolateral Na⁺-K⁺-ATPase in exchange for potassium. In the DCT2, sodium is also absorbed through the thiazide-sensitive NaCl co-transporter (TSC).² The critical role of ENaC in sodium homeostasis has been emphasized by identification of gain-of-function mutations in the C-terminus of the β or the γ subunit in patients with Liddle syndrome, a severe form of hypertension caused by sodium retention.^{3,4}

Pseudohypoaldosteronism type 1 (PHA-1), conversely, is a severe salt-wasting syndrome

Received October 21, 2009. Accepted August 1, 2010.

Published online ahead of print. Publication date available at www.jasn.org.

R.P. and Q.W. contributed equally to this work.

B.M.C.'s current affiliation is Water and Salt Research Center, Department of Anatomy, Aarhus University, Aarhus, Denmark; A.M.Z.'s current affiliation is Department of Medicine, University of Cambridge, Cambridge, United Kingdom; C.R.'s current affiliation is Department of Pharmacology and Toxicology, University of Lausanne, Lausanne, Switzerland.

Correspondence: Dr. Birgitte Mønster Christensen, Water and Salt Research Center, Department of Anatomy, Aarhus University, Wilhelm Meyers Allé 3, 8000 Aarhus C, Denmark. Phone: +45-89423007; Fax: +45-86198664; E-mail: bmc@ana.au.dk; or Dr. Edith Hummler, Department of Pharmacology and Toxicology, University of Lausanne, Rue du Bugnon 27, CH-1005 Lausanne, Switzerland. Phone: +41216925357; Fax: +41216925355; E-mail: Edith.Hummler@unil.ch

Copyright © 2010 by the American Society of Nephrology

characterized by urinary loss of sodium and reduced potassium excretion despite elevated levels of aldosterone. In humans, a life-threatening form of the disease is inherited as an autosomal recessive trait and is caused by loss-of-function mutations in any of the three ENaC subunits.⁵ Clinical symptoms of the disease are weight loss and dehydration, hypovolemia and hypotension, hyponatremia, hyperkalemia, and metabolic acidosis accompanied by elevated plasma aldosterone levels.⁶ Complete knockout (KO) of each of the ENaC subunits resulted in an early and lethal PHA-1 phenotype.^{7–9} Previously, a CD-specific conditional KO for α ENaC was generated, and, surprising, these mice were able to maintain water, sodium, and potassium balance even after 1 week of salt restriction, 23 hours of water deprivation, or 4 days of potassium loading.¹⁰ In this study, we investigated the implication of the CNT and CD for the ENaC-mediated sodium reabsorption using mice that express the Cre recombinase from the *Aqp2* promoter (*Aqp2::iCre*) and conditional alleles of α ENaC (*Scnn1a^{lox/lox}*).^{11,12} Aquaporin 2 (AQP2) is a water channel expressed along the CNT and CD.^{13,14} Our data indicate that α ENaC expression within the CNT is important for sodium and potassium balance.

RESULTS

Inactivation of α ENaC in the CD and the CNT

KO and control mice were born consistent with Mendelian inheritance (*Scnn1a^{lox/-}/Aqp2::iCre* 25.9%, *Scnn1a^{lox/+}/Aqp2::iCre* 29.4%, *Scnn1a^{lox/+}* 21.8%, and *Scnn1a^{lox/-}* 22.8%; $n = 197$). To verify the deletion of α ENaC expression in these CNT/CD-specific KO mice (*Scnn1a^{lox/-}/Aqp2::iCre*), we dissected the CNT and the CCD and analyzed these by Western blot (Figure 1). The previously described CD-specific KO mice (*Scnn1a^{lox/lox}/HoxB7::Cre*) were used as positive (and negative) control.¹⁰ In the CNT and the CCD of all control groups, a band at approximately 95 kD corresponding to full-length α ENaC protein was observed (Figure 1A, lanes 1 and 2, and B, lanes 1 and 2 and lanes 5 and 6). This band was absent in the CCD of the CD-specific KO mice (Figure 1A, lane 4)¹⁰ and in the CCD of the CNT/CD-specific KO mice (Figure 1B, lanes 4 and 8). We observed a faint band corresponding to full-length α ENaC protein in the CNT of the CNT/CD-specific KO mice (Figure 1B, lanes 3 and 7), suggesting that the inactivation of α ENaC in the CNT was not complete. We further observed a band just above 26 kD in the CNT of the CD-specific KO mice (Figure 1A, lane 3) and in the CNT of the control littermates




Figure 1. Loss of α ENaC expression in CD and partial loss in CNT of CNT/CD-specific KO mice. (A and B) Western blot of microdissected CNT and CCD from CD-specific KO mice and their corresponding controls (A) and from the CNT/CD-specific KO mice and littermate controls (B). Anti- α ENaC antibody recognizes a band at approximately 95 kD corresponding to full-length α ENaC and a band just above 26 kD that may correspond to a cleaved fragment of α ENaC (arrowheads*). Some unspecific bands are also observed (arrowheads). The blots are reprobed with anti-actin antibody to examine the amount of protein loaded on the gels.

of the CNT/CD-specific KO mice (Figure 1B, lanes 1 and 5) that most likely corresponds to a cleavage product of the α subunit.^{15,16} It was not observed in the CNT of the CNT/CD-specific KO mice (Figure 1B, lanes 3 and 7).

Western blots of microdissected tubules from sodium-restricted CNT/CD-specific KO mice and littermate control mice showed that the approximately 95 kD band corresponding to full-length α ENaC protein was absent in the CCD of the CNT/CD-specific KO mice (Figure 2). In the CNT of the CNT/CD-specific KO mice, this band was weakly expressed or absent (Figure 2). Thus, on a standard or salt-restricted diet (Figures 1 and 2), α ENaC protein is barely detectable in the CNT and not detected in the CCD of KO mice.

To investigate further the recombination efficiency, we performed immunohistochemistry on kidneys from sodium-deprived CNT/CD-specific KO mice and control mice. Double labeling showed almost complete co-localization of AQP2 and Cre recombinase in the principal cells along the CD (Figure 3A). Cellular counting in the CCD revealed that 96% of the AQP2-positive cells showed co-staining with the Cre recombinase. In the CNT, cellular counting showed that approximately 70% of AQP2-positive cells were co-labeled with Cre recombinase (Figure 3B). Double labeling of Cre recombinase and H^+ -ATPase (marker for intercalated cells) revealed no Cre expression in the intercalated cells (Figure 3C).

Within the CNT, immunolabeling was performed with antibodies recognizing α ENaC, AQP2, and TSC. The anti-TSC antibody was used as a marker of the DCT.² In control mice, α ENaC was expressed along the early and late CNT (Figure 3, D through F). The early CNT, which was identified at the transition from the TSC-positive DCT to the TSC-negative CNT, contained none or only a few AQP2-positive cells in both con-




Figure 2. Sodium-restricted CNT/CD-specific KO mice show near-loss of α ENaC protein expression in CNT. (A through D) Western blots of microdissected proximal tubules (Prox), CNT, and CCD from four different KO and control mice that are subjected to sodium-deficient diet. Anti- α ENaC antibody recognizes a band at approximately 95 kDa corresponding to full-length α ENaC (top blots). The blots are reprobbed with anti-actin antibody (bottom blots).

control and KO mice (Figure 3, D through I). Consistent with this, α ENaC was also detected in such tubules, which were both AQP2 and TSC negative and likely represent early parts of the CNT with undetectable AQP2 expression. In KO mice, α ENaC was partly absent in the CNT (Figure 3, G through I) with some expression remaining in both late and early CNT. Approximately 70% of all early CNT cells (both CNT and intercalated cells) in the control were α ENaC positive, in comparison with only 50% in the KO situation. In the late CNT, the percentage of α ENaC-positive CNT cells (control 63%, KO 18%; identified as TSC negative and AQP2 positive) was reduced by approximately 70% in the KO. In summary, the CNT/CD-specific α ENaC KO mice have reduced numbers of α ENaC-expressing cells in both the late and early CNT, with a more severe reduction (up to 70%) in the late CNT, and corresponding to an increase in AQP2-expressing cells in the late CNT.

CNT/CD-Specific α ENaC KO Mice Exhibit a PHA-1 Phenotype under Standard Salt Diet

Following a standard salt diet, mice did not exhibit reduced body weight. In contrary, the urinary sodium excretion was

significantly higher in the KO mice ($P < 0.05$; Table 1), whereas the urinary potassium excretion was unchanged. Food and, thus, sodium intake were not altered (Table 1). The increased urine excretion ($P < 0.01$; Table 1) was accompanied by lower urinary osmolality ($P < 0.01$; Table 1) and higher water intake ($P < 0.05$; Table 1). The CNT/CD-specific α ENaC KO mice presented with significantly lower serum sodium concentrations ($P < 0.05$; Table 1) and hyperkalemia revealed by significantly higher blood potassium concentrations ($P < 0.05$; Table 1). Plasma aldosterone was measured in CNT/CD-specific α ENaC KO and control mice that were homozygous for a specific renin allele (*Ren-2^{-/-}*),¹⁷ and the CNT/CD-specific α ENaC KO mice presented with significantly higher plasma aldosterone levels ($P < 0.05$; Table 2). BP was slightly reduced in the KO mice, without any significant difference (Table 1). No significant changes were observed in the heart rate (HR; Table 1).

Sodium-Deficient Diet Induces Severe Renal Sodium Loss

The CNT/CD-specific α ENaC KO and control mice were challenged with a sodium-deficient diet for 4 consecutive days. Already after 1 day, the KO mice lost significant body weight, whereas the control mice kept or even gained body weight ($P < 0.001$; Figure 4A). No difference was observed in food intake (KO [$n = 8$] 0.15 ± 0.01 versus control [$n = 10$] 0.14 ± 0.01 g/g body wt), and loss of body weight was paralleled by a severe urinary sodium loss (Figure 4B). The cumulative sodium balance showed that the control mice were able to retain their sodium, whereas the KO mice continued to excrete sodium (KO 0.25 ± 0.014 mmol; control 0.07 ± 0.005 mmol; $P < 0.001$, day 4; Figure 4B). After 1 day of sodium-deficient diet, the urinary potassium excretion was significantly increased in the KO mice ($P < 0.05$), a difference that vanished during the subsequent days (Figure 4C). The KO mice continued to have a significant higher urinary excretion during the first 3 days on the sodium-deficient diet (Figure 5A). The urine osmolality was significantly lower in the KO mice on the sodium-deficient diet ($P < 0.001$, day 4; Figure 5B). Water intake was not different (Figure 5C). After 4 days, a significant increase was found in blood potassium levels (KO [$n = 8$] 6.4 ± 0.2 versus control [$n = 10$] 5.8 ± 0.2 mM; $P < 0.05$, day 4) and plasma aldosterone levels ($P < 0.01$; Table 2).

When the mice were followed for a period of 15 days upon salt-deprivation in standard cages, the KO mice lost weight continuously. The weight loss reached more than 10% of their initial weight ($P < 0.001$; Figure 6). Moreover, plasma aldosterone levels were significantly elevated ($P < 0.001$; Table 2).

CNT/CD-Specific α ENaC KO Mice Are not Able to Eliminate a Potassium Load

The ability of the KO mice to eliminate a potassium load was tested by challenging the animals with a diet containing 5% potassium for 2 consecutive days. The KO mice had a significant lower urinary potassium excretion ($P < 0.01$, day 2; Figure 7A) and significantly higher levels of potassium in the




Figure 3. Residual expression of α ENaC in CNT of CNT/CD-specific KO mice. (A through I) Immunohistochemistry using whole kidney sections from sodium-restricted control (D through F) and KO mice (A through C and G through I). (A and B) Double labeling of whole kidney sections from a KO mouse is performed with polyclonal primary antibodies that recognize Cre recombinase (brown) and AQP2 (gray-blue). Along the CD, nuclear Cre recombinase staining is observed in AQP2-expressing cells (A, arrows). Within the CNT, some AQP2-expressing cells are negative for Cre recombinase (arrowhead, B). (C) Double labeling with primary antibodies recognizing Cre recombinase (brown) and H^+ -ATPase (gray-blue) shows no co-localization of the two proteins (arrows). (D through I) Serial kidney sections from a control mouse (D through F) and KO mouse (G through I) incubated with anti-TSC antibody (D and G), anti- α ENaC antibody (E and H), and anti-AQP2 antibody (F and I). In control mice, α ENaC labeling is observed in the CNT (AQP2 positive and TSC negative, D through F, *) including the early CNT, which is identified adjacent to the transition from the TSC-positive DCT to the TSC-negative CNT (D through F, ■). In the KO mice, α ENaC is absent from part of the CNT (G through I, Δ) but present in other parts (G through I, *), including the early CNT (G through I, inset, ■).

blood ($P < 0.05$; Table 3), whereas the serum sodium concentration was significantly reduced ($P < 0.05$; Table 3). After 1 day of a high-potassium diet, urinary sodium excretion was not affected but was significantly reduced after day 2 ($P < 0.05$, day 2; Figure 7B). Fractional excretion of sodium ($FE(Na^+)$) was not different (Table 3), but KO mice showed significantly lower urinary chloride excretion (KO [$n = 11$] 1.1 ± 0.2 versus control [$n = 11$] 2.5 ± 0.4 mmol; $P < 0.05$, day 2), whereas no difference in serum chloride concentration was observed (Table 3). The concentration of bicarbonate was significantly lower in the KO mice (Table 3), indicating metabolic acidosis. Serum osmolality (Table 3), urine osmolality (KO [$n = 11$] 1198 ± 54 mmol/kg H_2O ; control [$n = 11$] 1306 ± 62 mmol/kg H_2O), urine output (KO [$n = 11$] 0.14 ± 0.03 ml/g

body wt per 24 hours; control [$n = 11$] 0.21 ± 0.04 ml/g body wt per 24 hours), and water intake (KO [$n = 11$] 0.28 ± 0.05 ml/g body wt per 24 hours; control [$n = 11$] 0.37 ± 0.04 ml/g body wt per 24 hours) did not reveal significant differences after 2 days of a high-potassium diet.

DISCUSSION

Constitutive KO of either β or γ ENaC causes an early lethal phenotype as a result of disturbances in the electrolyte balance.^{7,9} α ENaC deficiency also induces death shortly after birth, with a lung, skin, and kidney phenotype.^{8,18} Mice in which α ENaC is deleted specifically in the CD are viable and do not show disturbances in sodium and potassium balance even when subjected to challenging diets.¹⁰ This suggested that the CNT is important in controlling ENaC-mediated sodium reabsorption in the kidney. To investigate this further, we generated mice in which the α ENaC gene was deleted in the CD and partly in the CNT. These mice are viable until adulthood and exhibit normal BP, although they show increased urinary sodium excretion, urine output, and plasma aldosterone, leading to hyponatremia and hyperkalemia under standard diet. After sodium restriction, the mice develop a severe salt-losing phenotype and show a continuous life-threatening reduction of body weight.

Partial Inactivation of α ENaC in the CNT Is Sufficient to Induce a Severe Salt-Losing Syndrome

Whereas α ENaC was deleted efficiently in the CCD principal cells, approximately 30% of the late CNT cells are not targeted and still express α ENaC protein. This may explain the remaining α ENaC expression in microdissected CNT of some animals under normal and sodium-deprived diet. Similar to our findings, CNT/CD-specific mineralocorticoid receptor (MR) KO mice (using the same *Aqp2::iCre* transgene) showed complete deletion of MR in the CD, whereas deletion of the MR protein in the early and late CNT was equally partial, thus following the AQP2 expression pattern in these segments.¹¹

The absence of AQP2 expression in the early CNT is therefore consistent with previous observations. In vasopressin-deficient Brattleboro rats, the initial portion of the CNT lacks detectable levels of AQP2, whereas long-term

Table 1. Urinary and blood parameters from mice kept on a standard-salt diet

Parameter	KO	Control
Body weight (g)		
females	21.30 ± 0.66 (n = 13)	22.10 ± 0.65 (n = 24)
males	27.70 ± 0.89 (n = 13)	28.20 ± 1.08 (n = 15)
Food intake (g/g body wt per 24 h)	0.13 ± 0.01 (n = 8)	0.12 ± 0.01 (n = 10)
Na ⁺ intake (mmol/24 h)	0.27 ± 0.03 (n = 8)	0.26 ± 0.03 (n = 10)
Urinary Na ⁺ (mmol/24 h)	0.23 ± 0.01 (n = 31) ^a	0.19 ± 0.01 (n = 46)
Urinary Na ⁺ (mM)	136 ± 11 (n = 31)	155 ± 7 (n = 46)
Urinary K ⁺ (mmol/24 h)	0.33 ± 0.04 (n = 31)	0.28 ± 0.03 (n = 46)
Urinary K ⁺ (mM)	213 ± 38 (n = 31)	242 ± 23 (n = 46)
Urinary osm (mosm/kgH ₂ O)	1809 ± 119 (n = 26) ^b	2442 ± 133 (n = 39)
Urine output (ml/g body wt per 24 h)	0.085 ± 0.008 (n = 31) ^b	0.055 ± 0.004 (n = 46)
Water intake (ml/g body wt per 24 h)	0.25 ± 0.03 (n = 31) ^a	0.16 ± 0.01 (n = 46)
Serum osm (mosm/kgH ₂ O)	309 ± 4 (n = 5)	313 ± 34 (n = 6)
Serum Na ⁺ (mM)	145.8 ± 1.0 (n = 5) ^a	149.8 ± 1.1 (n = 6)
Plasma/serum K ⁺ (mM)	6.3 ± 0.2 (n = 10) ^a	5.6 ± 0.1 (n = 16)
Mean BP (mmHg)	127 ± 2 (n = 13)	135 ± 3 (n = 19)
Mean HR (beats/min)	632 ± 18 (n = 13)	630 ± 13 (n = 19)

Data are means ± SEM.

^aP < 0.05.

^bP < 0.01.

Table 2. Plasma aldosterone (pg/ml) in mice kept on a standard-salt diet or a sodium-deficient diet

Diet	KO	Control
Standard diet (<i>Ren-2</i> ^{-/-} gene mice)	1915 ± 396 (n = 5) ^a	381 ± 55 (n = 8)
Sodium-deficient diet, 4 days (<i>Ren-2</i> ^{-/-} gene mice)	3085 ± 400 (n = 4) ^b	1064 ± 213 (n = 6)
Sodium-deficient diet, 15 days (<i>Ren-2</i> ^{+/-} gene mice)	20,984 ± 2775 (n = 7) ^c	1218 ± 304 (n = 7)

Data are means ± SEM.

^aP < 0.05.

^bP < 0.01.

^cP < 0.001.

vasopressin treatment induced its expression throughout the CNT.¹⁹ Moreover, in *TRPV5::EGFP* transgenic mice, some tubule segments were enhanced green fluorescent protein and calbindin positive but negative for AQP2 and TSC, thus likely representing early CNT.²⁰

We observed that the Cre recombinase protein was expressed along the CD, consistent with CNT/CD-specific MR KO mice.¹¹ Immunohistochemistry also showed that αENaC was not expressed anymore in the early part of the CCD in the CNT/CD-specific αENaC KO mice (Supplemental Figure 1).

The CNT Is Critical for the ENaC-Mediated Sodium Reabsorption

The increased urinary sodium excretion, hyponatremia, and hyperkalemia observed in the KO mice indicate impaired sodium reabsorption in the CNT and are therefore consistent with the absence of ENaC in this segment. Moreover, the KO mice exhibited decreased urine osmolality and increased urine output/water intake. The changes in water balance could be explained by a reduced renal urine-con-

centrating ability, leading to increased urine output and causing increased water intake. The renal effect can be due to impaired ENaC-mediated sodium reabsorption in the CNT and CD, which leads to a reduced driving force for osmotic reabsorption of water in these segments. Apparently, the urine-concentrating ability would be affected only when both the CNT and the CD are targeted because no change in water balance was seen in the CD-specific αENaC KO mice.¹⁰ An impaired urine-concentrating ability was previously described in aldosterone synthase-deficient mice.²¹ The changes in water balance could also be due to the activation of the renin-angiotensin-aldosterone system as shown by elevated plasma aldosterone concentrations, which could lead to a stimulation of thirst and result in increased water intake and subsequently increased urine output.

It was previously shown that CD-specific αENaC KO mice do not exhibit a phenotype even after challenging diets.¹⁰ In contrast, we observed that sodium restriction caused a continuous loss of body weight and sodium in the KO mice. The body weight of the KO mice did not stabilize after 15 days of sodium restriction. A similar weight loss in response to a low-sodium diet was reported in transgenic mice expressing low levels of βENaC.²² Moreover, the severe salt loss in humans

with the recessive form of PHA-1 does not improve with age.⁶ The CNT/CD-specific αENaC KO mice were hyperkalemic in contrast to the CD-specific αENaC KO mice. Both CNT/CD-specific and CD-specific αENaC KO mice decreased their urine osmolality upon sodium restriction, but only the CNT/CD-specific αENaC KO mice exhibited a significant lower urine osmolality compared with their littermate controls. Challenging the CD-specific αENaC KO mice with a potassium-rich diet for 4 days resulted in increased sodium and potassium in both plasma and urine but not different from the control.¹⁰ In contrast, the CNT/CD-specific αENaC KO mice were not able to excrete a potassium load, as shown by significantly higher potassium concentration in blood and decreased urinary potassium excretion. The potassium-loaded CNT/CD-specific αENaC KO mice also showed indications of metabolic acidosis. Thus, in contrast to CD-specific αENaC KO mice, the CNT/CD-specific αENaC KO mice exhibited symptoms of PHA-1, supporting that the CNT is critical for the ENaC-mediated sodium reabsorption. This is consistent with previous suggestions that the late DCT and CNT, rather than the CD, are the main regulators of ENaC-mediated sodium




Figure 4. Loss of body weight and sodium in sodium-deprived CNT/CD-specific KO mice. (A) KO ($n = 18$) and control mice ($n = 23$) are subjected to a sodium-deficient diet for 4 days, and body weight is measured daily. Body weight is presented as percentage of the initial weight. (B and C) Cumulative sodium (B) and potassium (C) balance in KO mice ($n = 18$, ■) and control mice ($n = 23$, □) subjected to a sodium-deficient diet for up to 4 days. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

and potassium homeostasis.^{23,24} The majority (90%) of the sodium delivered to aldosterone-sensitive distal segments is reabsorbed in the CNT and DCT.²⁵ Moreover, the CNT exhibit a higher ENaC activity compared with the CD of aldosterone-infused rats.²⁶ The apical ENaC expression has also been shown to be more pronounced in the CNT compared with the CD in mice on a moderately low-sodium diet (0.05%).²⁷ That a partial gene deletion in the CNT was sufficient to induce a severe salt-losing syndrome also shows that the functional reserve in the CNT is limited because the remaining α ENaC-positive CNT cells were unable to compensate fully for the loss of α ENaC in the other cells.

CNT/CD-Specific Deletion of α ENaC Is More Severe than Inactivation of the MR in the Same Segments

The actions of aldosterone are mediated through the MR. After binding of aldosterone to the MR, the hormone is translocated




Figure 5. Increased urine output and reduced osmolality in sodium-deprived CNT/CD-specific KO mice. (A through C) Urine excretion (A), urine osmolality (B), and water intake (C) are measured daily in KO mice ($n = 18$, ■) and control mice ($n = 23$, □) subjected to a sodium-deficient diet for up to 4 days. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

to the nucleus, where it controls the transcription of ENaC and other genes. In contrast to the CNT/CD-specific α ENaC KO mice, no disturbance in electrolyte and water balance was observed in CNT/CD-specific MR KO mice subjected to a normal-salt diet.¹¹ Thus, partial deletion of α ENaC in the CNT induces a more severe phenotype than inactivation of MR in the same segments. This may not be surprising because the MR protein is an upstream effector on sodium absorption compared with ENaC as an effector. Upon sodium restriction, the CNT/CD-specific MR KO mice also showed loss of body weight, increased sodium and urinary excretion, and significantly higher plasma aldosterone levels,¹¹ similar to CNT/CD-specific α ENaC KO mice.

In summary, gradual gene deletion in the CNT was sufficient to induce a severe salt-losing syndrome, confirming that the CNT is crucial for maintaining sodium and potassium balance.




Figure 6. Continuous loss of body weight in sodium-deprived CNT/CD-specific KO mice. KO ($n = 7$) and control mice ($n = 7$) are subjected to a sodium-deficient diet for 15 days in standard cages. Body weight is measured daily and is presented as percentage of the initial body weight (at day 0; $**P < 0.01$, $***P < 0.001$).

CONCISE METHODS

Generation of Transgenic Mice

To inactivate the *Scnn1a* gene in the CNT cells, we used mice expressing Cre recombinase under the control of the regulatory elements of the mouse *Aqp2* gene¹¹ and *Scnn1a*^{lox/lox} conditional

KO mice and the *Scnn1a*^{+/-} mice.^{8,12} CNT/CD-specific αENaC KO mice (*Scnn1a*^{lox/-}/*Aqp2::iCre*; KO group) and heterozygous (*Scnn1a*^{lox/+}/±*Aqp2::iCre* and *Scnn1a*^{lox/-}; control group) littermates were obtained by interbreeding *Scnn1a*^{+/-}/*Aqp2::iCre* mice with *Scnn1a*^{lox/lox} mice. Genotyping of the mice was performed by PCR analysis of tail biopsies described previously.¹² To genotype *Scnn1a*^{+/-} and *Aqp2::iCre* mice, the following primers were used: *Scnn1a*^{+/-}: primer #1: 5'-TTAAGGGTGCACACAGTGACGGC-3', #2: 5'-TTTGTACAGTCCTGCACGACGCG-3', and #3: 5'-AACTCCAGAAGTCCAGCTGGCTC-3', and *Aqp2::iCre*: #4: 5'-AAGTGCCACAGTCTAGCCTCT-3', #5: 5'-CCTGTTGTTCAGCTTGCACCAG-3', and #6: 5'-GGAGAACGCTATGACCCGAGT-3'.

Microdissection of Nephron Segments

Microdissection of kidneys was performed from KO mice ($n = 2$) and control littermates (*Scnn1a*^{lox/+} and *Scnn1a*^{lox/+}/*Aqp2::iCre*; $n = 2$, approximately 3 months of age). As additional controls, we included CD-specific KO mice (*Scnn1a*^{lox/lox}/*HoxB7::iCre*; $n = 2$) and their littermate controls (*Scnn1a*^{lox/lox}; $n = 2$, approximately 11 months of age).¹⁰ Microdissection was also performed on kidneys from KO mice ($n = 4$) and control littermates (*Scnn1a*^{lox/+}; $n = 4$, 2 to 2.5 months of age) that were subjected to a sodium-deficient diet in standard cages for 6 to 17 days. The experiment was performed four times and each time with one KO mouse and one control mouse in parallel. The kidney was perfused with DMEM/F-12 (1:1; 21041 medium; Invitrogen) completed with 40 μg/ml liberase blendzyme TM 2 or 30 μg/ml blendzyme (Roche Applied Science). Thin pyramids cut along the corticomedullary axis were incubated at 37 or 30°C for 40 minutes in the perfusion medium. The action of enzyme was stopped by washing the pyramids with ice-cold DMEM/F-12 (1:1) without blendzyme. Then, the medulla was removed under microscope and the CNT and CCD were microdissected in ice-cold DMEM/F-12 (1:1) without blendzyme. Pools of 10 to 20 microdissected tubules in 5 μl of DMEM/F12 (1:1) were then transferred to 5 μl of 2× concentrated protein sample buffer (9.6% [wt/vol] SDS, 13.8% [wt/vol] sucrose, 0.026% [wt/vol] bromphenol blue, and 4.2% [vol/vol] β-mercaptoethanol).




Figure 7. Retention of potassium in CNT/CD-specific KO mice upon potassium loading. KO mice ($n = 11$) and control mice ($n = 11$) are subjected to a high-potassium diet for 2 days. (A and B) Urinary potassium excretion (A) and urinary sodium excretion (B) are measured daily. $*P < 0.05$; $**P < 0.01$.

Western Blot Analysis of Microdissected Nephron Segments

Samples were heated at 95°C for 5 minutes and loaded and electrophoresed on an 8% SDS-PAGE. Proteins were then transferred to Protran nitrocellulose membrane (Schleicher & Schuell) or Amersham Hybond-ECL nitrocellulose membrane (Amersham), and Western blots were performed according to standard procedures. The membrane was first probed with affinity-purified αENaC antibody (1:100, 1:500, or 1:2000 dilution¹⁰); after stripping, anti-actin antibody (1:200 dilution; Sigma) was used. Blots were revealed with SuperSignal reagent (Pierce) or Amersham ECL Western blotting Detection Reagents (Amersham).

Immunohistochemistry

Kidneys from mice that were sodium-restricted for 4 consecutive days (*Scnn1a*^{lox/-}/*Aqp2::iCre* [$n = 2$] and *Scnn1a*^{lox/+}/*Aqp2::iCre* [$n = 3$]) were fixed by intravascular perfusion of 3% paraformaldehyde and subjected to paraffin embedding and sectioning (2-μm-thick sections).

Table 3. Functional data from mice subjected to a high-potassium diet for 48 hours

Parameter	KO	Control
Serum Na ⁺ (mM)	147.6 ± 1.4 (n = 8) ^a	151.5 ± 0.7 (n = 11)
Serum K ⁺ (mM)	7.7 ± 0.8 (n = 9) ^a	5.2 ± 0.2 (n = 11)
Serum HCO ₃ ⁻ (mM)	12.8 ± 1.1 (n = 7) ^b	18.3 ± 0.5 (n = 10)
Serum osm (mosm/kgH ₂ O)	329.0 ± 4.7 (n = 8)	325.0 ± 1.5 (n = 9)
Serum creatinine (mM)	19.1 ± 2.3 (n = 7) ^a	12.1 ± 1.5 (n = 10)
Serum Cl ⁻ (mM)	114.4 ± 1.4 (n = 7)	114.8 ± 0.7 (n = 10)
FE(Na ⁺) (%)	0.27 ± 0.05 (n = 7)	0.25 ± 0.02 (n = 10)

Data are means ± SEM.

^aP < 0.05.

^bP < 0.001.

Double Labeling

Double-labeling experiments were performed using (1) rabbit polyclonal Cre recombinase antibody (1:8000 dilution; Covance) and biotinylated rabbit polyclonal AQP2 antibody (7661AP; 1:250 dilution) and (2) Cre recombinase antibody (1:8000 dilution) and biotinylated rabbit polyclonal H⁺-ATPase antibody (H7659AP; 1:100 dilution²⁸). Sections were incubated overnight at 4°C with Cre antibody before undergoing incubation with horseradish peroxidase-conjugated goat anti-rabbit secondary antibody and visualization by 3,3'-diaminobenzidine (brown color). Sections were then incubated in 3.5% H₂O₂ in methanol to remove any remaining peroxidase from the first staining. After blocking for endogenous biotin (Biotin Blocking System; DakoCytomation), sections were then incubated overnight at 4°C with either biotinylated AQP2 or H⁺-ATPase antibodies. Labeling was visualized by use of Streptavidin horseradish peroxidase (Sigma) and Vector SG substrate (blue-gray color; Vector Laboratories).

Single Labeling of Serial Kidney Sections

Serial paraffin sections were incubated with (1) rabbit polyclonal αENaC antibody (1:800 dilution¹⁰), rabbit polyclonal AQP2 antibody (1:3000 dilution), and mouse monoclonal Calbindin D-28K antibody (1:20,000 dilution; Research Diagnostics) and (2) rabbit polyclonal TSC antibody (1:1000 dilution²⁹), rabbit polyclonal αENaC antibody (1:800 dilution¹⁰), and rabbit polyclonal AQP2 antibody (1:3000 dilution). Labeling was visualized by use of peroxidase-conjugated secondary antibody and 3,3'-diaminobenzidine. Sections were counterstained with hematoxylin. Light microscopy was carried out using a Leica DMRE microscope.

Quantification of αENaC-Positive Cells in the CNT

Cell counting was performed on kidney sections from *Scnn1a*^{lox/-}/*Aqp2::iCre* mice and *Scnn1a*^{lox/+}/*Aqp2::iCre* mice that were labeled with rabbit polyclonal αENaC antibodies and peroxidase-conjugated secondary antibodies. Counting was performed on electronic images taken with a ×25 objective. The number of αENaC-positive (labeled) and αENaC-negative (unlabeled) cells with a distinct nucleus was counted in the early CNT, adjacent to the transition from DCT to CNT (the transition was identified on serial sections labeled with TSC antibodies). The total number of cells counted was 65 in the *Scnn1a*^{lox/-}/*Aqp2::iCre* mice (n = 2) and 81 in the *Scnn1a*^{lox/+}/*Aqp2::iCre* mice (n = 2). Cellular counting was also performed in the late CNT (tubules were identified on serial sections labeled with AQP2 and TSC antibodies). The number of cells counted was 330 in the

Scnn1a^{lox/-}/*Aqp2::iCre* mice (n = 2) and 339 in the *Scnn1a*^{lox/+}/*Aqp2::iCre* mice (n = 3). The fraction of ENaC-positive cells was calculated from the number of positive cells divided by the total number of cells counted for each animal.

Quantification of Cre Recombinase-Positive Cells in the CCD and the CNT

The cellular counting was performed on kidney sections from *Scnn1a*^{lox/-}/*Aqp2::iCre* mice and *Scnn1a*^{lox/+}/*Aqp2::iCre* mice that were double labeled with rabbit polyclonal Cre recombinase antibody and biotinylated rabbit polyclonal AQP2 antibody. Counting was performed on electronic images taken with a ×63 objective. The number of AQP2-positive/Cre recombinase-positive and AQP2-positive/Cre recombinase-negative cells with a distinct nucleus was counted in the CCD (313 cells, n = 5 mice) and in the CNT (465 cells, n = 5 mice). The fraction of Cre recombinase-positive cells was calculated from the number of AQP2-positive/Cre recombinase-positive cells divided by the total number of AQP2-positive cells counted for each animal.

Experimental Protocols

Sodium-Deficient Diet in Metabolic Cages

For each metabolic cage study, experimental mice and controls from the same litter were used. Six- to 12-week-old mice were placed in individual metabolic cages (Tecniplast, Buguggiate, Italy) and fed a standard-salt diet (0.23% sodium; Institut National de la Recherche Agronomique, Unité de Préparation des Aliments Expérimentaux, Jouy en Josas, France) for 2 days, followed by 4 consecutive days on a sodium-deficient diet (0% sodium; Institut National de la Recherche Agronomique, Unité de Préparation des Aliments Expérimentaux). During the experiment, the animals had free access to food and water. The diet was given as a mixture of food in gelatin and water (100 g food/60 ml water). Blood was collected from the tail vein of conscious mice at the end of the diet. The experiment was also performed with 8- to 12-week-old mice fed a standard-salt diet (0.17% sodium, given as powder food; Ssniff Spezialdiäten GmbH, Soest, Germany) for 2 days followed by 3 days on a sodium-deficient diet (<0.01% sodium; Ssniff Spezialdiäten GmbH).

Sodium-Deficient Diet in Standard Cages

Experimental (n = 7) and control littermate mice (n = 7, 8 to 12 weeks old) were fed a standard-salt diet (0.17% sodium; Ssniff Spezialdiäten GmbH), and the body weight was measured at day 0 to determine the reference weight. Then, mice were fed a sodium-deficient diet (<0.01% sodium; Ssniff Spezialdiäten GmbH) with free access to water, and their body weight was monitored daily (at the same time) for 15 consecutive days.

Blood Collection for Aldosterone Measurements

Control and *Scnn1a*^{lox/lox}/*Aqp2::iCre* (KO) mice (8 to 12 weeks old) that were homozygous for a specific renin allele (*Ren-2*^{-/-}) were kept in standard cages with free access to food and water. Thirteen mice (eight control and five experimental) were fed a standard-salt diet

(0.17% sodium; Ssniff Spezialdiäten GmbH) and 10 mice (six control and four KO) were fed a sodium-deficient diet (<0.01% sodium; Ssniff Spezialdiäten GmbH) for 4 consecutive days. At the end of experiment, blood samples were collected after decapitation. Plasma aldosterone levels were measured according to standard procedures using a RIA (Coat-A-Count RIA kit; Siemens Medical Solutions Diagnostics, Ballerup, Denmark). Mouse samples with values >1200 pg/ml were further diluted using a serum pool with a low aldosterone concentration (<50 pg/ml).

High-Potassium Diet

Data from two separate experiments were pooled. In the first series of experiments, mice were 4 months old; in the second series, mice were 2.5 to 12 months old. Experimental mice ($n = 5$ for first experiment; $n = 6$ for second experiment) and control mice ($n = 6$ for first experiment; $n = 5$ for second experiment) were placed in individual metabolic cages and fed a standard diet for 2 consecutive days (0.59% potassium in the first experiment and 0.95% potassium in the second experiment). This was followed by 2 days on a 5% potassium diet (the potassium was added as KCl). During the experiment, the animals had free access to food and water. The diets were given as a mixture of food in gelatin and water (100 g food/60 ml water). After the diet, blood was collected from the eye.

Urine and Serum/Plasma Analysis

Urine and serum/plasma osmolarity as well as sodium, potassium, chloride, creatinine, and bicarbonate composition were analyzed at the Laboratoire Central de Chimie Clinique (Centre Hospitalier Universitaire Vaudoise, Lausanne, Switzerland). The potassium values were corrected for the degree of hemolysis.

BP Measurements

The BP and HR were measured in KO ($n = 13$) and control mice ($n = 19$). The mice were kept on a normal-salt diet containing 0.23% of sodium with free access to tap water and were analyzed at the age of 4 to 6 months. BP and HR were recorded intra-arterially using a computerized data-acquisition system (Notocord Systems SA, Croissy, France).¹⁷ Briefly, for placement of the intra-arterial catheter, a mouse was anesthetized *via* inhalation of 1 to 2% halothane with oxygen. The right carotid artery was exposed for a length of approximately 4 mm. A silicone/PE10 catheter filled with 0.9% NaCl solution containing heparin (300 IU/ml) was inserted into the artery. After ligation, the catheter was subcutaneously tunneled to exit at the back of the neck and fixed with a piece of scotch tape and dental cement. The mouse was allowed 3 to 4 hours to recover from the anesthesia and placed in a Plexiglas tube for partial restriction of its movements. Thirty minutes later, the arterial line was connected to a pressure transducer; BP and HR were then monitored every 20 seconds for 15 to 20 minutes using the Notocord computerized data-acquisition system at a sampling rate of 500 Hz. Once BP measurement was completed, blood was sampled from the arterial catheter for analysis of serum sodium and potassium concentrations.

Statistical Analysis

Results are presented as mean \pm SEM. Data were analyzed by one-way ANOVA and unpaired *t* test. $P < 0.05$ was considered statistically significant.

ACKNOWLEDGMENTS

Financial support for this study was provided by the Swiss National Science Foundation, the Danish Medical Research Council, the Danish National Research Foundation, and the Leducq Foundation.

We are very grateful to Johannes Loffing (University of Fribourg/University of Zürich, Switzerland) for help with perfusion of the mouse kidneys. We also thank Nicole Fowler Jaeger, Anne-Marie Mérillat, Sandrine Egli, and Inger-Merete Paulsen for excellent technical assistance and Friedrich Beermann for critically reading the manuscript. We also thank Søren Nielsen (Water and Salt Research Center, University of Aarhus, Aarhus, Denmark) for providing the antibodies against AQP2 and H⁺-ATPase and Mark Knepper (National Institutes of Health, Bethesda, MD) for providing the antibody against TSC.

DISCLOSURES

None.

REFERENCES

- Verrey F, Hummler E, Schild L, Rossier BC: Mineralocorticoid action in the aldosterone-sensitive distal nephron. In: *The Kidney Physiology and Pathophysiology*, 4th Ed., edited by Alpern RJ, Hebert SC, Academic Press, 2008, pp 889–924
- Loffing J, Loffing-Cueni D, Valderrabano V, Klausli L, Hebert SC, Rossier BC, Hoenderop JG, Bindels RJ, Kaissling B: Distribution of transcellular calcium and sodium transport pathways along mouse distal nephron. *Am J Physiol Renal Physiol* 281: F1021–F1027, 2001
- Shimkets RA, Warnock DG, Bositis CM, Nelson-Williams C, Hansson JH, Schambelan M, Gill JR Jr, Ulick S, Milora RV, Findling JW: Liddle's syndrome: Heritable human hypertension caused by mutations in the beta subunit of the epithelial sodium channel. *Cell* 79: 407–414, 1994
- Hansson JH, Nelson-Williams C, Suzuki H, Schild L, Shimkets R, Lu Y, Canessa C, Iwasaki T, Rossier B, Lifton RP: Hypertension caused by a truncated epithelial sodium channel gamma subunit: Genetic heterogeneity of Liddle syndrome. *Nat Genet* 11: 76–82, 1995
- Chang SS, Grunder S, Hanukoglu A, Rosler A, Mathew PM, Hanukoglu I, Schild L, Lu Y, Shimkets RA, Nelson-Williams C, Rossier BC, Lifton RP: Mutations in subunits of the epithelial sodium channel cause salt wasting with hyperkalaemic acidosis, pseudohypoaldosteronism type 1. *Nat Genet* 12: 248–253, 1996
- Kuhnle U: Pseudohypoaldosteronism: mutation found, problem solved? *Mol Cell Endocrinol* 133: 77–80, 1997
- Barker PM, Nguyen MS, Gatzky JT, Grubb B, Norman H, Hummler E, Rossier B, Boucher RC, Koller B: Role of gammaENaC subunit in lung liquid clearance and electrolyte balance in newborn mice: Insights into perinatal adaptation and pseudohypoaldosteronism. *J Clin Invest* 102: 1634–1640, 1998
- Hummler E, Barker P, Gatzky J, Beermann F, Verdumo C, Schmidt A, Boucher R, Rossier BC: Early death due to defective neonatal lung

- liquid clearance in alpha-ENaC-deficient mice. *Nat Genet* 12: 325–328, 1996
9. McDonald FJ, Yang B, Hrstka RF, Drummond HA, Tarr DE, McCray PB Jr, Stokes JB, Welsh MJ, Williamson RA: Disruption of the beta subunit of the epithelial Na⁺ channel in mice: Hyperkalemia and neonatal death associated with a pseudohypoaldosteronism phenotype. *Proc Natl Acad Sci U S A* 96: 1727–1731, 1999
 10. Rubera I, Loffing J, Palmer LG, Frindt G, Fowler-Jaeger N, Sauter D, Carroll T, McMahon A, Hummler E, Rossier BC: Collecting duct-specific gene inactivation of alphaENaC in the mouse kidney does not impair sodium and potassium balance. *J Clin Invest* 112: 554–565, 2003
 11. Ronzaud C, Loffing J, Bleich M, Gretz N, Grone HJ, Schutz G, Berger S: Impairment of sodium balance in mice deficient in renal principal cell mineralocorticoid receptor. *J Am Soc Nephrol* 18: 1679–1687, 2007
 12. Hummler E, Merillat AM, Rubera I, Rossier BC, Beermann F: Conditional gene targeting of the Scnn1a (alphaENaC) gene locus. *Genesis* 32: 169–172, 2002
 13. Nielsen S, DiGiovanni SR, Christensen EI, Knepper MA, Harris HW: Cellular and subcellular immunolocalization of vasopressin-regulated water channel in rat kidney. *Proc Natl Acad Sci U S A* 90: 11663–11667, 1993
 14. Kishore BK, Mandon B, Oza NB, DiGiovanni SR, Coleman RA, Ostrowski NL, Wade JB, Knepper MA: Rat renal arcade segment expresses vasopressin-regulated water channel and vasopressin V2 receptor. *J Clin Invest* 97: 2763–2771, 1996
 15. Ergonul Z, Frindt G, Palmer LG: Regulation of maturation and processing of ENaC subunits in the rat kidney. *Am J Physiol Renal Physiol* 291: F683–F693, 2006
 16. Hughey RP, Bruns JB, Kinlough CL, Harkleroad KL, Tong Q, Carattino MD, Johnson JP, Stockand JD, Kleyman TR: Epithelial sodium channels are activated by furin-dependent proteolysis. *J Biol Chem* 279: 18111–18114, 2004
 17. Wang Q, Hummler E, Nussberger J, Clement S, Gabbiani G, Brunner HR, Burnier M: Blood pressure, cardiac, and renal responses to salt and deoxycorticosterone acetate in mice: Role of Renin genes. *J Am Soc Nephrol* 13: 1509–1516, 2002
 18. Charles RP, Guitard M, Leyvraz C, Breiden B, Haftek M, Haftek-Terreau Z, Stehle JC, Sandhoff K, Hummler E: Postnatal requirement of the epithelial sodium channel for maintenance of epidermal barrier function. *J Biol Chem* 283: 2622–2630, 2008
 19. Coleman RA, Wu DC, Liu J, Wade JB: Expression of aquaporins in the renal connecting tubule. *Am J Physiol Renal Physiol* 279: F874–F883, 2000
 20. Hofmeister MV, Fenton RA, Praetorius J: Fluorescence isolation of mouse late distal convoluted tubules and connecting tubules: Effects of vasopressin and vitamin D3 on Ca²⁺ signaling. *Am J Physiol Renal Physiol* 296: F194–F203, 2009
 21. Makhanova N, Lee G, Takahashi N, Sequeira Lopez ML, Gomez RA, Kim HS, Smithies O: Kidney function in mice lacking aldosterone. *Am J Physiol Renal Physiol* 290: F61–F69, 2006
 22. Pradervand S, Barker PM, Wang Q, Ernst SA, Beermann F, Grubb BR, Burnier M, Schmidt A, Bindels RJ, Gatzky JT, Rossier BC, Hummler E: Salt restriction induces pseudohypoaldosteronism type 1 in mice expressing low levels of the beta-subunit of the amiloride-sensitive epithelial sodium channel. *Proc Natl Acad Sci U S A* 96: 1732–1737, 1999
 23. Meneton P, Loffing J, Warnock DG: Sodium and potassium handling by the aldosterone-sensitive distal nephron: The pivotal role of the distal and connecting tubule. *Am J Physiol Renal Physiol* 287: F593–F601, 2004
 24. Loffing J, Korbmayer C: Regulated sodium transport in the renal connecting tubule (CNT) via the epithelial sodium channel (ENaC). *Pflugers Arch* 458: 111–135, 2009
 25. Malnic G, Klose RM, Giebisch G: Micropuncture study of renal potassium excretion in the rat. *Am J Physiol* 206: 674–686, 1964
 26. Frindt G, Palmer LG: Na channels in the rat connecting tubule. *Am J Physiol Renal Physiol* 286: F669–F674, 2004
 27. Loffing J, Pietri L, Aregger F, Bloch-Faure M, Ziegler U, Meneton P, Rossier BC, Kaissling B: Differential subcellular localization of ENaC subunits in mouse kidney in response to high- and low-Na diets. *Am J Physiol Renal Physiol* 279: F252–F258, 2000
 28. Christensen BM, Kim YH, Kwon TH, Nielsen S: Lithium treatment induces a marked proliferation of primarily principal cells in rat kidney inner medullary collecting duct. *Am J Physiol Renal Physiol* 291: F39–F48, 2006
 29. Biner HL, Arpin-Bott MP, Loffing J, Wang X, Knepper M, Hebert SC, Kaissling B: Human cortical distal nephron: Distribution of electrolyte and water transport pathways. *J Am Soc Nephrol* 13: 836–847, 2002

Supplemental information for this article is available online at <http://www.jasn.org/>.