

VHL Inactivation Induces HEF1 and Aurora Kinase A

Jianyong Xu,* Huapeng Li,* Bo Wang,* Yan Xu,* Jiayin Yang,* Xiaofei Zhang,* Sarah K. Harten,[†] Deepa Shukla,[†] Patrick H. Maxwell,[†] Duanqing Pei,* and Miguel A. Esteban*

*Key Laboratory of Regenerative Biology, South China Institute for Stem Cell Biology and Regenerative Medicine, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou, China; and [†]Centre for Cell Signaling and Molecular Genetics, Rayne Institute, University College London, London, United Kingdom

ABSTRACT

The ciliary hypothesis for cystic renal diseases postulates that most of these conditions result from abnormalities in the primary cilium, a microtubule-based structure that acts as a sensor for extracellular cues. Inactivation of the von Hippel-Lindau (*VHL*) tumor suppressor gene predisposes to renal cysts and clear cell renal cell carcinoma. *VHL* plays a critical role in the formation of primary cilia in kidney epithelium, but the underlying mechanisms are poorly understood. Here, we demonstrate that *VHL* inactivation induces HEF1/Cas-L/NEDD9 and Aurora kinase A via the stabilization of hypoxia-inducible factors 1 and 2. Aurora kinase A is a mitotic kinase commonly upregulated in cancer that causes regression of the primary cilium by promoting histone deacetylase-dependent tubulin depolymerization of the ciliary axoneme. HEF1/Cas-L/NEDD9 is a component of focal adhesions that has a prominent role in inducing metastasis and that colocalizes with Aurora kinase A at the centrosome, thereby enhancing the harmful effect of Aurora kinase A on the cilium. Suppression of this pathway improved the formation of primary cilia and reduced cell motility in *VHL*-defective renal cancer cells. Our results highlight the gatekeeper role of *VHL* in the kidney epithelium.

J Am Soc Nephrol 21: 2041–2046, 2010. doi: 10.1681/ASN.2010040345

Primary cilia are single hair-like organelles on the surface of most growth-arrested or differentiated mammalian cells.¹ They are microtubule-containing projections that work as mechanosensors, chemosensors, and osmosensors. Compared with related structures such as respiratory cilia, primary cilia lack the central pair of microtubules and are therefore static. Nevertheless, primary cilia activate multiple signals that influence proliferation, survival, cell polarity, and motility. This variety of functions is caused by specific proteins being located on or near the ciliary membrane, including the polycystins and components of the Hedgehog/Wnt signaling pathways.¹ The regulation of ciliary length is dictated by a bidirectional process of trafficking termed

intraflagellar transport, responsible for renewing components at the tip of cilia.² The centrosome is another point of regulation as cilia originate from microtubules nucleated by one of the two centrioles. In this regard, Golemis and collaborators showed that phosphorylation of histone deacetylase 6 (HDAC6) at the centrosome by Aurora kinase A (Aurka) induces tubulin deacetylation and destabilizes the ciliary axoneme.³ This is independent of Aurka activity on mitosis and is potentiated by NEDD9/Cas-L/HEF1 (hereafter referred to as NEDD9), which colocalizes with Aurka at the centrosome.³ NEDD9 also colocalizes with integrins at focal adhesions and regulates cell motility.⁴

Defects in genes involved in primary cilia assembly or function produce a series of diseases, termed ciliopathies, which have the formation of visceral cysts as unifying feature.^{1,5} Among other ciliopathies are autosomal dominant polycystic kidney disease, nephronophthisis, Bardet-Biedl disease, and von Hippel-Lindau (*VHL*) disease. The latter is an autosomal dominant syndrome characterized by cysts, especially in the pancreas and kidney, and a variety of tumors including clear cell renal cell carcinomas (CCRCCs) and pheochromocytomas.⁶ In accordance with Knudson's model for tumor suppressors, the development of pathology in *VHL* patients is linked to somatic inactivation of the remaining wild-type *VHL* allele in a susceptible cell type (e.g., kidney epithelial cells). Also, the kidneys of *VHL* patients have large numbers of early "second-hit" lesions that presumably precede the formation of cysts and CCRCCs.⁷ Importantly, *VHL* is also mu-

Received April 3, 2010. Accepted July 21, 2010.

Published online ahead of print. Publication date available at www.jasn.org.

J.X. and H.L. contributed equally to this work.

Correspondence: Dr. Miguel A. Esteban and Dr. Duanqing Pei, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China. Phone: 86-20-3205-1820; Fax: 86-20-3201-5299; E-mails: esteban@gibh.org and pei_duanqing@gibh.ac.cn

Copyright © 2010 by the American Society of Nephrology

tated in the majority of sporadic CCRCCs.⁶

The most studied function of VHL protein (pVHL) is the regulation of cellular responses to altered oxygenation through controlling the degradation of the α subunits of the transcription factor hypoxia inducible factor (HIF).⁸ In the presence of oxygen, HIF- α subunits are hydroxylated by the HIF prolyl hydroxylases,⁹ and this generates a binding site for pVHL. Under hypoxia or when pVHL is inactivated, stabilized HIF translocates to the nucleus, binds HIF- β , and induces many genes.^{6,8} pVHL has additional HIF-unrelated functions such as controlling extracellular matrix assembly.¹⁰ We and others have also reported that pVHL regulates primary cilia through both HIF-dependent and -independent mechanisms.^{11–14} Better understanding of how pVHL exerts this role may provide ways to prevent or cure CCRCCs.

Here we hypothesized that activation of the NEDD9–AurkA pathway may underlie an HIF-negative effect on the cilium in *VHL*-defective CCRCC cells. Supporting this idea, AurkA is a HIF-1 α target¹⁵ and is up-regulated in CCRCCs,¹⁶ although an association with *VHL* status had not yet been established. Moreover, NEDD9 was originally termed renal carcinoma antigen NY-REN-1,¹⁷ and previous studies showed that NEDD9 mRNA is induced by hypoxia in CD133+ cells¹⁸ and by pVHL inactivation in CCRCC cells.¹⁹ To test this, we generated pools of the *VHL*-defective CCRCC cell line RCC10 that overexpress the longer isoform of pVHL (pVHL30) or control empty vector. Re-expression of *VHL* reduced HIF, NEDD9, and AurkA, as analyzed by Western blotting and real-time RT-PCR (qPCR; Figure 1, A and B). The well-known HIF- α target VEGF and the *VHL*-regulated tight junction protein claudin 1²⁰ were used as qPCR controls. Both isoform 1 and 2 of NEDD9 were down-regulated by pVHL at the mRNA level, but the closely related p130CAS showed no differential effect (Figure 1C). Regulation of NEDD9/AurkA by pVHL was




Figure 1. Inactivation of *VHL* induces NEDD9 and AurkA in CCRCC cells. (A) Western blot using the indicated antibodies and lysates from two different pools of *VHL*-defective RCC10 cells infected with a retrovirus producing pVHL30 or control empty vector. Tubulin was used as loading control (also hereafter). (B) qPCR for the indicated targets in the same RCC10 pools; mean values of three independent experiments and the SD are shown (this applies hereafter to all data containing error bars). * $P < 0.05$ as analyzed by *t* test (also applies hereafter). (C) qPCR for NEDD9 isoforms 1 and 2 and the NEDD9-related molecule p130CAS in the same RCC10 pools. (D) Left: immunofluorescence photographs of *VHL*-defective RCC10 pools infected with empty vector or *VHL*30 and labeled for AurkA. Magnified images of the same field (only for *VHL*-defective cells) show AurkA staining cells at different stages of the cell cycle. Scale bars correspond to 10 μ m. Right: quantification of AurkA positive cells in three immunofluorescence experiments (performed in triplicate and counting 10 fields on each coverslip) in *VHL*-defective RCC10 pools infected with retroviruses producing pVHL30 or empty vector. ** $P < 0.01$ (also hereafter). (E) Western blot using the indicated antibodies (left) and qPCR for the indicated genes (right); two different pools of *VHL*-defective RCC4 cells were infected as above. (F) Western blot using the indicated antibodies and lysates from paired CCRCC tumor samples and surrounding normal tissue of two patients with *VHL*-defective sporadic CCRCC.

additionally shown by immunofluorescence in RCC10 cells (Figure 1D) and by Western blotting and qPCR in RCC4 cells (Figure 1E). Western blotting of paired tumor and normal tissue samples from patients with *VHL*-de-

fective sporadic CCRCC further confirmed these observations (Figure 1F). We studied whether activation of NEDD9 and AurkA in *VHL*-defective RCC10 cells was HIF mediated. We overexpressed a series of previously re-

ported disease-associated pVHL mutants that retain different abilities to down-regulate HIF.^{6,21} The shorter isoform of pVHL (pVHL19), which has the ability to down-regulate HIF but lacks other functions associated to pVHL30,²² was used as an additional control. Type 2C pVHL variants are associated with high risk of pheochromocytoma *in vivo* but no other manifestations, whereas type 1 and 2B pVHL variants associate with renal manifestations.⁶ Type 2C mutants effectively down-regulated HIF- α subunits, NEDD9 and Aurka, but types 1 and 2B did not (Figure 2A). Next we transfected *VHL*-defective RCC10 cells with small interfering RNA (siRNA) oligos for HIF-1 α and/or HIF-2 α and observed that Aurka is regulated by both HIF- α subunits, whereas NEDD9 is regulated only by HIF-2 α (Figure 2B; Supplementary Figure 1); siRNA oligos against the firefly luciferase gene were the control. HIF-2 α and NEDD9 were up-regulated by HIF-1 α siRNA. HIF-2 α siRNA also moderately increased HIF-1 α , consistent with previous reports showing that HIF- α subunits influence each other.²³ The regulatory effect of HIF-2 α on NEDD9 and Aurka was verified overexpressing a constitutively stable HIF-2 α that bears mutations in the proline residues hydroxylated by the HIF prolyl hydroxylases^{24,25} (Figure 2C). Moreover, of four putative hypoxia responsive elements in the NEDD9 proximal promoter, chromatin immunoprecipitation (ChIP) showed specific binding of HIF-2 α to region 4 (-633 to 628 upstream of the transcription initiation; Figure 2D). This was validated by cotransfection of a constitutively stable HIF-2 α and a luciferase reporter plasmid containing the NEDD9 promoter with or without a mutation (GCGTG to GTTTG) in this hypoxia responsive element (Figure 2E).

We evaluated the contribution of NEDD9 and Aurka to primary cilia abnormalities in RCC10 cells. Overexpression of NEDD9 or Aurka reduced cilia formation in *VHL*-defective RCC10 cells stably transfected with *VHL* (RCC10/*VHL*) (Figure 3, A and B); increased expression of NEDD9 and Aurka was verified by Western blot-




Figure 2. Stabilization of HIF mediates the induction of NEDD9 and Aurka in *VHL*-defective RCC10 cells. (A) Western blot using the indicated antibodies and lysates from *VHL*-defective RCC10 pools infected with retroviruses producing pVHL19, disease specific-pVHL mutants, or empty vector. (B) Western blot using the indicated antibodies and lysates from *VHL*-defective RCC10 cells transfected with siRNA oligos (oligo pairs A, see Supplementary Methods) for HIF-1 α , HIF-2 α , and the firefly luciferase gene (control); two different pools were transfected in parallel. Ctrl indicates control (also hereafter). (C) Western blot using the indicated antibodies and lysates from two different pools of an RCC10 cell clone stably expressing *VHL*²⁵ (RCC10/*VHL*) infected with retroviruses producing a constitutively stable form of HIF-2 α or empty vector. (D) Top: schematic representation of putative hypoxia responsive element sites in the proximal NEDD9 promoter. Bottom: semiquantitative RT-PCR with ChIP samples obtained using HIF-2 α or control antibodies; diluted chromatin (1:10) was used for the input. (E) Left: activity of the same promoter cloned upstream of the luciferase gene into pGL3Basic and cotransfected into HEK293T cells together with empty pcDNA3 vector or constitutively stable forms of HIF-1 α (HIF-1 α DM) and HIF-2 α (HIF-2 α DM). Right: activity of the same promoter containing mutations in the indicated putative hypoxia responsive element sites and cotransfected in HEK293T cells with empty pcDNA3 vector or a constitutively stable HIF-2 α . WT, wild type; MUT, mutated. GCGTG sites were mutated to GTTTG using site-directed mutagenesis.

ting. Conversely, two small molecule Aurka inhibitors or the HDAC inhibitor trichostatin (TSA) increased the number of cilia in *VHL*-defective RCC10 cells com-

pared with the control (Figure 3, A and B). siRNA for NEDD9 effectively increased cilia formation in *VHL*-defective RCC10 cells as well (Figure 3C), and the combined




Figure 3. Manipulation of the NEDD9–AurkA pathway influences primary cilia formation in *VHL*-defective RCC10 cells. (A) Left: Western blot showing increased expression of AurkA and NEDD9 in RCC10/*VHL* cells infected with AurkA and NEDD9 respectively. Right: quantification of primary cilia in RCC10/*VHL* cells overexpressing AurkA, NEDD9, or both, and in *VHL*-defective RCC10 cells treated for 72 hours with VX680 (250 nM), ZM447439 (2.5 μ M), or TSA (100 nM). (B) Immunofluorescence photographs of *VHL*-defective RCC10 or RCC10/*VHL* cells infected or treated as indicated; scale bar corresponds to 10 μ m. Values correspond to three independent experiments done in duplicate—16 fields were counted in each coverslip; this applies hereafter to all related experiments. (C) Left: Western blot using lysates from *VHL*-defective RCC10 cells transiently transfected with siRNA oligos for NEDD9 or control. Right: quantification of primary cilia in *VHL*-defective RCC10 cells transfected with siRNA for HIF-1 α (100 nM), HIF-2 α (100 nM), HIF-1 α (50 nM) + HIF-2 α (50 nM), and NEDD9 (100 nM) compared with control (100 nM).

knock down of both HIF- α subunits was more effective than HIF-1 α alone (Figure 3C). This cooperative effect highlights a

role for HIF-2 α without contradicting our previous findings that *VHL* inactivation alters cilia mainly through HIF-1 α .¹¹ The

fact that HIF-2 α siRNA alone has no effect on ciliogenesis may be explained by the incomplete reduction of NEDD9 expression compared with NEDD9 siRNA (Figures 2B and 3C; Supplementary Figure 1). The latter may imply that additional mechanisms regulate NEDD9 in *VHL*-defective cells, but it could also be a threshold effect. Besides, it is possible that up-regulation of HIF-1 α triggered by HIF-2 α counteracts cilia formation indirectly. Notably, under the conditions tested, we did not detect any change in cell proliferation after overexpression of NEDD9/AurkA or treatment with inhibitors and siRNA oligos (Supplementary Figure 2, A–C), reinforcing the idea that NEDD9/AurkA affect ciliogenesis independently of cell mitosis.³ Additionally, because NEDD9 is also an important regulator of cell motility, we studied whether manipulating its expression would influence the motility of RCC10 cells. As reported for other CCRCC cell lines,²⁶ *VHL*-defective RCC10 cells expressing pVHL30 were less motile than the control, which we assessed using wound healing and Transwell assays (Figure 4A). We then observed that overexpressing NEDD9 increased the motility of *VHL*-defective RCC10 expressing pVHL30 and knock down of NEDD9 or HIF-2 α reduced the motility of *VHL*-defective RCC10 expressing the empty vector (Figure 4, B–E). The effects on cell motility seemed independent of cell proliferation as well (Supplementary Figure 2D).

In summary, our data show that the NEDD9–AurkA pathway is induced through HIF in CCRCC cells and has significant impact on two relevant features of *VHL*-defective cells: the suppression of primary cilia that *in vivo* can lead to premalignant cysts and the increased motility that can lead to metastasis. This supports the idea that AurkA and HDAC inhibitors may be useful in treating sporadic CCRCC and *VHL* disease. Indeed, AurkA is a promising therapeutic target in cancer in general,²⁷ and TSA is already used in the clinic for unrelated malignancies.²⁸ Our findings may also have broader implications, because activation of HIF is common in cancer because of oncogenic signaling or abnormal oxygenation.⁸ This could explain why, in




Figure 4. NEDD9-Aurka regulate motility in *VHL*-defective RCC10 cells. (A) Left: quantification of gap closure in three independent experiments (measured in triplicate) after wounding monolayers of *VHL*-defective RCC10 cells (infected with *VHL30* or empty vector) with a tip edge (cells were washed and fresh medium added afterward). Right: cell migration through Transwell filters using similar RCC10 cell populations. (B) Selected phase contrast photographs of a similar experiment using *VHL*-defective RCC10 cells treated as indicated. (C) Same as in B but using RCC10/*VHL* cells infected with NEDD9 or empty vector. (D) Quantification of three independent experiments as in B and C measured in triplicate. (E) Quantification of cell migration through Transwell filters (three experiments in triplicate) using *VHL*-defective RCC10 cells or RCC10/*VHL* cells transfected and infected as indicated.

some contexts, cancer cells lose primary cilia or metastasize. Notably, the pollutant-responsive transcription factor, aryl hydrocarbon receptor, which like HIF

dimerizes with HIF- β s (also termed ARNT), also increases NEDD9 expression.²⁹ While our work was under review, Dubois and collaborators³⁰ re-

ported that HIF- α can induce NEDD9 in colon cancer cells. In the future, it will be interesting to determine whether other mechanisms known to affect ciliogenesis are regulated by *VHL* in a HIF-dependent or -independent manner.

CONCISE METHODS

Cell Culture and Plasmids

VHL-defective RCC10 and RCC4 CCRCC lines and RCC10/*VHL* (clone 90) were reported previously.¹¹ Experimental tissue culture conditions are described in the Supplementary Information. Retroviral infections were performed as described²⁵ but using HEK293T and a complementary packaging vector (pCL) for viral production. pBABE retroviral vectors expressing human NEDD9 (isoform 1, NM_006403.3) and Aurka (isoform 1, NM_198433.1) were prepared using cDNA from *VHL*-defective RCC10 cells as a template and confirmed by sequencing. Other retroviral vectors including constitutively stable versions of HIF-1 α (carrying mutations at prolines 402 and 564) and HIF-2 α (prolines 405 and 531) have been described before.¹¹ Aurka inhibitors ZM447439 and VX680 were purchased from Tocris Bioscience and Kava Tech, respectively, and diluted in DMSO; TSA was purchased from Sigma and also diluted in DMSO. Uncoated Transwell filters were purchased from Millipore; cell migration (4×10^5 cells were added in the top chamber) was allowed to proceed for 2 hours before fixing cells with methanol and cleaning the upper chamber.

Antibodies

Monoclonal anti-NEDD9 was purchased from Abcam, rabbit polyclonal anti-HIF1 α and anti-HIF2 α was purchased from BD Biosciences and Novus Biologicals, respectively, rabbit polyclonal anti-Aurka was purchased from Cell Signaling, and mouse monoclonal anti-acetylated tubulin was purchased from Sigma. Immunofluorescence studies were performed as described,²⁵ and a Leica TCS SP2 Spectral Confocal System microscope was used for detection. Western blotting was performed using ECL Plus (Amersham).

siRNA Transfection, qPCR, and ChIP

siRNA oligos for HIF-1 α , HIF-2 α , and NEDD9 are described in the Supplementary Information. All siRNA oligos were purchased from Guangzhou Ribobio. Cells were transfected using LipofectAMINE 2000 (Invitrogen) and a reverse transfection protocol as indicated by the manufacturer.

qPCR was performed using SYBR Green (Takara) and an ABI7300 machine. Samples were measured in triplicate and normalized on the basis of β -actin values. For ChIP, we followed the X-ChIP protocol from Abcam; ChIP grade anti-HIF-2 α was also from Abcam. qPCR and ChIP primer sequences are listed in the Supplementary Information.

ACKNOWLEDGMENTS

This work was supported by Chinese Academy of Sciences Grant KSCX2-YW-R-244 and start up funding from the Guangzhou Institutes of Biomedicine and Health to M.A.E.

DISCLOSURES

None.

REFERENCES

- Gerdes JM, Davis EE, Katsanis N: The vertebrate primary cilium in development, homeostasis, and disease. *Cell* 137: 32–45, 2009
- Rosenbaum JL, Witman GB: Intraflagellar transport. *Nat Rev Mol Cell Biol* 3: 813–825, 2002
- Pugacheva EN, Jablonski SA, Hartman TR, Henske EP, Golemis EA: HEF1-dependent Aurora A activation induces disassembly of the primary cilium. *Cell* 129: 1351–1363, 2007
- Sattler M, Salgia R, Shrikhande G, Verma S, Uemura N, Law SF, Golemis EA, Griffin JD: Differential signaling after beta1 integrin ligation is mediated through binding of CRKL to p120(CBL) and p110(HEF1). *J Biol Chem* 272: 14320–14326, 1997
- Pazour GJ: Intraflagellar transport and cilia-dependent renal disease: The ciliary hypothesis of polycystic kidney disease. *J Am Soc Nephrol* 15: 2528–2536, 2004
- Kaelin WG Jr: Molecular basis of the VHL hereditary cancer syndrome. *Nat Rev Cancer* 2: 673–682, 2002
- Mandriota SJ, Turner KJ, Davies DR, Murray PG, Morgan NV, Sowter HM, Wykoff CC, Maher ER, Harris AL, Ratcliffe PJ, Maxwell PH: HIF activation identifies early lesions in VHL kidneys: Evidence for site-specific tumor suppressor function in the nephron. *Cancer Cell* 1: 459–468, 2002
- Zhang H, Semenza GL: The expanding universe of hypoxia. *J Mol Med* 86: 739–746, 2008
- Schofield CJ, Ratcliffe PJ: Oxygen sensing by HIF hydroxylases. *Nat Rev Mol Cell Biol* 5: 343–354, 2004
- Ohh M, Yauch RL, Lonergan KM, Whaley JM, Stemmer-Rachamimov AO, Louis DN, Gavin BJ, Kley N, Kaelin WG Jr, Iliopoulos O: The von Hippel-Lindau tumor suppressor protein is required for proper assembly of an extracellular fibronectin matrix. *Mol Cell* 1: 959–968, 1998
- Esteban MA, Harten SK, Tran MG, Maxwell PH: Formation of primary cilia in the renal epithelium is regulated by the von Hippel-Lindau tumor suppressor protein. *J Am Soc Nephrol* 17: 1801–1806, 2006
- Lutz MS, Burk RD: Primary cilium formation requires von hippel-lindau gene function in renal-derived cells. *Cancer Res* 66: 6903–6907, 2006
- Schermer B, Ghenoiu C, Bartram M, Muller RU, Kotsis F, Hohne M, Kuhn W, Rapka M, Nitschke R, Zentgraf H, Fliegau M, Omran H, Walz G, Benzing T: The von Hippel-Lindau tumor suppressor protein controls ciliogenesis by orienting microtubule growth. *J Cell Biol* 175: 547–554, 2006
- Thoma CR, Frew IJ, Hoerner CR, Montani M, Moch H, Krek W: pVHL and GSK3beta are components of a primary cilium-maintenance signalling network. *Nat Cell Biol* 9: 588–595, 2007
- Klein A, Flugel D, Kietzmann T: Transcriptional regulation of serine/threonine kinase-15 (STK15) expression by hypoxia and HIF-1. *Mol Biol Cell* 19: 3667–3675, 2008
- Kurahashi T, Miyake H, Hara I, Fujisawa M: Significance of Aurora-A expression in renal cell carcinoma. *Urol Oncol* 25: 128–133, 2007
- Scanlan MJ, Gordan JD, Williamson B, Stockert E, Bander NH, Jongeneel V, Gure AO, Jager D, Jager E, Knuth A, Chen YT, Old LJ: Antigens recognized by autologous antibody in patients with renal-cell carcinoma. *Int J Cancer* 83: 456–464, 1999
- Martin-Rendon E, Hale SJ, Ryan D, Baban D, Forde SP, Roubelakis M, Sweeney D, Moukayed M, Harris AL, Davies K, Watt SM: Transcriptional profiling of human cord blood CD133+ and cultured bone marrow mesenchymal stem cells in response to hypoxia. *Stem Cells* 25: 1003–1012, 2007
- Harten SK, Esteban MA, Maxwell PH: Identification of novel VHL regulated genes by transcriptomic analysis of RCC10 renal carcinoma cells. *Adv Enzyme Regul* 49: 43–52, 2009
- Harten SK, Shukla D, Barod R, Hergovich A, Balda MS, Matter K, Esteban MA, Maxwell PH: Regulation of renal epithelial tight junctions by the von Hippel-Lindau tumor suppressor gene involves occludin and claudin 1 and is independent of E-cadherin. *Mol Biol Cell* 20: 1089–1101, 2009
- Hergovich A, Lisztwan J, Barry R, Ballschmieter P, Krek W: Regulation of microtubule stability by the von Hippel-Lindau tumour suppressor protein pVHL. *Nat Cell Biol* 5: 64–70, 2003
- Lolkema MP, Gervais ML, Sniijckers CM, Hill RP, Giles RH, Voest EE, Ohh M: Tumor suppression by the von Hippel-Lindau protein requires phosphorylation of the acidic domain. *J Biol Chem* 280: 22205–22211, 2005
- Raval RR, Lau KW, Tran MG, Sowter HM, Mandriota SJ, Li JL, Pugh CW, Maxwell PH, Harris AL, Ratcliffe PJ: Contrasting properties of hypoxia-inducible factor 1 (HIF-1) and HIF-2 in von Hippel-Lindau-associated renal cell carcinoma. *Mol Cell Biol* 25: 5675–5686, 2005
- Kondo K, Kim WY, Lechpammer M, Kaelin WG Jr: Inhibition of HIF2alpha is sufficient to suppress pVHL-defective tumor growth. *PLoS Biol* 1: E83, 2003
- Esteban MA, Tran MG, Harten SK, Hill P, Castellanos MC, Chandra A, Raval R, O'Brien TS, Maxwell PH: Regulation of E-cadherin expression by VHL and hypoxia-inducible factor. *Cancer Res* 66: 3567–3575, 2006
- Kamada M, Suzuki K, Kato Y, Okuda H, Shuin T: von Hippel-Lindau protein promotes the assembly of actin and vinculin and inhibits cell motility. *Cancer Res* 61: 4184–4189, 2001
- Ochi T, Fujiwara H, Yasukawa M: Aurora-A kinase: A novel target both for cellular immunotherapy and molecular target therapy against human leukemia. *Expert Opin Ther Targets* 13: 1399–1410, 2009
- Kelly WK, Marks PA: Drug insight: Histone deacetylase inhibitors—development of the new targeted anticancer agent suberoylanilide hydroxamic acid. *Nat Clin Pract Oncol* 2: 150–157, 2005
- Bui LC, Tomkiewicz C, Chevallier A, Pierre S, Bats AS, Mota S, Raingeaud J, Pierre J, Diry M, Transy C, Garlatti M, Barouki R, Coumoul X: Nedd9/Hef1/Cas-L mediates the effects of environmental pollutants on cell migration and plasticity. *Oncogene* 28: 3642–3651, 2009
- Kim SH, Xia D, Kim SW, Holla V, Menter DG, Dubois RN: Human enhancer of filamentation 1 is a mediator of hypoxia-inducible factor-1alpha-mediated migration in colorectal carcinoma cells. *Cancer Res* 70: 4054–4063, 2010

Supplemental information for this article is available online at <http://www.jasn.org/>.