

Long-Term Cancer Risk of Immunosuppressive Regimens after Kidney Transplantation

Martin P. Gallagher,* Patrick J. Kelly,[†] Meg Jardine,* Vlado Perkovic,* Alan Cass,* Jonathan C. Craig,[†] Josette Eris,[‡] and Angela C. Webster^{†§}

*Renal Division, George Institute for International Health, Camperdown, New South Wales, Australia; [†]School of Public Health, University of Sydney, Sydney, New South Wales, Australia; [‡]Statewide Renal Services, Royal Prince Alfred Hospital, Camperdown, New South Wales, Australia; and [§]Australia and New Zealand Dialysis and Transplant Registry, Adelaide, South Australia, Australia

ABSTRACT

Cancer is a widely recognized complication of transplantation, and the effects of various immunosuppressive drugs on cancer risk remains controversial. This randomized trial allocated 489 recipients of first cadaveric renal transplants to one of three groups: Azathioprine and prednisolone, cyclosporine monotherapy, or cyclosporine monotherapy followed by a switch to azathioprine and prednisolone after 3 months. Here, we report cancer outcomes by non-skin cancer (including melanoma) and skin cancer (excluding melanoma) for 481 patients during a median follow-up of 20.6 years. A total of 226 patients developed at least one cancer: 95 with non-skin cancer and 171 with skin cancer. In the intention-to-treat analysis, mean times to first non-skin cancer (16.0, 15.3, and 15.7 years for groups 1 through 3, respectively) and first skin cancer (13.6, 14.3, and 15.2 years, respectively) were not different among the three groups or between any subgroup. In multivariate analyses, non-skin cancer associated with increasing age and previous smoking history, whereas skin cancer associated with increasing age, nonbrown eye color, fairer skin, and a functioning transplant. Treatment allocation did not associate with development of either form of cancer in multivariate analyses. In conclusion, these immunosuppressive regimens, widely used in recent decades, carry similar risks for carcinogenicity after kidney transplantation.

J Am Soc Nephrol 21: 852–858, 2010. doi: 10.1681/ASN.2009101043

Kidney transplantation is the preferred treatment for end-stage kidney disease because of improved patient survival and quality of life and lower treatment costs compared with dialysis. Cancer is a widely recognized complication of transplantation^{1,2} and is likely to become more common among transplant recipients as donation criteria are extended to allow older donors, the age of waiting list patients increases, and transplant recipients live longer.

The relative carcinogenicity of the specific immunosuppressive agents or combinations of agents is not well understood. One reason for this is that randomized trials of various immunosuppressive regimens have not reported cancer outcomes or followed kidney transplant recipients long enough for cancers to develop. Second, al-

though observational data from registries have made an important contribution to our knowledge,^{1–3} their validity in answering drug intervention questions is limited because of the unequal

Received October 12, 2009. Accepted February 16, 2010.

Published online ahead of print. Publication date available at www.jasn.org.

The interpretation and reporting of the data are the responsibility of the authors and in no way should be seen as an official policy or interpretation of the Australia and New Zealand Dialysis and Transplant Registry.

Correspondence: Dr. Martin P. Gallagher, George Institute for International Health, P.O. Box M201 Missenden Road, Camperdown, NSW 2050, Australia. Phone: +61-2-9993-4552; Fax: +61-2-9767-6254; E-mail: mgallagher@george.org.au

Copyright © 2010 by the American Society of Nephrology

distribution of confounders (both known and unknown). These factors combine to limit our knowledge and impair clinicians' attempts to mitigate patients' cancer risk.

The Australian Multicenter Trial of Cyclosporine Withdrawal randomly assigned patients to three arms: Standard therapy of azathioprine and prednisolone (AP), long-term cyclosporine (CY), and short-term cyclosporine followed by withdrawal and maintenance azathioprine and prednisolone (WDL). We previously reported graft and patient survival outcomes at 15 and 20 years of median follow-up.^{4,5} In this study, we compare incident cancer rates and spectrum of cancers across the three arms using the extremely prolonged follow-up provided by registry linkage, the unbiased distribution of confounders arising from the trial randomization, and extensive baseline data on cancer risk.

RESULTS

A total of 489 patients were randomly allocated to one of the three treatment groups between August 1983 and November 1986. Follow-up was available for 481 (98%) patients; the median follow-up of surviving patients was 20.6 years (interquartile range 20.0 to 21.5 years) after randomization. The baseline characteristics of the three treatment groups, including measured risk factors for non-skin and skin malignancies, were similar (Table 1). Following provision according to the study protocol, more recipients in the AP group received antithymocyte globulin (ATG) than in either of the CY group or the WDL group. A total of 226 (46%) patients developed at least 1 cancer. Ninety-five (19%) patients had at least one non-skin cancer (NSC) and 171 (35%) had at least one skin cancer exclud-

Table 1. Characteristics of trial participants

Characteristic	AP	CY	WDL	Total
No. randomly assigned	158	166	165	489
Male gender (n [%])	88 (56)	98 (59)	93 (56)	
Age at transplantation (years; n [%])				
<35	40 (25)	50 (30)	46 (28)	136
35 to 44	39 (25)	34 (20)	37 (22)	110
45 to 54	38 (24)	42 (25)	57 (35)	137
≥55	41 (26)	40 (24)	24 (15)	105
Years on dialysis (median [range])	1.24 (0.20 to 21.70)	1.16 (0.10 to 11.50)	1.21 (0.10 to 8.60)	1.22 (0.00 to 21.70)
History of ischemic heart disease (n [%])	22 (14)	29 (17)	22 (13)	73
BMI at transplantation (kg/m ² ; [n])				
<25	98 (63)	109 (67)	112 (70)	319
≥25	57 (37)	53 (33)	47 (30)	157
Smoking (ever/never; n)	81/77	96/67	87/76	262/220
Latitude of primary residence (n [%])				
>35°S ^a	27 (18)	26 (16)	30 (19)	83
30 to 35°S	70 (46)	77 (48)	72 (46)	219
<30°S	55 (36)	57 (36)	56 (35)	168
Eye color (brown/not brown; n)	60/98	66/99	49/114	175/311
Skin type (n [%])				
fair and freckles	16 (10)	10 (6)	8 (5)	34
fair, no freckles	27 (17)	27 (16)	24 (15)	78
olive	66 (42)	69 (42)	78 (48)	213
darker skin	49 (31)	59 (36)	54 (33)	162
Previous skin cancer (yes/no; n)	3/155	3/163	1/163	7/481
Cold ischemia time (hours; median [range])	22 (7 to 60)	22 (4 to 30)	23 (8 to 39)	22 (4 to 60)
HLA mismatches (n [%])				
0	2 (1)	2 (1)	1 (1)	5
1 to 2	52 (33)	55 (33)	58 (36)	165
≥3	102 (65)	108 (65)	100 (63)	310
2 DR mismatches	18 (12)	19 (12)	20 (13)	57
PRA (current; n)				
≤20%	132 (89)	138 (87)	130 (85)	400
>20%	16 (11)	21 (13)	23 (15)	60
Received ATG (n [%])	94 (59)	9 (5)	8 (5)	111

^aIncludes those with predominant residential history of NZ, northern and southern Europe and Pacific Islands.

BMI, body mass index; PRA, panel-reactive antibodies.

ing melanoma (SCEM; 60% squamous cell carcinomas and 40% basal cell carcinomas) during the study follow-up (Table 2). Adenocarcinomas made up the largest group of NSCs (37%) followed by SCEMs (20%)

Risk for Cancer by Randomized Group

The mean time to first NSC was not different among treatment groups: AP 16.0 years (95% confidence interval [CI] 15.0 to 17.0 years), CY 15.3 years (95% CI 14.5 to 16.1 years), and WDL 15.7 years (95% CI 15.0 to 16.4 years). The overall cumulative incidence of NSC at 20 years after transplantation was 27% and not different among the three treatment groups (27, 27, and 28%, respectively; *P* = 0.96; Figure 1A). The proportions of adenocarcinomas and squamous cell carcinomas within each treatment group were not different (*P* = 0.44 and *P* = 0.1, respectively).

The mean time to the first SCEM was not different among groups: AP 13.6 years (95% CI 12.4 to 14.8 years), CY 14.3 years (95% CI 13.0 to 15.6 years), and WDL 15.2 years (95% CI 14.0 to 16.4 years). The overall cumulative incidence of skin cancer was 48% at 20 years and was not different among the three treatment groups (50, 48, and 46%; *P* = 0.69; Figure 1B). Seven trial participants had a history of skin cancer before

Table 2. Cancer types across all treatment groups out to 20 years after randomization

Cancer Type	AP	CY	WDL	Total
NSC				
total recipients	29	35	31	95
adenocarcinoma	8	15	12	35
colon	4	4	1	9
prostate	0	3	2	5
kidney	0	2	1	3
breast	0	1	2	3
esophagus/stomach	1	1	1	3
other sites	3	3	2	8
unknown primary	0	1	3	4
squamous cell carcinoma	4	11	4	19
cervix	1	3	0	4
lung	0	3	1	4
head and neck	2	3	3	8
other	1	2	0	3
melanoma	7	1	1	9
lymphoma (all)	2	3	4	9
leukemia	1	0	1	2
transitional cell carcinoma	3	0	2	5
Kaposi	0	1	1	2
multiple myeloma	1	0	0	1
other/unknown	3	4	6	13
Skin cancer				
total recipients	51	65	55	171
squamous cell carcinoma	32	36	34	102
basal cell carcinoma	19	28	21	68
other	0	1	0	1




Figure 1. Cumulative incidence of cancer out to 20 years posttransplant. (A) Cumulative incidence of NSC or melanoma, by treatment allocation, over time since randomization. (B) Cumulative incidence of SCEM, by treatment allocation, over time since randomization.

transplantation, but none went on to develop an additional SCEM after transplantation.

Analyses of NSC incidence by baseline subgroups of gender, age, dialysis duration, smoking history, HLA matching, and panel-reactive antibody levels suggested no differences between any treatment groups (Figure 2A). Similarly, for SCEM, analysis by baseline subgroups of gender, age, dialysis duration, smoking history, HLA mismatching, panel-reactive antibody levels, skin type, and eye color did not suggest any differences between any treatment groups (Figure 2B).

Risk for Cancer by Multivariate Modeling

In the univariate analysis of risk factors, only increasing age and history of smoking were significantly associated with the development of NSC after transplantation. Gender, exposure to ATG, body mass index, history of ischemic heart disease, and graft failure showed no association. After allowing for their independent effects in a multivariate model, age and smoking history continued to be significantly associated with NSC de-




Figure 2. Treatment effects upon cancer risk by potential effect modifiers. (A) Effects of randomized study of immunosuppressive treatments on NSC, stratified by potential effect modifiers. *Mismatch is the number of HLA mismatches between donor and recipient (grouped as shown); PRA is the panel-reactive antibody values for recipients (grouped as shown). (B) Effects of randomized study of immunosuppressive treatments on SCEM, stratified by potential effect modifiers. *Mismatch is the number of HLA mismatches between donor and recipient (grouped as shown); PRA is the panel-reactive antibody values for recipients (grouped as shown).

development, but there was no demonstrable effect of immunosuppressive treatment allocation on NSC risk (Figure 3). Using these variables to define groups at low and high risk for NSC after transplantation, the predicted incidence varies between these groups by approximately six-fold over time (Table 3).

In SCEM, the univariate modeling showed that continued transplant function, increased age at transplantation, nonbrown eye color, and fairer skin type all were significantly associated with SCEM development after transplantation. Receipt of ATG, history of smoking, and allocated immunosuppressive treatment were not. After allowing for their independent effects in the multivariate model, male gender, continued transplant function, increased age at transplantation, nonbrown eyes, and fairer skin




Figure 3. Risk factors for NSA and SCEM by multivariate modeling.

type showed significant association with SCEM (Figure 3). Using these variables to define groups at low and high risk for SCEM after transplantation, the predicted incidence varies from 10- to 28-fold over time (Table 3).

DISCUSSION

These results suggest that patients who were treated with these three immunosuppressive regimens, commonly used in renal transplantation for the past 30 years, show no difference in the development of both NSC and skin cancer out to 20 years after transplantation. We demonstrated no difference in drug effects for any patient subgroups, but the risk for cancer was strongly associated with patient characteristics known at the time of transplantation. As in the general population, age and smoking history are the major determinants of posttransplantation NSC risk. The results also highlight the inexorable rise in skin cancer during time after transplantation and the reduction in risk after graft loss, presumably mediated through a reduced need for subsequent immunosuppression.

Despite the existing knowledge regarding cancer risk, no immunosuppressive treatment strategies have been proved to reduce this risk. Much debate has focused on the effects of the various immunosuppressive agents and dose regimens, with concerns raised about azathioprine,^{4,6,7} cyclosporine,^{8,9} mycophenolate mofetil,⁸ tacrolimus,¹⁰ and triple therapy.¹¹ Our data suggest that azathioprine- and cyclosporine-based regimens are associated with similar overall long-term cancer risks, suggesting that the risk may be mediated by the total burden of immunosuppression more than the agent. It remains possible that small differences in drug effects do exist but that we did not have the statistical power to detect them even after 20 years of follow-up. The clinical implications of such small differences, if they do exist undetected by our analysis, is questionable, however, because most modern transplant regimens use combinations of antiprolifera-

Table 3. Predicted risk for NSC and skin cancer over time after transplantation

Risk Stratum	Group Characteristics	No. (95% CI) with Cancers Predicted per 100 People Treated		
		5 years	10 years	20 years
NSC				
low	<35 years old, never smoked	2.5 (0.9 to 4.1)	4.9 (2.1 to 7.7)	12.0 (5.8 to 18.0)
high	≥55 years old, previous or current smoker	17.0 (9.8 to 24.0)	32.0 (20.0 to 41.0)	61.0 (43.0 to 74.0)
SCEM				
low	Female, <35 years old, brown eyes, dark skin	1.9 (0.7 to 3.1)	4.2 (1.7 to 6.6)	9.3 (4.1 to 14.0)
high	Male, ≥55 years old, nonbrown eyes, fair skin	56.0 (29.0 to 73.0)	84.0 (55.0 to 95.0)	99.0 (82.0 to 100.0)

tive agents and calcineurin antagonists. Our analysis also underlines that other factors contribute much more to cancer risk after transplantation than the specific immunosuppressive therapy and that risk estimation is possible.

One recent publication reported a possible reduction in cancer risk at 24 months of follow-up with sirolimus,¹² but this has not been confirmed by other trials¹³ or a meta-analysis.¹⁴ Apart from the association of anti-CD3 mAbs with posttransplantation lymphoproliferative disease,¹⁵ the clinical evidence for one treatment regimen over another in reducing posttransplantation malignancy is varied and not compelling. This analysis suggests that longer term follow-up of recent comparative immunosuppressive trials (e.g., Efficacy Limiting Toxicity Elimination [ELITE]-Symphony Trial¹³) could provide important data on cancer outcomes and that better markers of the degree of immunosuppression may also aid in reducing cancer risk.

A strength of these data is that they arose from a randomized trial, using a crossed synthesis design made possible using the Australian and New Zealand Transplant Registry, demonstrating that it is feasible and informative to obtain long-term results from trials with almost complete outcome data. The randomized allocation of immunosuppressive treatment ensures that the measure of no difference should be the least biased estimate of effect possible. Nonetheless, there are some limitations to the results of this trial. The trial was designed to test a primary outcome of graft survival, although cancer incidence was a prespecified outcome and important cancer risk factors were collected at baseline. The increase in cancer risk after transplantation has been widely documented, but the reported rates do vary and this is likely a function of regional variations in cancer epidemiology as well as differing study methods. Patients in this study come from a largely white population arising from the broader Australian general population, who have among the highest rates of skin cancer in the world. Thus, the absolute estimates of risk may differ for transplant recipients in different settings, but it is likely that the relative differences will hold true. In addition, some of the immunosuppressive regimens used in the trial, most notably cyclosporine monotherapy, the absence of mandated cyclosporine drug monitoring, and the absence of new antiproliferative agents (e.g., mycophenolate), differ from current immunosuppressive practice, so caution should be exercised when extrapolating the results to contemporary immunosuppressive protocols. It is likely, however, that today's immuno-

suppressive regimens, with their lower acute rejection rates, represent more potent immunosuppression.

Although the large transplant registries have provided an invaluable real-world insight into posttransplantation malignancy,^{1,2,16} their value in clarifying the effects of individual drugs and drug regimens is limited by selection bias as a result of the absence of random allocation to treatments and limited recording of drug exposures. The major burden of cancer after transplantation is seen more than 5 years after engraftment, beyond the scope and affordability of routine clinical trial follow-up. Cancer event rates seen in our study were approximately 1.25% per annum for NSC and 2.5% per annum for SCEM, making it difficult and expensive to power a study using cancers as an outcome. The rates of cancer seen in other studies have been higher¹⁷ but still low in absolute terms. In such a context, deriving long-term follow-up of large randomized trials from patient registries through similar cross-synthesis designs offers the best hope of identifying treatments to reduce the cancer burden in transplant recipients.

CONCISE METHODS

Details of the trial, including the extent of deviations from protocol therapy, have been previously reported^{4,5,18} but will be summarized in brief here.

Participants

Eligible participants were adult patients who were receiving primary, deceased-donor renal transplants. Exclusion criteria were insulin-dependent diabetes, abnormal liver function tests, previous malignancy, malabsorption, active infection, or a known contraindication to azathioprine. Patients were randomly assigned at the time of transplantation between 1983 and 1986 in seven centers throughout Australia. The randomization sequence was centrally generated by computer and stratified by center, with patient assignment by sequentially labeled, opaque, sealed envelopes delivered to each center. The study was open-label; patients, treating physicians, data entry staff, and data analysts were not blinded to treatment allocation.

Randomized Interventions

AP Group.

Participants who were allocated to AP received treatment with azathioprine and prednisolone with the option of induction ATG ther-

apy. Oral azathioprine (3 mg/kg) was given preoperatively followed by 2 mg/kg per d. Intravenous methylprednisolone was given preoperatively (1000 mg) and on day 1 (500 mg) followed by oral prednisolone daily tapering to maintenance doses of 10 to 15 mg/d. Two of the seven centers involved in the trial opted for routine use of prophylactic ATG (5 mg/kg) for all of their recipients, dosed every other day for a total of seven doses.

CY Group.

Participants who were allocated to CY received long-term cyclosporine monotherapy. Intravenous cyclosporine (5 mg/kg) was given preoperatively and on day 1 (4 mg/kg) followed by oral cyclosporine 12.5 mg/kg tapering to 7.5 mg/kg by 3 months after transplantation. Intravenous prednisolone was given at operation (1000 mg) and on day 1 (500 mg). Patients with clinical signs of nephrotoxicity had their cyclosporine dosage reduced to 7.5 mg/kg, with therapy changed to azathioprine and prednisolone when there was no improvement within 1 week. Patients who were receiving cyclosporine could receive additional maintenance prednisolone if they experienced three or more episodes of rejection.

WDL Group.

Participants who were allocated to WDL received identical treatment as CY for the first 3 months after transplantation. When at 3 months there was no evidence of rejection (either clinically or on renal biopsy) in the previous 2 weeks, treatment was changed to azathioprine (2 mg/kg per d) and prednisolone (20 mg/d). Therapy transition was completed after a 4-day overlap of the two strategies for the majority of recipients.

Outcomes

The primary and secondary outcomes of the study—death-censored graft survival, patient survival, and the combined end point of death or graft loss^{4,5,18}—have previously been reported. Cancer outcomes were prespecified in the original trial and were briefly reported at 15 and 20 years^{4,5} but were not previously analyzed and reported in detail. Cancer outcomes were measured by linking the trial database with the Australia and New Zealand Dialysis and Transplant Registry database, using data up to and including the census of December 2005.

Statistical Analysis

An intention-to-treat analysis of cancer outcomes by randomized treatment allocation was assessed using cumulative incidence curves and Cox proportional hazard models. Pair-wise differences between treatment groups were conducted both overall and by patient subgroups. In all analyses, recipients were censored from the analysis at their last known date of follow-up (alive, lost to follow-up, or death) when they did not experience the cancer event of interest.

A secondary analysis was conducted to examine the risk factors for NSC using a Cox proportional hazards model and similarly for SCEM. Age, gender, allocated treatment group, and graft status (functioning *versus* failed) were included in the multivariate models, regardless of their statistical significance. Graft status was a time-varying variable. Other variables that were considered possible risk factors

for cancer were included in the multivariable modeling at their univariate $P \leq 0.3$ and remained in the final model at the multivariable $P \leq 0.05$. Potential effect modification was investigated by testing for interaction between age and gender and also between graft status and all other variables remaining in the model. For both analyses, no interaction term had a $P < 0.05$; hence, no interaction terms were included in either final model. The assumption of proportion hazards was checked for each variable by including a time-varying covariate of the variable of interest, multiplied by $\log(\text{time})$. Schoenfeld residuals were also plotted to check potential deviation from the proportional hazards assumption. Statistical analyses was conducted in SAS 9.1.

ACKNOWLEDGMENTS

This study was supported by Sandoz out to 10 years of follow-up, but this analysis received no dedicated support. Authors have received support from the Australian National Heart Foundation Postdoctoral Fellowship (V.P.) and Australian National Health and Medical Research Council Fellowship (A.C.). The Australia and New Zealand Dialysis and Transplant Registry is funded by the Australian Department of Health and Ageing, the NZ Ministry of Health, and Kidney Health Australia.

Parts of the data reported here were supplied by the Australia and New Zealand Dialysis and Transplant Registry.

We thank those who instigated the original study, especially Bruce Hall, David Tiller, Ian Hardie, John Mahony, Tim Matthew, Geoffrey Thatcher, Peter Miach, Nip Thompson, and Ross Sheil, along with all of the clinicians at all of the centers participating in the trial, and Royal Prince Alfred Hospital (Sydney), Princess Alexandra Hospital (Brisbane), Queen Elizabeth Hospital (Adelaide), Royal North Shore Hospital (Sydney), Monash Medical Centre (Melbourne), Austin Hospital (Melbourne), and Sir Charles Gairdner Hospital (Perth).

DISCLOSURES

M.G. and V.P. have received speaking fees from pharmaceutical companies that manufacture immunosuppressive agents used in transplantation; J.E. has received travel assistance from and chairs advisory boards of pharmaceutical companies that manufacture immunosuppressive agents; and the Australia and New Zealand Dialysis and Transplant Registry has received contributions in the past from pharmaceutical companies that manufacture immunosuppressive agents.

REFERENCES

1. Vajdic CM, McDonald SP, McCredie MR, van Leeuwen MT, Stewart JH, Law M, Chapman JR, Webster AC, Kaldor JM, Grulich AE: Cancer incidence before and after kidney transplantation. *JAMA* 296: 2823–2831, 2006
2. Birkeland SA, Storm HH, Lamm LU, Barlow L, Blohme I, Forsberg B, Eklund B, Fjeldborg O, Friedberg M, Frödin L, *et al.*: Cancer risk after renal transplantation in the Nordic countries, 1964–1986. *Int J Cancer* 60: 183–189, 1995
3. Webster AC, Craig JC, Simpson JM, Jones MP, Chapman JR: Identifying high risk groups and quantifying absolute risk of cancer after

- kidney transplantation: A cohort study of 15,183 recipients. *Am J Transplant* 7: 2140–2151, 2007
- Gallagher M, Jardine M, Perkovic V, Cass A, McDonald S, Petrie J, Eris J: Cyclosporine withdrawal improves long-term graft survival in renal transplantation. *Transplantation* 87: 1877–1883, 2009
 - Gallagher MP, Hall B, Craig J, Berry G, Tiller DJ, Eris J: A randomized controlled trial of cyclosporine withdrawal in renal-transplant recipients: 15-Year results. *Transplantation* 78: 1653–1660, 2004
 - Scharf J, Nahir M, Eidelman S, Jacobs R, Levin D: Carcinoma of the bladder with azathioprine therapy. *JAMA* 237: 152, 1977
 - Krutchik AN, Buzdar AU, Tashima CK: Azathioprine and breast carcinoma. *JAMA* 239: 107–108, 1978
 - Guba M, Graeb C, Jauch KW, Geissler EK: Pro- and anti-cancer effects of immunosuppressive agents used in organ transplantation. *Transplantation* 77: 1777–1782, 2004
 - Dantal J, Hourmant M, Cantarovich D, Giral M, Blancho G, Dreno B, Souillou JP: Effect of long-term immunosuppression in kidney-graft recipients on cancer incidence: Randomised comparison of two cyclosporin regimens. *Lancet* 351: 623–628, 1998
 - Imao T, Ichimaru N, Takahara S, Kokado Y, Okumi M, Imamura R, Namba Y, Isaka Y, Nonomura N, Okuyama A: Risk factors for malignancy in Japanese renal transplant recipients. *Cancer* 109: 2109–2115, 2007
 - Penn I, First MR: Development and incidence of cancer following cyclosporine therapy. *Transplant Proc* 18: 210–215, 1986
 - Schena FP, Pascoe MD, Alberu J, del Carmen Rial M, Oberbauer R, Brennan DC, Campistol JM, Racusen L, Polinsky MS, Goldberg-Alberts R, Li H, Scarola J, Neylan JF: Conversion from calcineurin inhibitors to sirolimus maintenance therapy in renal allograft recipients: 24-Month efficacy and safety results from the CONVERT trial. *Transplantation* 87: 233–242, 2009
 - Ekberg H, Bernasconi C, Tedesco-Silva H, Vitko S, Hugo C, Demirbas A, Acevedo RR, Grinyo J, Frei U, Vanrenterghem Y, Daloze P, Halloran P: Calcineurin inhibitor minimization in the SYMPHONY study: Observational results 3 years after transplantation. *Am J Transplant* 9: 1876–1885, 2009
 - Webster AC, Lee VW, Chapman JR, Craig JC: Target of rapamycin inhibitors (TOR-I; sirolimus and everolimus) for primary immunosuppression in kidney transplant recipients. *Cochrane Database Syst Rev* CD004290, 2006
 - Cherikh WS, Kauffman HM, McBride MA, Maghirang J, Swinnen LJ, Hanto DW: Association of the type of induction immunosuppression with posttransplant lymphoproliferative disorder, graft survival, and patient survival after primary kidney transplantation. *Transplantation* 76: 1289–1293, 2003
 - Robson R, Cecka JM, Opelz G, Budde M, Sacks S: Prospective registry-based observational cohort study of the long-term risk of malignancies in renal transplant patients treated with mycophenolate mofetil. *Am J Transplant* 5: 2954–2960, 2005
 - Kasiske BL, Snyder JJ, Gilbertson DT, Wang C: Cancer after kidney transplantation in the United States. *Am J Transplant* 4: 905–913, 2004
 - Hall BM, Tiller DJ, Hardie I, Mahony J, Mathew T, Thatcher G, Miach P, Thomson N, Sheil AG: Comparison of three immunosuppressive regimens in cadaver renal transplantation: Long-term cyclosporine, short-term cyclosporine followed by azathioprine and prednisolone, and azathioprine and prednisolone without cyclosporine. *N Engl J Med* 318: 1499–1507, 1988

EMBARGOED FOR RELEASE until April 29, 2010 – 5:00 PM (EDT)

ASN Contact: Shari Leventhal • 202-416-0658 (p) • sleventhal@asn-online.org

CANCER RISK THE SAME FOR KIDNEY TRANSPLANT RECIPIENTS, NO MATTER THE DRUG

Washington, DC (April 23, 2010) — Drugs taken by kidney transplant recipients to prevent organ rejection carry similar risks of cancer, according to a study appearing in an upcoming issue of the *Journal of the American Society Nephrology* (JASN). The results suggest that no single medication is to blame for the higher incidence of cancer seen among patients who have undergone transplantation.

Individuals who receive a kidney transplant have an increased risk of developing cancer compared with the general population. Researchers suspected that the increased risk may come from immunosuppressive medications that patients must take long-term to prevent organ rejection. To investigate, Martin Gallagher, MBBS, FRACP (The George Institute for International Health in Australia) and his colleagues studied the incidence of cancer in transplant patients who took part in a randomized clinical trial 20 years ago, looking for any differences in cancer risk associated with different immunosuppressive drugs.

The researchers studied the incidence of cancer among 481 kidney transplant recipients in the Australian Multicentre Trial of Cyclosporine Withdrawal who each received one of three treatment regimens: azathioprine and prednisolone, cyclosporine monotherapy, or cyclosporine monotherapy followed by a switch to azathioprine and prednisolone after three months.

A total of 226 patients in the trial developed at least one cancer. By 20 years post transplant, 27% of patients developed non-skin cancer and 48% of patients developed skin cancer. One type of treatment did not have a greater effect on cancer timing or incidence than another, indicating that the therapies carry similar risks for cancer after kidney transplantation.

“We have shown no significant differences with a high degree of precision, allowing us to conclude that any differences in cancer risk from these different treatments are unlikely to be clinically significant,” said Dr. Gallagher. He added that this study provides the strongest evidence yet that no single immunosuppressive medication appears to drive the increase in cancer risk seen after transplantation.

Press Release

The study also indicates that certain patient characteristics that are known at the time of transplantation have a significant effect on recipients' increased risk of cancer. (Non-skin cancer was associated with increasing age and previous smoking history; skin cancer was associated with increasing age, non-brown eye color, fairer skin, and a functioning transplant.) Therefore, patients at especially high risk can be monitored more closely and use preventive measures to protect against cancer.

The authors noted that immune suppressive treatments have evolved since the trial was designed 20 years ago. It is likely that today's immunosuppressive regimens, which are better at preventing acute rejection, are more potent at immunosuppression.

Study co-authors include Meg Jardine, MBBS, PhD, FRACP, Vlado Perkovic, MBBS, PhD, FRACP, Alan Cass, MBBS, PhD, FRACP (The George Institute for International Health); Patrick Kelly, PhD, Jonathan Craig, MBBS, PhD (University of Sydney, in Australia); Josette Eris, MBBS, PhD, FRACP (Royal Prince Alfred Hospital, in Camperdown, Australia); and Angela Webster, MBBS, PhD, MRCP (University of Sydney and Australia and New Zealand Dialysis and Transplant (ANZDATA) Registry, in Adelaide, South Australia).

Disclosures: Dr. Gallagher and Dr. Perkovic have received speaking fees from Roche Pharmaceuticals. Dr. Eris has received travel assistance from and chairs advisory boards of Roche (chair), Novartis, Wyeth, and Janssen-Cilag; she has also received travel support from Novartis and Roche within the last three years. The ANZDATA Registry has received contributions from Roche, Wyeth and Novartis.

The article, entitled "Long-Term Cancer Risk of Immunosuppressive Regimens after Kidney Transplantation," will appear online at <http://jasn.asnjournals.org/> on April 29, 2010, doi 10.1681/ASN.2009101043.

The content of this article does not reflect the views or opinions of The American Society of Nephrology (ASN). Responsibility for the information and views expressed therein lies entirely with the author(s). ASN does not offer medical advice. All content in ASN publications is for informational purposes only, and is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. This content should not be used during a medical emergency or for the diagnosis or treatment of any medical condition. Please consult your doctor or other qualified health care provider if you have any questions about a medical condition, or before taking any drug, changing your diet or commencing or discontinuing any course of treatment. Do not ignore or delay obtaining professional medical advice because of information accessed through ASN. Call 911 or your doctor for all medical emergencies.

Founded in 1966, the American Society of Nephrology (ASN) is the world's largest professional society devoted to the study of kidney disease. Comprised of 11,000 physicians and scientists, ASN continues to promote expert patient care, to advance medical research, and to educate the renal community. ASN also informs policymakers about issues of importance to kidney doctors and their patients. ASN funds research, and through its world-renowned meetings and first-class publications, disseminates information and educational tools that empower physicians.

###