

patients with CKD, trading greater weight loss for increased risk may be the best option. Weight loss after bariatric surgery is associated with a number of improvements that can help resolve kidney disease—even ESRD, eliminating the need for transplantation.¹⁴

Bariatric surgery improves underlying causes of CKD (dyslipidemia, hypertension, urinary albumin excretion, atherosclerosis, and metabolic syndrome). It can also improve measures linked to kidney function (insulin resistance, adiponectin, TNF- α , IL-18, and C-reactive protein).^{16,17} For patients attempting to meet kidney transplant requirements through surgical weight loss, RYGB may provide the quickest and best weight loss outcomes. It is the most commonly performed procedure in patients with obesity and CKD; it enables them to undergo transplantation.¹⁸

Bariatric surgery in CKD patients requires a careful balance of risks and benefits.^{19,20} High-quality controlled trials are needed to identify and quantify those risks and benefits. To improve options for predicting surgical risk on a case-by-case basis in patients with obesity and diabetes, studies of GFR estimation techniques are necessary.

DISCLOSURES

None.

REFERENCES

- Turgeon NA, Perez S, Mondestin M, Davis SS, Lin E, Tata S, Kirk AD, Larsen CP, Pearson TC, Sweeney JF: The impact of renal function on outcomes of bariatric surgery. *J Am Soc Nephrol* 23: 885–894, 2012
- Poirier P, Cornier MA, Mazzone T, Stiles S, Cummings S, Klein S, McCullough PA, Ren Fielding C, Franklin BA; American Heart Association Obesity Committee of the Council on Nutrition, Physical Activity, and Metabolism: Bariatric surgery and cardiovascular risk factors: A scientific statement from the American Heart Association. *Circulation* 123: 1683–1701, 2011
- Buchwald H, Estok R, Fahrbach K, Banel D, Jensen MD, Pories WJ, Bantle JP, Sledge I: Weight and type 2 diabetes after bariatric surgery: Systematic review and meta-analysis. *Am J Med* 122: 248–256, 2009
- Delanghe JR, Cobbaert C, Harmoinen A, Jansen R, Laitinen P, Panteghini M: Focusing on the clinical impact of standardization of creatinine measurements: A report by the EFCC Working Group on Creatinine Standardization. *Clin Chem Lab Med* 49: 977–982, 2011
- Nair S, Mishra V, Hayden K, Lisboa PJ, Pandya B, Vinjamuri S, Hardy KJ, Wilding JP: The four-variable modification of diet in renal disease formula underestimates glomerular filtration rate in obese type 2 diabetic individuals with chronic kidney disease. *Diabetologia* 54: 1304–1307, 2011
- Praditpomsilpa K, Townamchai N, Chaiwatanarat T, Tiranathanagul K, Katawatin P, Susantitaphong P, Trakarnvanich T, Kanjanabuch T, Avihingsanon Y, Tungsanga K, Eiam-Ong S: The need for robust validation for MDRD-based glomerular filtration rate estimation in various CKD populations. *Nephrol Dial Transplant* 26: 2780–2785, 2011
- Michels WM, Grootendorst DC, Verduijn M, Elliott EG, Dekker FW, Krediet RT: Performance of the Cockcroft-Gault, MDRD, and new CKD-EPI formulas in relation to GFR, age, and body size. *Clin J Am Soc Nephrol* 5: 1003–1009, 2010
- Oh SJ, Lee JI, Ha WC, Jeong SH, Yim HW, Son HS, Sohn TS: Comparison of cystatin C- and creatinine-based estimation of glomerular filtration rate according to glycaemic status in type 2 diabetes [published online ahead of print March 13, 2012]. *Diabet Med* doi:10.1111/j.1464-5491.2012.03628.x
- Jesudason DR, Clifton P: Interpreting different measures of glomerular filtration rate in obesity and weight loss: Pitfalls for the clinician [published online ahead of print December 20, 2011]. *Int J Obes (Lond)* doi:10.1038/ijo.2011.242
- Matlaga BR, Shore AD, Magnuson T, Clark JM, Johns R, Makary MA: Effect of gastric bypass surgery on kidney stone disease. *J Urol* 181: 2573–2577, 2009
- Nasr SH, D'Agati VD, Said SM, Stokes MB, Langoza MV, Radhakrishnan J, Markowitz GS: Oxalate nephropathy complicating Roux-en-Y Gastric Bypass: An underrecognized cause of irreversible renal failure. *Clin J Am Soc Nephrol* 3: 1676–1683, 2008
- Penniston KL, Kaplon DM, Gould JC, Nakada SY: Gastric band placement for obesity is not associated with increased urinary risk of urolithiasis compared to bypass. *J Urol* 182: 2340–2346, 2009
- Semins MJ, Asplin JR, Steele K, Assimos DG, Lingeman JE, Donahue S, Magnuson T, Schweitzer M, Matlaga BR: The effect of restrictive bariatric surgery on urinary stone risk factors. *Urology* 76: 826–829, 2010
- Currie A, Chetwood A, Ahmed AR: Bariatric surgery and renal function. *Obes Surg* 21: 528–539, 2011
- Buchwald H, Avidor Y, Braunwald E, Jensen MD, Pories W, Fahrbach K, Schoelles K: Bariatric surgery: A systematic review and meta-analysis. *JAMA* 292: 1724–1737, 2004
- Ahmed MH, Byrne CD: Bariatric surgery and renal function: A precarious balance between benefit and harm. *Nephrol Dial Transplant* 25: 3142–3147, 2010
- Chauhan V, Vaid M, Gupta M, Kalanuria A, Parashar A: Metabolic, renal, and nutritional consequences of bariatric surgery: Implications for the clinician. *South Med J* 103: 775–783, 2010
- Modanlou KA, Muthyala U, Xiao H, Schnitzler MA, Salvalaggio PR, Brennan DC, Abbott KC, Graff RJ, Lentine KL: Bariatric surgery among kidney transplant candidates and recipients: Analysis of the United States renal data system and literature review. *Transplantation* 87: 1167–1173, 2009
- Gore JL: Obesity and renal transplantation: Is bariatric surgery the answer? *Transplantation* 87: 1115, 2009
- Williams ME, Magerowski GA, Blackburn GL: Nutritional management of obesity and the metabolic syndrome in the patient with chronic kidney disease. In: *Nutritional Management of Renal Disease*, 3rd Ed., edited by Kopple JD, Massry JD, Kalatar Zadeh K, Amsterdam, Academic Press, In press.

See related article, "The Impact of Renal Function on Outcomes of Bariatric Surgery," on pages 885–894.

New Observational Data Demonstrate that Mortality Is Lower in Patients Receiving More Frequent Dialysis

David C. Wheeler and Ben Caplin

Centre for Nephrology, University College London Medical School, London, United Kingdom

J Am Soc Nephrol 23: 770–773, 2012.
doi: 10.1681/ASN.2012030311

Published online ahead of print. Publication date available at www.jasn.org.

Correspondence: Dr. David C. Wheeler, Centre for Nephrology, University College London Medical School, Royal Free Campus, Rowland Hill Street, London NW3 2PF, United Kingdom. E-mail: d.wheeler@ucl.ac.uk

Copyright © 2012 by the American Society of Nephrology

Despite improvements in medical care and technology, the life expectancy of patients with ESRD receiving hemodialysis in the United States remains high, with an annual mortality rate of around 20%/yr.¹ Conventional dialysis regimens based on three times weekly treatment sessions are reasonably acceptable to patients and are cost effective, but may not deliver optimum therapy. The obvious question is whether delivery of more dialysis improves survival among these patients.² Observational studies dating back to the 1990s predicted this may be the case, with better survival reported among patients on a three times weekly regimen who received a larger delivered dose, as assessed by solute clearance.^{3,4}

The first randomized controlled trial designed to test the hypothesis that delivering more dialysis would improve outcomes, the Hemodialysis (HEMO) study, was negative.⁵ In this study, a modest increase in delivered dialysis dose, which was assessed in terms of solute clearance as measured by KT/V and achieved in the context of a three times weekly regimen, showed no clear benefits in terms of patient survival.

Increasing the frequency of dialysis sessions will enhance solute clearance and salt and water removal. Observational studies again indicate this strategy is associated with better outcomes.^{6,7} Three recent randomized controlled trials have tested the hypothesis that delivering more frequent treatment improves outcomes in hemodialysis patients. Using standard thrice weekly dialysis as the comparator, these trials have assessed short daily regimens⁸ or long nocturnal regimens delivered six times weekly.^{9,10} The trials were not powered to determine effects on mortality, although mortality was part of a composite end point in two studies,^{8,9} but did show benefits on surrogate end points including BP,⁸⁻¹⁰ hyperphosphatemia,⁸⁻¹⁰ and left ventricular hypertrophy,^{8,9} as well as improvements in physical health scores.⁸ The frequency of vascular access-related events increased in more frequently dialyzed patients in two of the studies,^{8,9} although there was no difference in access survival.

In these trials, random allocation to the two interventions (standard versus more frequent dialysis regimens) reduces the chances of there being any systematic differences between the two groups, making the controls an appropriate substitute for the treated group in all ways except for the intervention of interest. However, such randomized trials not only require substantial resources, but are also difficult to conduct, particularly when one or both of the interventions require major changes to patient lifestyle. From a patient's perspective, allowing the play of chance to determine allocation to a treatment regimen that will have major lifestyle implications may not be appealing and may limit willingness to enroll. It is therefore not surprising that it has been difficult to conduct trials of different dialysis regimens powered to assess hard clinical end points.

Accepting these difficulties, what more can we learn from the observational data? In this issue of *JASN*, Weinhandl *et al.*¹¹ present a large observational study in which they compare outcomes in 1873 home hemodialysis patients receiving daily treatment to outcomes in 9365 in-center controls receiving a standard

thrice weekly regimen. The authors use a complex algorithm, which includes up to 17 different patient factors, in an attempt to ensure that the two groups were as precisely matched as possible. In an intention-to-treat analysis, they demonstrate a lower hazard ratio for all-cause mortality (0.87; 95% confidence interval, 0.78–0.97) among those patients receiving home dialysis. The authors reasonably conclude that daily dialysis, delivered at home, is associated with a modest survival benefit.

The paper provides the largest and most robust observational data to date addressing this question. However, as the authors concede, the study has limitations. A significant proportion of patients discontinued home dialysis, suggesting that this therapy was not acceptable in the longer term. All home patients, and presumably none of the in-center subjects, used a specific machine, the Nx stage system. Thus, the improvement in outcome could have resulted from a technical difference in the equipment rather than the frequency of dialysis. In addition, as the authors discussed, because no data on compliance are available, they can only assume that those patients assigned to a daily home regimen actually received more dialysis. Furthermore, as in any retrospective study, potential bias in data collection is an important consideration. For example, the completeness of comorbidity data may systematically differ between the two groups.

The key question in interpreting this study, however, is whether the control group receiving in-center treatment represents an appropriate comparator population for the home dialysis group. In other words, were the two groups comparable in all ways other than exposure to the different dialysis regimens? The authors identified five control subjects for each home dialysis case, but even with data from a quarter of a million dialysis patients available from the US Renal Data Systems data set, only about one third of cases were matched on all factors and almost 9% were matched on less than half. This means that despite matching, the home dialysis group was not only a year younger but also less deprived socioeconomically, as demonstrated by the lower number dually eligible for both Medicare and Medicaid. Although the authors report that adjustment for matching factors did not alter the relative hazard ratio for death, these observations highlight the challenge of identifying appropriate controls in this type of study.

As acknowledged by the authors and in common with all observational studies, a further limitation is the extent of unmeasured confounding. No statistical method can deal with the possibility that unquantified factors associated with both the exposure (home hemodialysis) and the outcome (death) account for the differences in mortality observed between the two groups. In an effort to address this possibility, the authors attempt to quantify the magnitude of effect at various levels of differential prevalence that such an unmeasured factor would need to exert to account for the differences in outcome. Although the authors should be commended for their attempts to illustrate what is effectively a shortcoming in all observational studies, the

situation may be more complex because several small confounders, particularly when combined with measurement error, can substantially alter the estimate of effects due to the exposure.¹²

The hazard ratio in the study by Weinhandl *et al.* is closer to unity than that reported in previous observational studies, as referenced in the paper, which had less rigorous control for confounding factors. Therefore, do we now have an accurate estimate of the beneficial effect of daily home versus in-center thrice weekly dialysis or would further adjustment for multiple small causes of confounding shift the hazard ratio even closer to 1? Can we ever assume that estimates of effects from observational studies would be reproduced in prospective randomized controlled trials? We should remember that there are well known examples in the literature where this has not been the case.¹³ Such disparity has been attributed to a failure to adequately adjust for multiple (time dependent, socioeconomic) confounders in the observational studies.¹³ Indeed it has been suggested that “. . . useful observational studies on the effects of therapy will be the exception. . .”¹⁴ as the selection of intervention by patient characteristics will play such an overwhelming part in determining exposure and it is almost impossible to control for. In reality, systematic comparison of observational versus randomized controlled evidence has not identified large differences in the effect estimates reached using the two techniques.^{15–17} However, there have been no comparisons of observational and randomized evidence in the context of dialysis. Of all the clinical scenarios in which confounding may influence the interpretation of evidence collected in the context of observational versus randomized trials, those involving invasive procedures or lifestyle disruption might be most prone.¹⁶

We should perhaps accept that we will never get a randomized controlled trial comparing daily home hemodialysis versus thrice weekly center-based treatment powered to assess differences in mortality. The observational study by Weinhandl *et al.* is informative and provides the best data to date on the subject, despite the above limitations. Although meta-analysis of independent randomized trials will remain the gold standard for assessing the effectiveness of dialysis interventions, the accompanying study is consistent with the hypothesis that daily dialysis therapy is of benefit to patients. Unless data from clinical trials become available in the future, decisions regarding the provision of daily dialysis delivered at home are likely to be based on these observational data.

DISCLOSURES

D.C.W. has received honoraria from Fresenius.

REFERENCES

1. US Renal Data Systems: Annual report 2011. Available at: www.usrds.org/adr.aspx. Accessed March 11, 2012
2. Gorodetskaya I, Zenios S, McCulloch CE, Bostrom A, Hsu CY, Bindman AB, Go AS, Chertow GM: Health-related quality of life and estimates of utility in chronic kidney disease. *Kidney Int* 68: 2801–2808, 2005
3. Owen WF Jr, Lew NL, Liu Y, Lowrie EG, Lazarus JM: The urea reduction ratio and serum albumin concentration as predictors of mortality in patients undergoing hemodialysis. *N Engl J Med* 329: 1001–1006, 1993
4. McClellan WM, Soucie JM, Flanders WD: Mortality in end-stage renal disease is associated with facility-to-facility differences in adequacy of hemodialysis. *J Am Soc Nephrol* 9: 1940–1947, 1998
5. Eknoyan G, Beck GJ, Cheung AK, Daugirdas JT, Greene T, Kusek JW, Allon M, Bailey J, Delmez JA, Depner TA, Dwyer JT, Levey AS, Levin NW, Milford E, Ornt DB, Rocco MV, Schulman G, Schwab SJ, Teehan BP, Toto R; Hemodialysis (HEMO) Study Group: Effect of dialysis dose and membrane flux in maintenance hemodialysis. *N Engl J Med* 347: 2010–2019, 2002
6. Pauly RP, Gill JS, Rose CL, Asad RA, Chery A, Pierratos A, Chan CT: Survival among nocturnal home haemodialysis patients compared to kidney transplant recipients. *Nephrol Dial Transplant* 24: 2915–2919, 2009
7. Johansen KL, Zhang R, Huang Y, Chen SC, Blagg CR, Goldfarb-Rumyantsev AS, Hoy CD, Lockridge RS Jr, Miller BW, Eggers PW, Kutner NG: Survival and hospitalization among patients using nocturnal and short daily compared to conventional hemodialysis: A USRDS study. *Kidney Int* 76: 984–990, 2009
8. Chertow GM, Levin NW, Beck GJ, Depner TA, Eggers PW, Gassman JJ, Gorodetskaya I, Greene T, James S, Larive B, Lindsay RM, Mehta RL, Miller B, Ornt DB, Rajagopalan S, Rastogi A, Rocco MV, Schiller B, Sergeeva O, Schulman G, Ting GO, Unruh ML, Star RA, Kliger AS; FHN Trial Group: In-center hemodialysis six times per week versus three times per week. *N Engl J Med* 363: 2287–2300, 2010
9. Rocco MV, Lockridge RS Jr, Beck GJ, Eggers PW, Gassman JJ, Greene T, Larive B, Chan CT, Chertow GM, Copland M, Hoy CD, Lindsay RM, Levin NW, Ornt DB, Pierratos A, Pipkin MF, Rajagopalan S, Stokes JB, Unruh ML, Star RA, Kliger AS, Kliger A, Eggers P, Briggs J, Hostetter T, Narva A, Star R, Augustine B, Mohr P, Beck G, Fu Z, Gassman J, Greene T, Daugirdas J, Hunsicker L, Larive B, Li M, Mackrell J, Wiggins K, Sherer S, Weiss B, Rajagopalan S, Sanz J, Dellagrottaglie S, Kariisa M, Tran T, West J, Unruh M, Keene R, Schlarb J, Chan C, McGrath-Chong M, Frome R, Higgins H, Ke S, Mandaci O, Owens C, Snell C, Eknoyan G, Appel L, Cheung A, Derse A, Kramer C, Geller N, Grimm R, Henderson L, Prichard S, Roecker E, Rocco M, Miller B, Riley J, Schuessler R, Lockridge R, Pipkin M, Peterson C, Hoy C, Fensterer A, Steigerwald D, Stokes J, Somers D, Hilkin A, Lilli K, Wallace W, Franzwa B, Waterman E, Chan C, McGrath-Chong M, Copland M, Levin A, Sioson L, Cabezon E, Kwan S, Roger D, Lindsay R, Suri R, Champagne J, Bullas R, Garg A, Mazzoato A, Spanner E, Rocco M, Burkart J, Moossavi S, Mauck V, Kaufman T, Pierratos A, Chan W, Regozo K, Kwok S; Frequent Hemodialysis Network (FHN) Trial Group: The effects of frequent nocturnal home hemodialysis: The Frequent Hemodialysis Network Nocturnal Trial. *Kidney Int* 80: 1080–1091, 2011
10. Culleton BF, Walsh M, Klarenbach SW, Mortis G, Scott-Douglas N, Quinn RR, Tonelli M, Donnelly S, Friedrich MG, Kumar A, Mahallati H, Hemmelgarn BR, Manns BJ: Effect of frequent nocturnal hemodialysis vs conventional hemodialysis on left ventricular mass and quality of life: A randomized controlled trial. *JAMA* 298: 1291–1299, 2007
11. Weinhandl ED, Liu J, Gilbertson DT, Arneson TJ, Collins AJ: Survival in daily home hemodialysis and matched thrice-weekly in-center hemodialysis patients. *J Am Soc Nephrol* 23: 895–904, 2012
12. Fewell Z, Davey Smith G, Sterne JA: The impact of residual and unmeasured confounding in epidemiologic studies: A simulation study. *Am J Epidemiol* 166: 646–655, 2007
13. Lawlor DA, Davey Smith G, Kundu D, Bruckdorfer KR, Ebrahim S: Those confounded vitamins: What can we learn from the differences between observational versus randomised trial evidence? *Lancet* 363: 1724–1727, 2004

14. Vandembroucke JP: When are observational studies as credible as randomised trials? *Lancet* 363: 1728–1731, 2004
15. Benson K, Hartz AJ: A comparison of observational studies and randomized, controlled trials. *N Engl J Med* 342: 1878–1886, 2000
16. Concato J, Shah N, Horwitz RJ: Randomized, controlled trials, observational studies, and the hierarchy of research designs. *N Engl J Med* 342: 1887–1892, 2000
17. Ioannidis JPA, Haidich AB, Pappa M, Pantazis N, Kokori SI, Tektonidou MG, Contopoulos-Ioannidis DG, Lau J: Comparison of evidence of treatment effects in randomized and nonrandomized studies. *JAMA* 286: 821–830, 2001

See related article, "Survival in Daily Home Hemodialysis and Matched Thrice-Weekly In-Center Hemodialysis Patients," on pages 895–904.