

- the progression of renal dysfunction and fibrosis in WKY rats with established anti-glomerular basement membrane nephritis. *J Am Soc Nephrol* 17: 674–685, 2006
17. Sharma U, Rhaleb NE, Pokharel S, Harding P, Rasoul S, Peng H, Carretero OA: Novel anti-inflammatory mechanisms of N-Acetyl-Ser-Asp-Lys-Pro in hypertension-induced target organ damage. *Am J Physiol Heart Circ Physiol* 294: H1226–H1232, 2008
 18. Sun Y, Yang F, Yan J, Li Q, Wei Z, Feng H, Wang R, Zhang L, Zhang X: New anti-fibrotic mechanisms of n-acetyl-seryl-aspartyl-lysyl-proline in silicon dioxide-induced silicosis. *Life Sci* 87: 232–239, 2010
 19. Chen YW, Liu BW, Zhang YJ, Chen YW, Dong GF, Ding XD, Xu LM, Pat B, Fan JG, Li DG: Preservation of basal AcSDKP attenuates carbon tetrachloride-induced fibrosis in the rat liver. *J Hepatol* 53: 528–536, 2010
 20. Tang SCW, Leung JCK, Chan LYY, Eddy AA, Lai KN: Angiotensin converting enzyme inhibitor but not angiotensin receptor blockade or statin ameliorates murine adriamycin nephropathy. *Kidney Int* 73: 288–299, 2008
 21. Krupa A, Jenkins R, Luo DD, Lewis A, Phillips A, Fraser D: Loss of MicroRNA-192 promotes fibrogenesis in diabetic nephropathy. *J Am Soc Nephrol* 21: 438–447, 2010
 22. Wang G, Kwan BC, Lai FM, Choi PC, Chow KM, Li PK, Szeto CC: Intrarenal expression of microRNAs in patients with IgA nephropathy. *Lab Invest* 90: 98–103, 2010
 23. Long J, Wang Y, Wang W, Chang BH, Danesh FR: MicroRNA-29c is a signature microRNA under high glucose conditions that targets Sprouty homolog 1, and its in vivo knockdown prevents progression of diabetic nephropathy. *J Biol Chem* 286: 11837–11848, 2011

See related article, “MicroRNA-324-3p Promotes Renal Fibrosis and Is a Target of ACE Inhibition,” on pages 1496–1505.

Aquaporin 2: Not Just for Moving Water

Jeff M. Sands

Renal Division, Departments of Medicine and Physiology, Emory University School of Medicine, Atlanta, Georgia

J Am Soc Nephrol 23: 1443–1444, 2012.
doi: 10.1681/ASN.2012060613

Aquaporin 2 (AQP2) is widely recognized for its role in vasopressin-stimulated water transport across the collecting duct and hence in the production of concentrated urine.^{1,2} AQP2 is primarily expressed in the apical plasma membrane and subapical vesicles of the collecting duct, although it has also been detected in the basolateral plasma membrane.³ In response to vasopressin binding to the V₂-vasopressin receptor, AQP2 is trafficked from the subapical vesicles to the apical plasma membrane; AQP2 is endocytosed and recycled into subapical vesicles when the vasopressin stimulus ends.¹ Water exits collecting duct cells through AQP3 and AQP4, located in

the basolateral plasma membrane, resulting in the transcellular reabsorption of water.⁴

In the current issue of *JASN*, Chen *et al.*⁵ provide evidence for a novel role for AQP2 in promoting cell migration and epithelial morphogenesis. Their rationale for hypothesizing that AQP2 may do more than transport water merits explanation, as it is quite clever. They noted that the phenotype of AQP2-null and -transgenic mice includes the severe urine concentrating defect that one would anticipate, but also includes renal tubular abnormalities and neonatal mortality from renal failure.^{6–8} Mice lacking AQP3 or AQP4 (or AQP1, which is not expressed in the collecting duct) also have a severe urine concentrating defect, but do not have the tubular abnormalities or neonatal mortality.⁹ Thus, they reasoned the phenotype of the AQP2-null mouse was not simply the result of polyuria and a severe urine concentrating defect but must result from another, previously unrecognized, function of AQP2.

Insight into what this novel role may be came from Chen *et al.* identifying a potential integrin-binding site in AQP2, which is not present in other aquaporins. In addition, Tamma *et al.*¹⁰ recently showed that integrin signaling modulates AQP2 trafficking. This suggested to Chen *et al.* that an interaction between AQP2 and an integrin at this site may contribute to the tubular abnormalities seen in AQP2-null mice.

What are integrins? Integrins are a large family of cell surface adhesion receptors that transduce signals coordinately with growth factors and the extracellular matrix.¹¹ Integrins are expressed in the collecting duct and play an important role in kidney development and repair.¹² More specifically, one of the integrins, integrin β 1, plays an important role in collecting duct development and the maintenance of tubular integrity.^{13,14} Thus, Chen *et al.* explored whether the potential integrin-binding site in AQP2 could play a role in the abnormalities found in AQP2-null mice.

In their study, Chen *et al.* confirmed that AQP2-null mice have tubular defects that were apparent as early as postnatal 7 days of age. They also found abnormal subcellular distribution of integrin β 1 in the AQP2-null mice, with integrin β 1 being expressed primarily at the basolateral plasma membrane. These *in vivo* studies are important for establishing a potential physiologic role for an AQP2– β 1 integrin interaction.

To further explore the functional significance of an AQP2– β 1 integrin interaction, Chen *et al.* proceeded to study cultured cells. They showed that AQP2 interacts with integrins through the integrin-binding site in AQP2, Arg-Gly-Asp (RGD), which is the same motif identified by Tamma *et al.*¹⁰ They then observed that AQP2 promotes epithelial cell migration, which is an important mechanism of tissue repair following injury, in both MDCK and LLC-PK1 cells, and that promigration effects requires both AQP2 and β 1 integrin. Finally, they proceeded to show that AQP2 promotes epithelial cell migration by facilitating the turnover of β 1 integrin in the focal adhesions.

In the present study, Chen *et al.* elucidate a novel role for AQP2 in modulating β 1 integrin trafficking and surface expression, and turnover at the focal adhesions. This mechanism

Published online ahead of print. Publication date available at www.jasn.org.

Correspondence: Dr. Jeff M. Sands, Emory University School of Medicine, Renal Division, 1639 Pierce Drive, NE, WMB Room 338, Atlanta, GA 30322. Email: jeff.sands@emory.edu

Copyright © 2012 by the American Society of Nephrology

allows AQP2 to contribute to cell motility and epithelial morphogenesis and may allow AQP2 to play an important role in the development and maintenance of tubular structure. These findings may also explain why AQP2-null mice have abnormal tubule development and neonatal mortality, in contrast to other AQP-null mice. The piecing together of the uniquely abnormal phenotype of AQP2-null mice and the integrin binding site, and then demonstrating the functional consequences of the AQP2- β 1 integrin interaction, provides a significant advance in our understanding of AQP2 biology and suggests a previously unrecognized role for AQP2 in tubule development.

Although AQP2-null mice demonstrate neonatal mortality, there are humans with AQP2 mutations who do survive to adulthood. At first glance, this may suggest that the abnormalities in the AQP2-null mice do not occur in people. However, there is an important difference between AQP2-null mice and humans with AQP2 mutations: the mice completely lack AQP2, whereas the humans have a mutated form of AQP2 that does not transport water. Because the RGD motif that binds β 1 integrin is located in the second extracellular loop of AQP2,^{5,10} one wonders whether the mutant AQP2 proteins present in humans maintain this newly identified role of AQP2 in regulating epithelial cell migration through interaction with β 1 integrin, even if they cannot transport water? It would be interesting to test whether known disease-causing AQP2 mutations interact with β 1 integrin and maintain a promigratory effect on epithelial cell mobility.

ACKNOWLEDGMENTS

This review was supported by NIH Grants R01-DK41707, R01-DK89828, and R21-DK91147.

DISCLOSURES

None.

REFERENCES

- Brown D: The ins and outs of aquaporin-2 trafficking. *Am J Physiol Renal Physiol* 284: F893–F901, 2003
- Sands JM, Layton HE, Fenton RA: Urine concentration and dilution. In: *Brenner and Rector's The Kidney*, 9th Ed., edited by Taal MW, Chertow GM, Marsden PA, Skorecki K, Yu ASL, Brenner BM, Elsevier, Philadelphia, 2011, 326–352
- van Balkom BWM, van Raak M, Breton S, Pastor-Soler N, Bouley R, van der Sluijs P, Brown D, Deen PMT: Hypertonicity is involved in redirecting the aquaporin-2 water channel into the basolateral, instead of the apical, plasma membrane of renal epithelial cells. *J Biol Chem* 278: 1101–1107, 2003
- Nielsen S, Frøkiaer J, Marples D, Kwon TH, Agre P, Knepper MA: Aquaporins in the kidney: From molecules to medicine. *Physiol Rev* 82: 205–244, 2002
- Chen Y, Rice W, Gu Z, Li J, Huang J, Brenner MB, Van Hoek AN, Xiong J, Gundersen GG, Norman JC, Hsu VW, Fenton RA, Brown D, Lu HAJ: Aquaporin 2 promotes cell migration and epithelial morphogenesis. *J Am Soc Nephrol* 23: 1506–1517, 2012
- Rojek A, Führtbauer EM, Kwon TH, Frøkiaer J, Nielsen S: Severe urinary concentrating defect in renal collecting duct-selective AQP2 conditional-knockout mice. *Proc Natl Acad Sci U S A* 103: 6037–6042, 2006
- Yang B, Gillespie A, Carlson EJ, Epstein CJ, Verkman AS: Neonatal mortality in an aquaporin-2 knock-in mouse model of recessive nephrogenic diabetes insipidus. *J Biol Chem* 276: 2775–2779, 2001
- Yang BX, Zhao D, Qian LM, Verkman AS: Mouse model of inducible nephrogenic diabetes insipidus produced by floxed aquaporin-2 gene deletion. *Am J Physiol Renal Physiol* 291: F465–F472, 2006
- Fenton RA, Knepper MA: Mouse models and the urinary concentrating mechanism in the new millennium. *Physiol Rev* 87: 1083–1112, 2007
- Tamma G, Lasorsa D, Ranieri M, Mastrofrancesco L, Valenti G, Svelto M: Integrin signaling modulates AQP2 trafficking via Arg-Gly-Asp (RGD) motif. *Cell Physiol Biochem* 27: 739–748, 2011
- Arcangeli A, Becchetti A: Complex functional interaction between integrin receptors and ion channels. *Trends Cell Biol* 16: 631–639, 2006
- Kreidberg JA, Symons JM: Integrins in kidney development, function, and disease. *Am J Physiol Renal Physiol* 279: F233–F242, 2000
- Wu W, Kitamura S, Truong DM, Rieg T, Vallon V, Sakurai H, Bush KT, Vera DR, Ross RS, Nigam SK: Beta1-integrin is required for kidney collecting duct morphogenesis and maintenance of renal function. *Am J Physiol Renal Physiol* 297: F210–F217, 2009
- Zhang X, Mernaugh G, Yang DH, Gewin L, Srichai MB, Harris RC, Iturregui JM, Nelson RD, Kohan DE, Abrahamson D, Fässler R, Yurchenco P, Pozzi A, Zent R: beta1 integrin is necessary for ureteric bud branching morphogenesis and maintenance of collecting duct structural integrity. *Development* 136: 3357–3366, 2009

See related article, "Aquaporin 2 Promotes Cell Migration and Epithelial Morphogenesis," on pages 1506–1517.

Lifetime Risk of ESRD: A Meaningful Concept?

Paul Eggers

Kidney and Urology Epidemiology, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, Maryland

J Am Soc Nephrol 23: 1444–1446, 2012.
doi: 10.1681/ASN.2012070733

Turin and colleagues¹ present some fascinating estimates of both the lifetime and short-term risks of ESRD in this issue of *JASN*, exploiting the excellent epidemiologic database in Alberta, Canada. Based on a total population of nearly 2.9 million persons with nearly 26 million person-years of follow-up, they have been able to calculate actual lifetime event rates of ESRD (2.6% and 1.8% for men and women, respectively) rather than the synthetic estimates based on observed incidence rates applied to hypothetical populations.²

The advantage of this approach is not so much in increased accuracy of the estimate as in the ability to array the predictions

Published online ahead of print. Publication date available at www.jasn.org.

Correspondence: Dr. Paul Eggers, Kidney and Urology Epidemiology, NIDDK, Room 615, 6707 Democracy Boulevard, Bethesda, MD 20892. Email: eggersp@extra.niddk.nih.gov

Copyright © 2012 by the American Society of Nephrology