

- Kusek JW, Keane MG, Wolf M: FGF23 induces left ventricular hypertrophy. *J Clin Invest* 121: 4393–4408, 2011
13. Doi S, Zou Y, Togao O, Pastor JV, John GB, Wang L, Shiizaki K, Gotschall R, Schiavi S, Yorioka N, Takahashi M, Boothman DA, Kuro-o M: Klotho inhibits transforming growth factor- β 1 (TGF- β 1) signaling and suppresses renal fibrosis and cancer metastasis in mice. *J Biol Chem* 286: 8655–8665, 2011
14. Guan X, Nie L, He T, Yang K, Xiao T, Wang S, Huang Y, Zhang J, Wang J, Sharma K, Liu Y, Zhao J: Klotho suppresses renal tubulo-interstitial fibrosis by controlling basic fibroblast growth factor-2 signalling. *J Pathol* 234: 560–572, 2014
15. Zhou L, Li Y, Hao S, Zhou D, Tan RJ, Nie J, Hou FF, Kahn M, Liu Y: Multiple genes of the renin-angiotensin system are novel targets of Wnt/ β -catenin signaling. *J Am Soc Nephrol* 26: 107–120, 2015

See related article, “Klotho Protects Against Indoxyl Sulphate-Induced Myocardial Hypertrophy,” on pages 2434–2446.

Not All Deaths in CKD Are from a Broken Heart

Germaine Wong^{*†‡} and Amit X. Garg^{§||}

^{*}Sydney School of Public Health, University of Sydney, Sydney, New South Wales, Australia; [†]Centre for Transplant and Renal Research, Westmead Hospital, Westmead, New South Wales, Australia; [‡]Centre for Kidney Research, Kids Research Institute, The Children’s Hospital at Westmead, Westmead, New South Wales, Australia; [§]Institute for Clinical Evaluative Sciences, Ontario, Canada; and ^{||}Division of Nephrology, Department of Medicine, Western University, London, Canada

J Am Soc Nephrol 26: 2307–2308, 2015.
doi: 10.1681/ASN.2015050531

CKD is common, but outcomes remain poor, with a high incidence of death as eGFR declines.¹ Two studies published in this issue of *JASN* examine the association between reduced kidney function and specific causes of death.^{2,3} Navaneethan *et al.*² used electronic health records and a linked mortality registry from the state of Ohio in the United States to study adults with two eGFR values <60 ml/min per 1.73 m². Thompson *et al.*³ used administrative health care data and linked laboratory information from the province of Alberta in Canada to study all adult deaths stratified by recent eGFR and urine protein results. Reliably ascertaining an individual’s cause of death is notoriously difficult, even in prospective cohort studies with central adjudication. In both of these retrospective analyses, administrative personnel recorded the cause of death in routine care using the International Classification of Diseases 10th Revision system. With this method of

ascertainment, we expect that many deaths were classified incorrectly. Reasons for misclassification include scenarios with multiple causes of death, ill-defined causes of death, and a lack of autopsy data. Nonetheless, these studies by Navaneethan *et al.*² and Thompson *et al.*³ provide several important messages for our consideration.

Both studies confirm that a greater proportion of deaths is attributable to cardiovascular diseases as eGFR declines. Some would say this association between low eGFR and a higher risk of death “from a broken heart” is already well appreciated by the nephrology community.⁴ As might be expected, in the Alberta study, after adjustment for age and sex, 33% of deaths were attributed to cardiovascular disease when the eGFR was 45–59 ml/min per 1.73 m², and 40% of deaths were attributed to cardiovascular disease when the eGFR was 15–29 ml/min per 1.73 m².³ Similarly, in the Ohio study, the 3-year probability of mortality from cardiovascular disease increased in a graded manner as eGFR declined (from approximately 3% when the eGFR was 60 ml/min per 1.73 m² to 7.5% when the eGFR was 10 ml/min per 1.73 m²).² The Alberta study provides new insights into the nature of these cardiovascular deaths. As eGFR declined, a greater proportion of people died from cardiac failure and valvular disease rather than ischemic heart disease. This finding is consistent with prior data of chronic fluid overload coupled with anemia and chronic hypertension producing a high cardiac output state, leading to mechanical trauma.⁵ Valvular calcification is also prominent in CKD, where it is attributed to chronic inflammation and impairments of calcium phosphate metabolism.⁶

There are other prominent causes of death in CKD beyond a broken heart. In the Alberta study, infection accounted for 3% of deaths in those with an eGFR of 45–59 ml/min per 1.73 m² and 5% of deaths in those with an eGFR of 15–29 ml/min per 1.73 m². A similar graded relationship between CKD stage and severe infection requiring hospitalization was also observed in a recent meta-analysis.⁷ Better infection control may prevent some of these deaths. For example, in the study by Navaneethan *et al.*,² approximately 10% of all CKD deaths were attributed to diabetic-related infections. Perhaps better diabetic foot care will prevent some deaths.

In the Ohio study, the 3-year probability of mortality from cancer was approximately 2.5%, and this percentage remained unchanged through all stages of eGFR. Similar findings were observed in the Alberta cohort. Large population-based cohort studies suggest a graded association between reduced eGFR and cancer risk.^{8,9} However, the direction and the causality of this association require clarification. Although in the Ohio study, approximately 20% of those with CKD had cancer, it is unclear whether these cancers developed before or after the onset of CKD. Cancers, such as multiple myeloma and lymphoma, are established risk factors for CKD, whereas 1 in 10 patients has CKD in the setting of solid-organ malignancies.¹⁰ In other analyses restricted to those with cancer, CKD was associated with at least a 2-fold higher risk of death from cancer.¹¹

Where do we go from here? These two studies shed new light on the association between reduced eGFR and cause of death.^{2,3}

Published online ahead of print. Publication date available at www.jasn.org.

Correspondence: Dr. Germaine Wong, Sydney School of Public Health, University of Sydney, Sydney, NSW, Australia 2006. Email: germaine.wong@health.nsw.gov.au

Copyright © 2015 by the American Society of Nephrology

While we continue to search for therapies to prevent cardiovascular deaths in CKD, we also need to focus on strategies to prevent nonvascular deaths from infection and cancer. For example, the benefit of screening to reduce cancer-related deaths in CKD remains largely unknown.¹² These studies serve as a call to bridge the gap between pathogenic understanding, experimental data, clinical testing, and high-quality care. We urgently need proven strategies to prevent deaths in CKD. Sadly, the results of intervention trials in CKD (for example, of antiplatelet and lipid-lowering therapies to prevent major cardiovascular events) have been disappointing or modest at best.^{13,14} Let us stop the heart break by working together on large-scale, practice-changing research. Finally, not all deaths should be prevented at the expense of quality of life, and we need to listen carefully to our patients and their families to ensure that we provide care according to their values and preferences.

ACKNOWLEDGMENTS

We thank Drs. Jonathan Craig and Jessica Sontrop for their advice. A.X.G. was supported by the Dr. Adam Linton Chair in Kidney Health Analytics.

DISCLOSURES

None.

REFERENCES

- Coresh J, Byrd-Holt D, Astor BC, Briggs JP, Eggers PW, Lacher DA, Hostetter TH: Chronic kidney disease awareness, prevalence, and trends among U.S. adults, 1999 to 2000. *J Am Soc Nephrol* 16: 180–188, 2005
- Navaneethan S, Schold J, Arrigain S, Jolly S, Nally J: Cause-specific deaths in non-dialysis dependent chronic kidney disease. *J Am Soc Nephrol* 26: 2512–2520, 2015
- Thompson S, James M, Wiebe N, Hemmelgarn B, Manns B, Klarenbach S, Tonelli M; Alberta Kidney Disease Network: Cause of death in patients with reduced kidney function. *J Am Soc Nephrol* 26: 2504–2511, 2015
- Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY: Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med* 351: 1296–1305, 2004
- Abreo G, Lenihan DJ, Nguyen P, Runge MS: High-output heart failure resulting from a remote traumatic aorto-caval fistula: Diagnosis by echocardiography. *Clin Cardiol* 23: 304–306, 2000
- Rouached M, El Kadiri Boutchich S, Al Rifai AM, Garabédian M, Fournier A: Prevalence of abnormal serum vitamin D, PTH, calcium, and phosphorus in patients with chronic kidney disease: Results of the study to evaluate early kidney disease. *Kidney Int* 74: 389–390, 2008
- McDonald HI, Thomas SL, Nitsch D: Chronic kidney disease as a risk factor for acute community-acquired infections in high-income countries: A systematic review. *BMJ Open* 4: e004100, 2014
- Lowrance WT, Ordoñez J, Udaltsova N, Russo P, Go AS: CKD and the risk of incident cancer. *J Am Soc Nephrol* 25: 2327–2334, 2014
- Wong G, Hayen A, Chapman JR, Webster AC, Wang JJ, Mitchell P, Craig JC: Association of CKD and cancer risk in older people. *J Am Soc Nephrol* 20: 1341–1350, 2009
- Launay-Vacher V, Oudard S, Janus N, Gligorov J, Pourrat X, Rixe O, Morere JF, Beuzeboc P, Deray G; Renal Insufficiency and Cancer Medications (IRMA) Study Group: Prevalence of Renal Insufficiency in cancer patients and implications for anticancer drug management: The renal insufficiency and anticancer medications (IRMA) study. *Cancer* 110: 1376–1384, 2007
- Iff S, Craig JC, Turner R, Chapman JR, Wang JJ, Mitchell P, Wong G: Reduced estimated GFR and cancer mortality. *Am J Kidney Dis* 63: 23–30, 2014
- Wong G, Chapman JR, Craig JC: Cancer screening in renal transplant recipients: What is the evidence? *Clin J Am Soc Nephrol* 3[Suppl 2]: S87–S100, 2008
- Baigent C, Landray MJ, Reith C, Emberson J, Wheeler DC, Tomson C, Wanner C, Krane V, Cass A, Craig J, Neal B, Jiang L, Hooi LS, Levin A, Agodoa L, Gaziano M, Kasiske B, Walker R, Massy ZA, Feldt-Rasmussen B, Krairittichai U, Ophascharoensuk V, Fellström B, Holdaas H, Tesar V, Wiecek A, Grobbee D, de Zeeuw D, Grönhagen-Riska C, Dasgupta T, Lewis D, Herrington W, Maffham M, Majoni W, Wallendszus K, Grimm R, Pedersen T, Tobert J, Armitage J, Baxter A, Bray C, Chen Y, Chen Z, Hill M, Knott C, Parish S, Simpson D, Sleight P, Young A, Collins R; SHARP Investigators: The effects of lowering LDL cholesterol with simvastatin plus ezetimibe in patients with chronic kidney disease (Study of Heart and Renal Protection): A randomised placebo-controlled trial. *Lancet* 377: 2181–2192, 2011
- Palmer SC, Di Micco L, Razavian M, Craig JC, Perkovic V, Pellegrini F, Jardine MJ, Webster AC, Zoungas S, Strippoli GF: Antiplatelet agents for chronic kidney disease. *Cochrane Database Systematic Review* 2: CD008834, 2013

See related articles, “Cause of Death in Patients with Reduced Kidney Function,” and “Cause-Specific Deaths in Non-Dialysis-Dependent CKD,” on pages 2504–2511 and 2512–2520, respectively.

Anti-phospholipase A₂ Receptor Antibody and Immunosuppression in Membranous Nephropathy: More Evidence for Pathogenicity of Anti-phospholipase A₂ Receptor Autoantibodies

Paul E.C. Brenchley

Renal Research Labs, Manchester Royal Infirmary and Institute of Cardiovascular Sciences, University of Manchester, Manchester, United Kingdom

J Am Soc Nephrol 26: 2308–2311, 2015.
doi: 10.1681/ASN.2015020181

In the 6 years since the discovery of anti-phospholipase A₂ receptor (anti-PLA₂R) antibodies in the majority of patients

Published online ahead of print. Publication date available at www.jasn.org.

Correspondence: Prof. Paul E.C. Brenchley, Renal Research Labs, Manchester Royal Infirmary and Institute of Cardiovascular Sciences, University of Manchester, Oxford Road, Manchester M13 9WL, UK. Email: paul.brenchley@manchester.ac.uk

Copyright © 2015 by the American Society of Nephrology