

Fibroblast Growth Factor-23 May Follow Cardiovascular Disease Rather than Causing It in Chronic Kidney Disease

We read the article “Fibroblast growth factor-23 and risks of cardiovascular and noncardiovascular diseases: A meta-analysis” by Marthi *et al.*¹ with great interest. The study provided an important demonstration that serum higher fibroblast growth factor-23 (FGF-23) level was positively associated with cardiovascular and noncardiovascular outcomes in populations with or without CKD.¹ In agreement with Marthi *et al.*,¹ our previous meta-analysis also pointed out that there might be predictive effects of FGF-23 on cardiovascular diseases (CVDs; including myocardial infarction, atrial fibrillation, myocardial ischemia, heart failure, and stroke) and mortality in patients with CKD.²

However, the absence of an exposure-response relationship and similarly sized associations between FGF-23 and different outcomes may not be strong enough to suggest a noncausal relationship between FGF-23 and CVD. Indeed, interactions between FGF-23 and CVD are complicated and remain inconclusive. Experimental studies have revealed that FGF-23 could induce left ventricular hypertrophy.³ Both FGF-23 overexpression and *Klotho* deficiency showed cardiac remodeling effects in CKD.⁴ However, Slavic *et al.*⁵ found that genetic ablation of *FGF-23* or *Klotho* did not modulate heart hypertrophy, indicating other mediators as the real culprits. Recently, left ventricular hypertrophy in transgenic mice was found to elevate myocardial and serum levels of FGF-23.³ Taken together, the mutual promotion between FGF-23 and CVD indicates a complex interaction between mineral metabolism and the cardiovascular system in CKD. FGF-23 may follow CVD rather than causing it in CKD.

In conclusion, the finding of this meta-analysis¹ is very important, because it brings us a deep and broad view of FGF-23 in CVD and mortality in patients with or without kidney disease. More mechanisms between FGF-23 and CVD will need to be studied to better understand the effect of FGF-23 on cardiovascular health in patients with CKD.

Published online ahead of print. Publication date available at www.jasn.org.

Correspondence: Dr. Cheng Xue, Division of Nephrology, Changzheng Hospital, Second Military Medical University, Fengyang Road 415, Shanghai 200000, China. Email: xuecheng@smmu.edu.cn

Copyright © 2018 by the American Society of Nephrology

ACKNOWLEDGMENTS

The work was supported by National Natural Science Foundation of China grant 81700579.

DISCLOSURES

None.

REFERENCES

1. Marthi A, Donovan K, Haynes R, Wheeler DC, Baigent C, Rooney CM, et al.: Fibroblast growth factor-23 and risks of cardiovascular and noncardiovascular diseases: A meta-analysis. *J Am Soc Nephrol* 29: 579–590, 2018
2. Xue C, Yang B, Zhou C, Dai B, Liu Y, Mao Z, et al.: Fibroblast growth factor 23 predicts all-cause mortality in a dose-response fashion in pre-dialysis patients with chronic kidney disease. *Am J Nephrol* 45: 149–159, 2017
3. Matsui I, Oka T, Kusunoki Y, Mori D, Hashimoto N, Matsumoto A, et al.: Cardiac hypertrophy elevates serum levels of fibroblast growth factor 23. *Kidney Int* 94: 60–71, 2018
4. Hu MC, Shi M, Cho HJ, Adams-Huet B, Paek J, Hill K, et al.: *Klotho* and phosphate are modulators of pathologic uremic cardiac remodeling. *J Am Soc Nephrol* 26: 1290–1302, 2015
5. Slavic S, Ford K, Modert M, Becirovic A, Handschuh S, Baierl A, et al.: Genetic ablation of *Fgf23* or *Klotho* does not modulate experimental heart hypertrophy induced by pressure overload. *Sci Rep* 7: 11298, 2017

See related Letters to the Editor, “Fibroblast Growth Factor-23 Is Not a Single Bystander in Chronic Kidney Disease Mortality,” and “Authors’ Reply,” on pages 2601 and 2602–2603, respectively.

Chenchen Zhou^{1,2}, Changlin Mei², Bing Dai², and Cheng Xue²

¹Department of Nephrology, Yueyang Hospital of Integrated Traditional Chinese and Western Medicine, Shanghai University of Traditional Chinese Medicine, Shanghai, China; and

²Division of Nephrology, Changzheng Hospital, Second Military Medical University, Shanghai, China

J Am Soc Nephrol 29: 2602, 2018.

doi: <https://doi.org/10.1681/ASN.2018050517>

Authors’ Reply

The published observational data show similarly sized epidemiologic associations between increased fibroblast growth factor-23 (FGF-23) concentration and risk of a range of cardio-

Published online ahead of print. Publication date available at www.jasn.org.

Correspondence: Dr. William Herrington, Nuffield Department of Population Health, University of Oxford, Richard Doll Building, OX3 7LF Oxford, United Kingdom. Email: will.herrington@ndph.ox.ac.uk

Copyright © 2018 by the American Society of Nephrology