

Joseph Chilcot¹ and Ken Farrington^{2,3}¹Psychology Department, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom;²Renal Unit, Lister Hospital, Stevenage, United Kingdom; and³Postgraduate Medicine, The University of Hertfordshire, Hatfield, United Kingdom

JASN 30: 1339–1340, 2019.

doi: <https://doi.org/10.1681/ASN.2019030308>

Authors' Reply

We thank Chilcot and Farrington¹ for their interest in our paper² and thoughtful comments surrounding the treatment of depression in hemodialysis patients. We agree with the letter writers that our observational study was not designed to evaluate the effect of antidepressant pharmacotherapy (versus no treatment) on mortality among individuals with hemodialysis-dependent ESRD. Under our active comparator new-user design, we could only conclude that the initiation of a higher (citalopram and escitalopram) versus lower (fluoxetine, fluvoxamine, paroxetine, and sertraline) QT-prolonging potential selective serotonin reuptake inhibitor (SSRI) was associated with a higher risk of sudden cardiac death.² Our study was designed to reflect a clinician's decision to prescribe an SSRI with higher versus lower QT-prolonging potential to a hemodialysis patient (*i.e.*, a treatment choice encountered in real-world practice).^{2–4} The study provides population-specific safety information that clinicians can consider when prescribing SSRI therapy to hemodialysis patients.

ACKNOWLEDGMENTS

Dr. Assimon and Dr. Flythe are supported by grant R03 HS026801 from the Agency for Healthcare Research and Quality. Dr. Flythe is supported by grant K23 DK109401 from the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health.

DISCLOSURES

Dr. Assimon and Dr. Flythe have received investigator-initiated research funding from the Renal Research Institute, a subsidiary of Fresenius Medical Care, North America. In the last 2 years, Dr. Flythe has received speaking honoraria from American Renal Associates; the American Society of Nephrology; Dialysis Clinic, Inc.; the National Kidney Foundation; and multiple universities. Dr. Flythe is on the medical advisory board of NxStage Medical, Inc. and has received consulting fees from Fresenius Medical Care, North America.

REFERENCES

1. Chilcot J, Farrington K: The case for selective withdrawal of antidepressants in patients with advanced kidney disease. *J Am Soc Nephrol* 30: 1339, 2019
2. Assimon MM, Brookhart MA, Flythe JE: Comparative cardiac safety of selective serotonin reuptake inhibitors among individuals receiving maintenance hemodialysis. *J Am Soc Nephrol* 30: 611–623, 2019
3. Velentgas P, Dreyer NA, Nourjah P, Smith SR, Torchia MM, editors: *Developing a Protocol for Observational Comparative Effectiveness Research: A User's Guide*, Rockville, MD, Agency for Healthcare Research and Quality, 2013
4. Brookhart MA: Counterpoint: The treatment decision design. *Am J Epidemiol* 182: 840–845, 2015

Magdalene M. Assimon¹ and Jennifer E. Flythe^{1,2}¹University of North Carolina Kidney Center, Division of Nephrology and Hypertension, Department of Medicine, UNC School of Medicine, Chapel Hill, North Carolina; and²Cecil G. Sheps Center for Health Services Research, University of North Carolina, Chapel Hill, North Carolina

JASN 30: 1340, 2019.

doi: <https://doi.org/10.1681/ASN.2019040431>

Published online ahead of print. Publication date available at www.jasn.org.

Correspondence: Dr. Magdalene M. Assimon, University of North Carolina Kidney Center, Division of Nephrology and Hypertension, Department of Medicine, University of North Carolina School of Medicine, 7024 Burnett-Womack CB #7155, Chapel Hill, NC 27599-7155. Email: massimon@med.unc.edu

Copyright © 2019 by the American Society of Nephrology