

**EMBARGOED FOR RELEASE until JANUARY 20, 2011 – 5:00 PM (EST)**

**ASN Contact:** Shari Leventhal • 202-416-0658 (p) • [sleventhal@asn-online.org](mailto:sleventhal@asn-online.org)

## **GENE'S MUTATIONS PLAY A MAJOR ROLE IN ONE CAUSE OF KIDNEY FAILURE**

**Screening may be warranted when the disease runs in families**

**Washington, DC (January 14, 2011)** — Mutations in a gene called *INF2* are by far the most common cause of a dominantly inherited condition that leads to kidney failure, according to a study appearing in an upcoming issue of the *Journal of the American Society Nephrology* (JASN). The results may help with screening, prevention, and therapy.

Focal segmental glomerulosclerosis (FSGS) attacks the kidney's filtering system and causes serious scarring. Approximately 20,000 persons are currently living with kidney failure due to FSGS in the United States, with an associated annual cost of more than \$3 billion. In addition, studies have shown that the incidence of FSGS is increasing. FSGS is a common cause of kidney failure in adults and the second leading cause in children. During the last decade, researchers have identified several genes that are mutated in patients with hereditary FSGS and have gained a better understanding of the mechanisms behind the disease's development. These advances are useful for genetic counseling and for developing strategies aimed at prevention and treatment.

One of the genes mutated in patients with hereditary, autosomal dominant FSGS is the *INF2* gene, initially discovered by Elizabeth J. Brown and Martin R. Pollak (Harvard Medical School, Boston)<sup>1</sup>, which produces a protein that helps maintain the structure of specialized kidney cells called podocytes. Corinne Antignac MD, PhD, Olivia Boyer, MD (Hôpital Necker-Enfants Malades and Université Paris Descartes, in Paris, France) and their colleagues conducted a study to confirm *INF2*'s importance in the development of FSGS and to better determine the prevalence of *INF2* mutations in a worldwide group of pediatric and adult patients.

The investigators screened 54 families (78 patients) with a history of autosomal dominant FSGS and detected mutations in the *INF2* gene in 17% of them. The mutations were located in one particular region of the gene that corresponds to a part of the *INF2* protein that interacts with podocyte proteins. This information might shed light on the mechanism behind *INF2*'s involvement in the development of FSGS and could be helpful as researchers design drugs to prevent or treat the disease. One of these mutations appeared in only one of 84 patients with sporadic (nonhereditary) FSGS.

“*INF2* is a major gene of autosomal dominant FSGS. Screening for *INF2* mutations needs to be strongly considered in patients with an autosomal dominant familial history of FSGS,” the authors concluded.

Study co-authors include Geneviève Benoit, MD, Olivier Gribouval, Fabien Nevo, Marie-Josèphe Tête, MD, Jean-Pierre Grunfeld, MD, Christophe Legendre, MD, Dominique Joly, MD, PhD, Marie-Claire Gubler, MD, Géraldine Mollet, PhD (Hôpital Necker-Enfants Malades); Jacques Dantal, MD, PhD (CHU Hôtel Dieu, in Nantes, France); Brigitte Gilbert-Dussardier, MD (CHU La Milétrie, in Poitiers, France); Guy Touchard, MD (Hopital Jean Bernard, in Poitiers, France); Alexandre Karras, MD (Hôpital Européen Georges Pompidou, in Paris, France); Claire Presne, MD (CHU d’Amiens, in Amiens, France); Philippe Rieu, MD (Hôpital du Kremlin-Bicêtre, France); Nabil Mohsin, MD (Royal Hospital, in Muscat, Oman); Isabelle Broutin, PhD (CNRS, in Paris, France); Thierry Hannedouche, MD, PhD (Hôpitaux Universitaires de Strasbourg, France); and Valérie Moal, MD (Hôpital de la Conception, in Marseille, France).

Disclosures: The authors reported no financial disclosures.

The article, entitled “Mutations in *INF2* Are a Major Cause of Autosomal Dominant Focal Segmental Glomerulosclerosis,” will appear online at <http://jasn.asnjournals.org/> on January 20, 2011, doi 10.1681/ASN.2010050518.

*The content of this article does not reflect the views or opinions of The American Society of Nephrology (ASN). Responsibility for the information and views expressed therein lies entirely with the author(s). ASN does not offer medical advice. All content in ASN publications is for informational purposes only, and is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. This content should not be used during a medical emergency or for the diagnosis or treatment of any medical condition. Please consult your doctor or other qualified health care provider if you have any questions about a medical condition, or before taking any drug, changing your diet or commencing or discontinuing any course of treatment. Do not ignore or delay obtaining professional medical advice because of information accessed through ASN. Call 911 or your doctor for all medical emergencies.*

*Founded in 1966 and comprised of more than 12,000 members, the American Society of Nephrology (ASN) leads the fight against kidney disease by educating health professionals, sharing new knowledge, advancing research, and advocating the highest quality care for patients.*

# # #

---

<sup>i</sup> Brown EJ, Schlöndorff JS, Becker DJ, Tsukaguchi H, Tonna SJ, Uscinski AL, Higgs HN, Henderson JM, Pollak MR. Mutations in the formin gene *INF2* cause focal segmental glomerulosclerosis. Renal Division, Children's Hospital, Boston, Massachusetts, USA. *Nat Genet.* 2010 Jan;42(1):72-6. Epub 2009 Dec 20. Erratum in: *Nat Genet.* 2010 Apr;42(4):361. Tonna, Stephen J.