

Table S1. Comparison of characteristics in antibody positive and negative patients without a PCR test

Variable		No swab, Ab- N=191 (%)	No swab, Ab+ N=44 (%)	p value
Gender	Female	67 (35.0)	17 (38.6)	0.73
	Male	124 (64.9)	27 (61.3)	
Age	Years (median)	69 (59-78)	66 (58-75)	0.25
Ethnicity	Black	22 (11.5)	7 (15.9)	0.84
	Caucasian	51 (26.4)	11 (25.0)	
	Indoasian	78 (40.8)	19 (43.2)	
	Other	40 (20.9)	7 (15.9)	
Cause of ESKD	APKD	12 (6.3)	4 (9.0)	0.12
	Diabetic nephropathy	74 (38.7)	11 (25.0)	
	Glomerulonephritis	35 (18.3)	8 (18.1)	
	Other	30 (15.7)	5 (11.3)	
	Unknown	34 (17.8)	13 (29.5)	
	Urological	6 (3.1)	3 (6.8)	
Time at ESKD	Years (median)	2.2 (0.92-4.2)	1.6 (0.94-4.2)	0.82
Unit	A	35 (18.3)	8 (18.2)	1.00
	B	156 (81.6)	38 (86.3)	
Lymphocyte Nadir	Median (10⁹/l)	1.2 (0.9-1.6)	1.1 (0.9-1.5)	0.52
CRP Peak	Median (mg/l)	7.7 (2.9-20.7)	7.2 (2.6-12.0)	0.87
Ferritin Peak	Median (µg/l)	391 (294-545)	431 (324-532)	0.57
Immunosuppressed	Yes	36 (18.8)	7 (15.9)	0.83